Financial Burdens of Cancer Treatment: A Systematic Review of Risk Factors and Outcomes

Grace L. Smith, MD, PhD, MPH^{a,b}; Maria A. Lopez-Olivo, MD, PhD^c; Pragati G. Advani, MD, DrPH^d; Matthew S. Ning, MD^a; Yimin Geng, MS, MSLIS^e; Sharon H. Giordano, MD, MPH^{a,f}; and Robert J. Volk, PhD^b

ABSTRACT

Background: Patients with cancer experience financial toxicity from the costs of treatment, as well as material and psychologic stress related to this burden. A synthesized understanding of predictors and outcomes of the financial burdens associated with cancer care is needed to underpin strategic responses in oncology care. This study systematically reviewed risk factors and outcomes associated with financial burdens related to cancer treatment. **Methods:** MEDLINE, Embase, PubMed, PsychINFO, and the Cochrane Library were searched from study inception through June 2018, and reference lists were scanned from studies of patient-level predictors and outcomes of financial burdens in US patients with cancer (aged ≥ 18 years). Two reviewers conducted screening, abstraction, and quality assessment. Variables associated with financial burdens were synthesized. When possible, pooled estimates of associations were calculated using random-effects models. **Results:** A total of 74 observational studies of financial burdens in 598,751 patients with cancer were identified, among which 49% of patients reported material or psychologic financial burdens (95% CI, 41%–56%). Socioeconomic predictors of worse financial burdens with treatment were lack of health insurance, lower income, unemployment, and younger age at cancer diagnosis. Compared with patients with health insurance, those who were uninsured demonstrated twice the odds of financial burdens (pooled odds ratio [OR], 2.09; 95% CI, 1.33–3.30). Financial burdens were most severe early in cancer treatment, did not differ by disease site, and were associated with worse health-related quality of life (HRQoL) and nearly twice the odds of cancer medication nonadherence (pooled OR, 1.70; 95% CI, 1.13–2.56). Only a single study demonstrated an association with increased mortality. Studies assessing the comparative effectiveness of interventions to mitigate financial burdens in patients with cancer were lacking. **Conclusions:** Evidence showed that financial burdens are common, disproportionately impacting younger and socioeconomically disadvantaged patients with cancer, across disease sites, and are associated with worse treatment adherence and HRQoL. Available evidence helped identify vulnerable patients needing oncology provider engagement and response, but evidence is critically needed on the effectiveness of interventions designed to mitigate financial burden and impact.

J Natl Compr Canc Netw 2019;17(10):1184–1192
doi: 10.6004/jnccn.2019.7305

^aDepartment of Radiation Oncology, ^bDepartment of Health Services Research, and ^cDepartment of General Internal Medicine, The University of Texas MD Anderson Cancer Center, Houston, Texas; ^dDivision of Cancer Epidemiology and Genetics, National Institutes of Health, National Cancer Institute, Bethesda, Maryland; and ^eResearch Medical Library, and ^fDepartment of Breast Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, Texas.

Background

Although the burden of cancer costs on the US health-care system is well-recognized,^{1,2} more recently, the personal “financial toxicity” associated with the cost of care has also been identified as a distinct consequence of cancer treatment.^{3–13} A recent systematic review that sought to define the prevalence of the types of financial hardships among patients with cancer showed that 40% to 50% of US patients experienced material or psychologic stressors related to the cost of treatment.¹³ The frequency of financial burdens among patients with cancer could also increase, especially for those facing escalating cancer therapy costs and barriers to insurance coverage.^{2,12,14,15}

As a result, cancer’s overall financial burden—including the direct costs of care, the indirect costs (eg, lost work, income, and savings), and the distress resulting from these financial stressors of cancer care—is now generally acknowledged as a priority concern faced by US patients as well as the oncology providers caring for them. Yet an effective response to this problem remains lacking, with approximately one-quarter to one-half of patients reporting inadequate oncology provider engagement with their financial burdens.^{16,17} To inform effective response and intervention in vulnerable patients,¹⁸ a comprehensive synthesis of the risk factors for and outcomes of financial burdens is needed. We therefore conducted a systematic review of these risk factors and outcomes across studies of US patients with cancer to identify the types of burdens in vulnerable patient subgroups and the consequent health impact.

Methods

Protocol and Registration

This protocol is registered in PROSPERO (CRD42017054521). Results are reported per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement.¹⁹

See JNCCN.org for supplemental online content.

Eligibility Criteria

Studies were searched that contained primary data analyses examining financial burdens among patients with cancer and the associations of burdens with socioeconomic, demographic, clinical, intervention, and outcome variables. Based on existing conceptual frameworks,^{12,20} studies of both material and psychological financial burdens in patients with cancer were searched. We included English-language, peer-reviewed, full-text studies conducted in US adults (aged ≥ 18 years) with cancers of any disease site and stage, and in any part of the cancer disease trajectory. Exclusion criteria were as follows: non-US studies (given major differences in health insurance and employment among countries), no original data analysis, and no patient-level financial burdens reported.

Search, Study Selection, and Data Collection

MEDLINE, Embase, PsychINFO, and the Cochrane Library were searched from study inception until June 2018, guided by a research librarian (Y. Geng), and reference lists of relevant reviews were manually examined. Search terms and strategy are delineated in eFigure 1 and eAppendix 1, available with this article at JNCCN.org. Two reviewers (G.L. Smith, P.G. Advani) performed title and abstract screening, full-text review, and consensus resolution of interrater discrepancies for inclusion (kappa, 0.79). Two reviewers for each study (G.L. Smith, P.G. Advani, M.S. Ning) abstracted and cross-checked study characteristics, including design, eligibility, and definition or measure of financial burdens directly reported from the text. Discrepancies were resolved through consensus.

Financial Burden, Risk Factor, and Outcome Measures

We extracted financial burden variables from each study, including measures of direct material burdens (eg, higher out-of-pocket [OOP] costs), indirect burdens (eg, lost savings and increased debt), and psychological burdens (eg, financial worries or distress). Key terms were health expenditures, healthcare costs, cost of illness, financing, fees, charges, bankruptcy, insolvency, economic hardship, economic burden, economic stress or distress, economic difficulty, OOP costs, costs, expenses, and spending. A complete list of search terms is included in supplemental eFigure 1. We extracted reported associations between independent variables (eg, risk factors or interventions) and financial burden variables, and associations between financial burden variables and outcome variables, statistical tests of associations, and magnitude/significance of effects (odds ratios [ORs], risk ratios, hazard ratios [HRs], and 95% CI).²¹

Risk of Bias Assessment

Reviewers independently ascertained the risk of bias using the NIH Quality Assessment Tool for Observational

Cohort and Cross-Sectional Studies (supplemental eTable 1),²² including overall study quality (ie, individual study rated as “good” if it met all applicable quality assessment criteria, “fair” if it met most of the criteria, and “poor” if it met a minority of the criteria). Strength of evidence was evaluated using Grading of Recommendations Assessment, Development, and Evaluation (GRADE) criteria.²¹

Data Synthesis

For studies reporting the frequency of financial burdens (based on the study’s specific measure of material or psychological financial burden), the total sample-weighted prevalence of financial burdens was calculated. Per Cochrane Consumers and Communication Review Group guidelines,²¹ we performed a qualitative synthesis of associations. Quantitative associations were pooled for insurance status and financial burdens,^{23–27} and for financial burdens and medication adherence,^{5,28,29} because these measures could be pooled across multiple studies. We converted reported ORs into log ORs to quantify the effect, using the generic inverse-variance method with a random-effects model to pool ORs. When needed, 1/OR was used for consistency of the referent group in pooled estimates. Adjusted estimates were used where possible for primary analyses to decrease potential confounder bias. Extracted 95% CIs were transformed into standard errors. Heterogeneity of the data was tested with Cochran’s *Q* test and interpreted using the *I*² index. When *I*² = 40%, heterogeneity was explored through sensitivity analyses by eliminating outliers (RevMan 5.3.5 [Reviewer Manager, Copenhagen]).³⁰

Results

Study Selection

Of 5,711 nonduplicate records screened, with 452 abstracts and/or full texts reviewed, 74 studies met the final inclusion criteria per PRISMA (Figure 1). Randomized studies were eligible, but none in the literature met inclusion criteria.

Study and Patient Characteristics

Studies represented an aggregate sample of 598,751 patients with cancer (Table 1). Of 566,531 patients for whom race/ethnicity was reported, 86% were categorized as white, non-Hispanic white, or Caucasian. Of 597,575 patients for whom sex was reported, 54% were female. Weighted mean age was 59.5 years (95% CI, 57.5–62.3). Additional study characteristics are reported in eAppendix 1 and supplemental eTable 2. In 25 reporting studies (of 20,272 patients), the sample-weighted prevalence was 49% (95% CI, 41%–56%) of patients with cancer with any financial burden (Figure 2).

Figure 1. PRISMA flow diagram of study selection.

Financial Burden Measures

Financial burden measures analyzed across studies were as follows: (1) direct material financial burdens of cancer treatment, such as higher OOP expenses, examined in 19% of the 74 studies (n=14); (2) indirect financial burdens, such as lost savings, work, and productivity, and inadequacy of financial resources to meet needs for food, housing, and transport as a result of cancer treatment, examined in 57% of the studies (n=42); and (3) psychological hardships resulting from financial burdens, such as the experience of distress, stress, hardship, and worry, examined in 14% of the studies (n=10) (Table 1). When grouped by type of financial burden, the prevalence ranged from 39% to 64% of patients with direct, indirect, or psychological burden, with no statistically significant difference in prevalence by burden type (see eAppendix 1).

Risk of Bias

Of 74 studies with observational study designs, the primary sources of bias were recall bias in retrospective and cross-sectional study designs and selection bias (additional details in eAppendix 1 and supplemental eTable 1). A total of 72% of the studies had a “good” quality assessment for risk of bias, with a clearly specified study population, participation by >50% of eligible persons, effect estimates provided with variances, a clearly defined

Table 1. Summary of Financial Toxicity Measures and Study Characteristics (N=74)

Study Characteristic	Studies, n (%)
Financial toxicity measure	
Direct material burden	14 (19%)
Measures	Higher total OOP expenses, higher OOP drug expenses, higher OOP-to-income ratio
Indirect burden	42 (57%)
Measures	Lost savings; debt; borrowing; loans; bankruptcy; losing or selling assets; lost/changed housing; inability to meet needs for food, housing, and transport; inability to meet financial demands (medical and insurance coverage, household finances and bills, daily living expenses, leisure spending); lost/changed work and productivity; inadequacy of financial support resources and information; lost/changed health insurance coverage
Psychological burden	10 (14%)
Measures	Distress, stress, hardship, burden, suffering, worry, difficulty, sacrifice, uncertainty, anxiety, worsened circumstances, lack of satisfaction
Study design	
Prospective cohort	10 (14%)
Retrospective cohort	9 (12%)
Cross-sectional	55 (74%)
Study setting	
Academic	21 (28%)
Community	8 (11%)
Academic and community	5 (7%)
Population-based	40 (54%)

Abbreviation: OOP, out-of-pocket.

risk factor and outcome measures consistently measured across study participants, and adjustment for confounders. The strength of pooled evidence ranged from very low to low (additional details in supplemental eTable 3).

Risk Factors Predicting Financial Burdens With Cancer Treatment

A total of 50 studies evaluated the risk factors for financial burdens, demonstrating similar risk factors for the different types of burdens (direct material, indirect, and psychological burdens) (supplemental eTable 4). Evidence supported multiple socioeconomic factors as independent risk factors. Lack of health insurance was associated with increased odds of (more severe) financial burdens.^{6,11,23–25,31–46} In the pooled analysis (of 4 of 15 studies examining insurance status),^{23,24,26,27} patients lacking any drug or health insurance coverage had

Figure 2. Pooled analyses showing sample-weighted prevalence of financial burdens. Weights are from random-effects analysis. Abbreviations: D, direct burdens; I, indirect burdens; P, psychological burdens.

twice the odds of experiencing financial burdens with cancer treatment, compared with patients who had partial/full insurance coverage (OR, 2.09; 95% CI, 1.33–3.30) (Figure 3). A limited number of studies evaluated the association by specific insurance type, showing that Medicare,^{35,38} supplemental,^{6,43} and commercial insurance⁴⁴ coverage were associated with decreased financial burdens compared with patients lacking insurance coverage and those with Medicaid.

Lower income was associated with increased odds of financial burdens,^{6,11,23,24,26,27,31–33,35–37,40–42,47–58} with a dose–response relationship across the income spectrum in a limited number of studies.^{23,52,53} Evidence also indicated that unemployment or job change^{11,24,31,32,36,37,49,51–53,57–65} and lower education were independent risk factors for worse financial burdens,^{23,35,40–42,48,51,61,66} although in 2 studies, education was not a risk factor.^{26,67}

Demographic risk factors for financial burdens included younger age, which was associated with worse financial burdens across the studies.^{11,14,24,25,27,31,32,38,40–42,46,48–50,53–55,57,58,61–63,65,68–72} Evidence supported a dose–response relationship across

the age spectrum.^{11,31,35,42,49} For example, the adjusted probability of “material hardship” decreased from 34% for patients aged 18 to 54 years to 23% for those aged 55 to 64 years, 18% for those aged 65 to 74 years, and 10% for those aged ≥ 75 years.¹¹ Guy et al⁵³ reported a contrasting finding of worse financial burdens in patients aged 50 to 64 versus those aged 18 to 39 years, but this contrasting finding may have been explained by financial burdens measured as OOP expenses as a percentage of family income instead of absolute OOP alone.¹¹ Non-white race and Hispanic ethnicity^{11,68} were associated with increased financial burdens compared with white race and non-Hispanic ethnicity, but the effect sizes of associations varied and race/ethnicity categories were heterogeneous across the studies (eg, ORs ranged from 1.41 to 8.0).^{11,14,24,25,27,31,32,35,37,41,52,68,71,73,74} Female versus male sex was associated with increased financial burdens in a limited number of studies.^{11,14,25,35,54,58,71}

Clinical risk factors for financial burdens, including disease and treatment factors, were investigated in multiple studies. Evidence indicated that no single cancer site, histology, or stage at diagnosis emerged as a consistent, independent risk factor for financial burdens

Figure 3. Pooled analyses showing association between insurance status and financial burdens. Abbreviation: OR, odds ratio.

across the studies. In a limited number of studies, treatment with chemotherapy was associated with increased risk of financial burdens,^{6,36,37,46,49,60,68} whereas receipt of radiation^{35,75} or oncologic surgery were not consistent risk factors. In a limited number of studies, greater treatment duration and intensity were associated with increased financial burdens,^{45,47,52,60} as were cancer recurrence or multiple cancer diagnoses.^{24,37,68} Non-cancer comorbidity and worse overall health and functional status were also factors associated with increased financial burdens in a limited number of studies.^{6,23,36,63}

Evidence across the studies supported a trajectory of financial burdens, with a greater magnitude (effect size) of burdens at cancer diagnosis/active treatment compared with the magnitude of burdens measured after treatment completion. Financial burdens decreased months to years after cancer treatment was completed.^{11,24,25,37,55,57,64,68} The severity of all types of burdens improved over time, including direct,^{57,64} indirect, and psychological.^{11,76} (supplemental eTable 4).

In a limited number of studies on the long-term trajectory of financial burdens, evidence indicated that certain financial burdens persisted years beyond initial treatment. Ell et al⁷⁶ reported that 49% of patients had financial stress at least 12 months after diagnosis, and Jagsi et al³⁷ reported that 33% had persistent financial

decline at 4 years after diagnosis.³⁷ Huntington et al⁵⁰ reported that a longer time from cancer diagnosis was associated with increased financial burdens in patients with chronic multiple myeloma undergoing long-term treatment. A single study implemented a financial navigation intervention. Among patients enrolled in the program, material financial burdens remained unchanged but psychological financial burdens decreased over time. However, this was a single-arm intervention without a “usual-care” group for statistical comparison.⁷⁷

Health Outcomes Associated With Financial Burdens

The association of financial burdens with health behaviors was addressed in 23 studies, and with quality of life and survival in another 23 studies (supplemental eTable 5). Financial burdens, including direct, indirect, and psychological burdens, all demonstrated associations with medical nonadherence (ie, delaying, foregoing, or not completing necessary medical care) to cancer-specific treatment and general medical care.

Cancer-specific treatment nonadherence was associated with financial burdens,^{5,26,28,29,44,70,75,78,79} with 2% to 10% of patients with higher financial burdens reporting nonadherence to chemotherapy and oncology appointments^{28,33,80} and 17% to 30% reporting nonadherence to oral maintenance cancer therapies,

such as imatinib for chronic myeloid leukemia and endocrine therapy for breast cancer.^{5,50,70} In the pooled analysis, patients with higher financial burdens (defined as having greater financial distress, copayments, or direct prescription costs) had nearly twice greater odds of cancer medication nonadherence (OR, 1.70; 95% CI, 1.13–2.56) (Figure 4; sensitivity analyses in eAppendix 1). Higher financial burdens were also associated with general medical nonadherence in a limited number of studies, including higher frequency of omitted or delayed subsequent cancer screening in patients with cancer and survivors; forgoing or delaying cancer survivorship care and follow-up, dental care, ophthalmologic care, prescription medications, and mental and primary medical care^{28,36,37,53,68,81,82}; and lower health self-efficacy and treatment decision-making.⁵⁸

Financial burdens, specifically indirect burdens and psychological burdens, were associated with decreased health-related quality of life (HRQoL) dimensions across the studies (supplemental eTable 5), with more than twice the odds of worse overall physical, mental, emotional, and social function and well-being and increased anxiety, depression, and worry. A dose-response relationship was present, with more severe financial burdens associated with incrementally worse HRQoL scores when measured on continuous scales.^{25,35,47,76,83,84}

In a single study of 231,596 patients with cancer, financial burden measured by bankruptcy was independently associated with increased mortality, with nearly twice the mortality risk compared with those not experiencing bankruptcy (HR, 1.79; 95% CI, 1.64–1.96).⁷¹ No other study in our review investigated the association of other financial burden measures with mortality.

Discussion

This systematic review of 74 studies identifying financial burdens associated with cancer treatment in >500,000 US patients highlights the ongoing critical need to mitigate this problem. The 49% frequency of financial burdens associated with cancer treatment reported in this review, derived across a variety of oncology settings, rivaled the burdens from common clinical toxicities associated with cancer treatment—for example, up to 28% incidence of cardiotoxicity and 34% to 51% incidence of peripheral neuropathy attributed to cancer therapies reported in prior studies.^{85–87} Our findings build upon one prior review that identified a 40% to 50% prevalence estimate of financial hardships from cancer treatment in US patients¹³ and a 28% to 73% prevalence estimate in another review of patients internationally,⁸⁸ validating prior estimates and including contemporary studies. Our study demonstrated that multiple types of burdens—direct material, indirect, and psychological burdens—were uniformly prevalent. This finding underscores the need to assess multidimensional financial burdens in vulnerable patients early in treatment and survivorship, as well as the need for multidimensional interventions to effectively mitigate burdens.

Across the studies, socioeconomic risk factors for direct, indirect, and psychological financial burdens were a lack of health insurance, lower income, and unemployment. Working-aged patients aged <65 years and patients without insurance or with Medicaid coverage (compared with private or Medicare coverage) had excess financial burdens. Multiple studies showed excess financial burdens among women (although the strengths of statistical significance in the *P* values in the studies approached the threshold of .05) and patients of

Figure 4. Pooled analyses showing association between financial burdens and cancer medication adherence. Abbreviation: OR, odds ratio.

nonwhite race and Hispanic ethnicity (although with a varying range of effect sizes in studies). These data identify the disproportionate risk of cancer's financial burdens in vulnerable patient groups and further trigger the hypothesis of whether financial burdens could be a direct and potentially targetable mediator of adherence and mortality disparities in US patients with cancer. Across the studies, the clinical risk factors of initial active treatment, higher treatment intensity, and chemotherapy indicated more severe financial burdens. Evidence did not support the notion that financial burdens differed across cancer disease sites, stages, or histologies, highlighting relevance across patient populations.

The dearth of studies identifying effective interventions reveals a current evidence gap and demonstrates a clinical need for comparative effectiveness studies of interventions to better equip oncology care teams to respond to patients' financial burdens and health impact. Our study's examination of health outcomes demonstrated that direct material, indirect, and psychological financial burdens adversely impacted cancer treatment adherence, including long-term maintenance therapy, and indirect and psychological burdens adversely impacted HRQoL. Additional study is needed on the impact of different types of financial burdens on cancer mortality, given the single study providing evidence of an association between bankruptcy and all-cause mortality.

Our review provides novel insight into the documented gap between patient-reported needs and current oncology practice, wherein oncology provider engagement is not matched with patient-reported financial burdens. In a survey of oncology providers from the iCan Care study, only 34% regularly had a member of their practice discuss financial burdens with patients.¹⁶ Of the patients surveyed who desired provider engagement, only 27% reported that engagement was adequate. Results from our review show that all types of financial burdens and the consequent treatment nonadherence and HRQoL declines could be meaningful targets for direct provider engagement and intervention. Bridging the observed gaps in care would require generalizable interventions implemented across a variety of oncology delivery settings and platforms, including early access to financial counseling,⁷⁷ healthcare and prescription cost assistance programs, enhanced provider cost communication, and patient decision tools.

Limitations of this review reflect those of observational studies. First, causality of associations was not definitively established due to potential attribution or recall bias in retrospective and cross-sectional studies, especially those with solely subjective financial burden measures such as a psychological burden. Countering this weakness was the consistency of risk factor and adherence associations with objectively validated measures, such as

direct material financial burden measures (eg, OOP expenses). The potential risks of recall, misclassification, and selection and attrition bias of the included studies are further detailed in eAppendix 1.

Regarding review-level limitations, because authors may be more likely to publish studies and selectively report outcomes that show statistical significance, the findings may overestimate the occurrence of financial burdens or may not reflect other risk factors and outcomes of financial burdens that may be observed in other populations, such as non-US patients. Pooled analyses were limited, since no current standard measure of financial burden has been established, there were only a limited number of studies whose definitions of financial burden could be quantitatively pooled. Our results provide a visual representation of the sign of effect, but the pooled magnitude of association should be interpreted with caution, recognizing the limitations of meta-analysis from observational studies and the pooling of ORs from different types of populations.^{89,90} The very low to low strength of evidence implied uncertainty about the estimates, and further research could have an important impact on our confidence in the estimate of effects.

Conclusions

Observational evidence supports the findings that cancer treatment's financial burdens are common—experienced in up to half of patients with cancer across disease sites—but that the risks disproportionately affect socioeconomically disadvantaged and working-aged patients. Evidence further shows that financial burdens are associated with worse treatment adherence and HRQoL. Results from a single study also suggest that financial burdens are associated with worse mortality. Synthesized results provide a basis for identifying patients at risk for financial toxicity who would benefit from oncology provider and care team engagement. However, due to the paucity of evidence identifying effective measures to mitigate financial burdens in patients, our results highlight a critical need for ongoing efforts to develop interventions to equip oncology care teams to respond to the meaningful adverse impact of financial burdens in patients with cancer.

Submitted August 1, 2018; accepted for publication March 29, 2019.

Author contributions: Study concept and design: Smith, Lopez-Olivo, Advani, Geng, Giordano, Volk. Data acquisition, analysis, and interpretation: All authors. Manuscript preparation: All authors. Critical revision: All authors. Final approval: All authors.

Disclosures: Dr. Smith has disclosed that she has received grant/research support from the NIH (K07 CA211804-01), the Center for Radiation Oncology Research, and the University Cancer Foundation via the Institutional Research Grant Program at The University of Texas MD Anderson Cancer Center. Dr. Lopez-Olivo has disclosed that she receives grant/research support from the Rheumatology Research Foundation (grant 14119629). Dr. Giordano has

disclosed that she has received grant/research support from the Susan G. Komen Foundation (SAC 150061). The remaining authors have not received any financial consideration from any person or organization to support the preparation, analysis, results, or discussion of this article.

Funding: This study was supported in part by The University of Texas MD Anderson Cancer Center Duncan Family Institute for Cancer Prevention and

Risk Assessment and the NCI Cancer Center Support grant, and grant P50 CA016672 (MD Anderson Cancer Center).

Correspondence: Grace L. Smith, MD, PhD, MPH, Department of Radiation Oncology and Department of Health Services Research, The University of Texas MD Anderson Cancer Center, 1400 Pressler Street, Houston, TX 77030. Email: glsmith@mdanderson.org

References

- Azzani M, Roslani AC, Su TT. The perceived cancer-related financial hardship among patients and their families: a systematic review. *Support Care Cancer* 2015;23:889–898.
- Miller KD, Siegel RL, Lin CC, et al. Cancer treatment and survivorship statistics, 2016. *CA Cancer J Clin* 2016;66:271–289.
- Institute of Medicine. Ensuring Patient Access to Affordable Cancer Drugs: Workshop Summary. Washington, DC: National Academies Press; 2014.
- Soni A. Trends in use and expenditures for cancer treatment among adults 18 and older, U.S. civilian noninstitutionalized population, 2001 and 2011. Statistical Brief #443. Rockville, MD: Agency for Healthcare Research and Quality; 2014.
- Dusetzina SB, Winn AN, Abel GA, et al. Cost sharing and adherence to tyrosine kinase inhibitors for patients with chronic myeloid leukemia. *J Clin Oncol* 2014;32:306–311.
- Davidoff AJ, Erten M, Shaffer T, et al. Out-of-pocket health care expenditure burden for Medicare beneficiaries with cancer. *Cancer* 2013; 119:1257–1265.
- Guy GP Jr, Ekwueme DU, Yabroff KR, et al. Economic burden of cancer survivorship among adults in the United States. *J Clin Oncol* 2013;31: 3749–3757.
- Yabroff KR, Lamont EB, Mariotto A, et al. Cost of care for elderly cancer patients in the United States. *J Natl Cancer Inst* 2008;100:630–641.
- Short PF, Moran JR, Puneekar R. Medical expenditures of adult cancer survivors aged <65 years in the United States. *Cancer* 2011;117: 2791–2800.
- Finkelstein EA, Tangka FK, Trogdon JG, et al. The personal financial burden of cancer for the working-aged population. *Am J Manag Care* 2009;15:801–806.
- Yabroff KR, Dowling EC, Guy GP Jr, et al. Financial hardship associated with cancer in the United States: findings from a population-based sample of adult cancer survivors. *J Clin Oncol* 2016;34:259–267.
- National Cancer Institute. Financial toxicity and cancer treatment (PDQ)—health professional version. Available at: <https://www.cancer.gov/about-cancer/managing-care/track-care-costs/financial-toxicity-hp-pdq>. Accessed February 14, 2018.
- Altice CK, Banegas MP, Tucker-Seeley RD, et al. Financial hardships experienced by cancer survivors: a systematic review. *J Natl Cancer Inst* 2016;109:djw205.
- Ramsey S, Blough D, Kirchhoff A, et al. Washington State cancer patients found to be at greater risk for bankruptcy than people without a cancer diagnosis. *Health Aff (Millwood)* 2013;32:1143–1152.
- Congressional Budget Office. Repealing the individual health insurance mandate: an updated estimate. Available at: <https://www.cbo.gov/publication/53300>. Accessed February 14, 2018.
- Jagsi R, Ward KC, Abrahamse PH, et al. Unmet need for clinician engagement about financial toxicity after diagnosis of breast cancer. *Cancer* 2018;124:3668–3676.
- Shih YT, Chien CR. A review of cost communication in oncology: patient attitude, provider acceptance, and outcome assessment. *Cancer* 2017; 123:928–939.
- O'Connor JM, Kircher SM, de Souza JA. Financial toxicity in cancer care. *J Community Support Oncol* 2016;14:101–106.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–269.
- Meeker CR, Geynisman DM, Egleston BL, et al. Relationships among financial distress, emotional distress, and overall distress in insured patients with cancer. *J Oncol Pract* 2016;12:e755–764.
- Ryan R; Cochrane Consumers and Communication Review Group. Cochrane Consumers and Communication Review Group: data synthesis and analysis. Available at: <http://cccr.cochrane.org/sites/cccr.org.cochrane.org/files/public/uploads/Analysis.pdf>. Accessed May 9, 2019.
- National Institutes of Health; National Heart, Lung, and Blood Institute. Study quality assessment tools. Available at: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>. Accessed February 14, 2018.
- Whitney RL, Bell JF, Reed SC, et al. Predictors of financial difficulties and work modifications among cancer survivors in the United States. *J Cancer Surviv* 2016;10:241–250.
- Banegas MP, Guy GP Jr, de Moor JS, et al. For working-age cancer survivors, medical debt and bankruptcy create financial hardships. *Health Aff (Millwood)* 2016;35:54–61.
- Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
- Pezzin LE, O'Neil MB, Nattinger AB. The economic consequences of breast cancer adjuvant hormonal treatments. *J Gen Intern Med* 2009; 24(Suppl 2):S446–450.
- Pisu M, Kenzik KM, Oster RA, et al. Economic hardship of minority and non-minority cancer survivors 1 year after diagnosis: another long-term effect of cancer? *Cancer* 2015;121:1257–1264.
- Bestvina CM, Zullig LL, Rushing C, et al. Patient-oncologist cost communication, financial distress, and medication adherence. *J Oncol Pract* 2014;10:162–167.
- Zullig LL, Peppercorn JM, Schrag D, et al. Financial distress, use of cost-coping strategies, and adherence to prescription medication among patients with cancer. *J Oncol Pract* 2013;9(Suppl 6):608–635.
- The Cochrane Collaboration. Cochrane Handbook for Systematic Reviews of Interventions. Version 6. Available at: <https://training.cochrane.org/handbook>. Accessed May 9, 2019.
- Veenstra CM, Regenbogen SE, Hawley ST, et al. A composite measure of personal financial burden among patients with stage III colorectal cancer. *Med Care* 2014;52:957–962.
- Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
- Markman M, Luce R. Impact of the cost of cancer treatment: an internet-based survey. *J Oncol Pract* 2010;6:69–73.
- Callahan C, Brintzenhofesoc K. Financial quality of life for patients with cancer: an exploratory study. *J Psychosoc Oncol* 2015;33:377–394.
- Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
- Jagsi R, Hawley ST, Abrahamse P, et al. Impact of adjuvant chemotherapy on long-term employment of survivors of early-stage breast cancer. *Cancer* 2014;120:1854–1862.
- Jagsi R, Pottow JAE, Griffith KA, et al. Long-term financial burden of breast cancer: experiences of a diverse cohort of survivors identified through population-based registries. *J Clin Oncol* 2014;32:1269–1276.
- Kircher SM, Johansen ME, Nimeiri HS, et al. Impact of Medicare Part D on out-of-pocket drug costs and medical use for patients with cancer. *Cancer* 2014;120:3378–3384.
- Lathan CS, Cronin A, Tucker-Seeley R, et al. Association of financial strain with symptom burden and quality of life for patients with lung or colorectal cancer. *J Clin Oncol* 2016;34:1732–1740.
- Lee M, Khan MM. Gender differences in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:384–393.
- Lee M, Salloum RG. Racial and ethnic disparities in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10: 534–544.
- Martin MY, Fouad MN, Oster RA, et al. What do cancer patients worry about when making decisions about treatment? Variation across racial/ethnic groups. *Support Care Cancer* 2014;22:233–244.
- Narang AK, Nicholas LH. Out-of-pocket spending and financial burden among Medicare beneficiaries with cancer. *JAMA Oncol* 2017;3:757–765.

44. Streeter SB, Schwartzberg L, Husain N, et al. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *Am J Manag Care* 2011;17(Suppl 5):SP38–44.
45. Tucker-Seeley RD, Abel GA, Uno H, et al. Financial hardship and the intensity of medical care received near death. *Psychooncology* 2015;24:572–578.
46. Wan Y, Gao X, Mehta S, et al. Indirect costs associated with metastatic breast cancer. *J Med Econ* 2013;16:1169–1178.
47. Abel GA, Albelda R, Khera N, et al. Financial hardship and patient-reported outcomes after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016;22:1504–1510.
48. Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
49. Regenbogen SE, Veenstra CM, Hawley ST, et al. The personal financial burden of complications after colorectal cancer surgery. *Cancer* 2014;120:3074–3081.
50. Huntington SF, Weiss BM, Vogl DT, et al. Financial toxicity in insured patients with multiple myeloma: a cross-sectional pilot study. *Lancet Haematol* 2015;2:e408–416.
51. Chongpison Y, Hornbrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
52. de Souza JA, Yap BJ, Wroblewski K, et al. Measuring financial toxicity as a clinically relevant patient-reported outcome: the validation of the Comprehensive Score for financial Toxicity (COST). *Cancer* 2017;123:476–484.
53. Guy GP Jr, Yabroff KR, Ekwueme DU, et al. Healthcare expenditure burden among non-elderly cancer survivors, 2008–2012. *Am J Prev Med* 2015;49(Suppl 5):S489–497.
54. Houts PS, Harvey HA, Simmonds MA, et al. Characteristics of patients at risk for financial burden because of cancer and its treatment. *J Psychosoc Oncol* 1985;3:15–22.
55. Jones WC, Parry C, Devine S, et al. Prevalence and predictors of distress in posttreatment adult leukemia and lymphoma survivors. *J Psychosoc Oncol* 2015;33:124–141.
56. Langa KM, Fendrick AM, Chernew ME, et al. Out-of-pocket health-care expenditures among older Americans with cancer. *Value Health* 2004;7:186–194.
57. Pisu M, Azuero A, Benz R, et al. Out-of-pocket costs and burden among rural breast cancer survivors. *Cancer Med* 2017;6:572–581.
58. Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
59. Blinder V, Patil S, Eberle C, et al. Early predictors of not returning to work in low-income breast cancer survivors: a 5-year longitudinal study. *Breast Cancer Res Treat* 2013;140:407–416.
60. Goodwin JA, Coleman EA, Sullivan E, et al. Personal financial effects of multiple myeloma and its treatment. *Cancer Nurs* 2013;36:301–308.
61. Bernard DSM, Farr SL, Fang Z. National estimates of out-of-pocket health care expenditure burdens among nonelderly adults with cancer: 2001 to 2008. *J Clin Oncol* 2011;29:2821–2826.
62. Ekwueme DU, Yabroff KR, Guy GP Jr, et al. Medical costs and productivity losses of cancer survivors—United States, 2008–2011. *MMWR Morb Mortal Wkly Rep* 2014;63:505–510.
63. Khera N, Chang YH, Hashmi S, et al. Financial burden in recipients of allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2014;20:1375–1381.
64. Meneses K, Azuero A, Hassey L, et al. Does economic burden influence quality of life in breast cancer survivors? *Gynecol Oncol* 2012;124:437–443.
65. Francoeur RB. Cumulative financial stress and strain in palliative radiation outpatients: the role of age and disability. *Acta Oncol* 2005;44:369–381.
66. de Souza JA, Wong YN. Financial distress in cancer patients. *J Med Person* 2013;11:73–77.
67. de Souza JA, Yap BJ, Hlubocky FJ, et al. The development of a financial toxicity patient-reported outcome in cancer: the COST measure. *Cancer* 2014;120:3245–3253.
68. Kent EE, Forsythe LP, Yabroff KR, et al. Are survivors who report cancer-related financial problems more likely to forgo or delay medical care? *Cancer* 2013;119:3710–3717.
69. Meeker CR, Wong YN, Eggleston BL, et al. Distress and financial distress in adults with cancer: an age-based analysis. *J Natl Compr Canc Netw* 2017;15:1224–1233.
70. Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
71. Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
72. Zheng Z, Yabroff KR, Guy GP Jr, et al. Annual medical expenditure and productivity loss among colorectal, female breast, and prostate cancer survivors in the United States. *J Natl Cancer Inst* 2015;108:djv382.
73. Wheeler SB, Spencer JC, Pinheiro LC, et al. Financial impact of breast cancer in black versus white women. *J Clin Oncol* 2018;36:1695–1701.
74. Palmer NR, Weaver KE, Hauser SP, et al. Disparities in barriers to follow-up care between African American and White breast cancer survivors. *Support Care Cancer* 2015;23:3201–3209.
75. Farias AJ, Hansen RN, Zeliadt SB, et al. The association between out-of-pocket costs and adherence to adjuvant endocrine therapy among newly diagnosed breast cancer patients. *Am J Clin Oncol* 2018;41:708–715.
76. Ell K, Xie B, Wells A, et al. Economic stress among low-income women with cancer: effects on quality of life. *Cancer* 2008;112:616–625.
77. Shankaran V, Leahy T, Steelquist J, et al. Pilot feasibility study of an oncology financial navigation program. *J Oncol Pract* 2018;14:e122–129.
78. Farias AJ, Du XL. Association between out-of-pocket costs, race/ethnicity, and adjuvant endocrine therapy adherence among Medicare patients with breast cancer. *J Clin Oncol* 2017;35:86–95.
79. Bender CM, Gentry AL, Brufsky AM, et al. Influence of patient and treatment factors on adherence to adjuvant endocrine therapy in breast cancer. *Oncol Nurs Forum* 2014;41:274–285.
80. Zafar SY, Peppercorn JM, Schrag D, et al. The financial toxicity of cancer treatment: a pilot study assessing out-of-pocket expenses and the insured cancer patient's experience. *Oncologist* 2013;18:381–390.
81. Nipp RD, Zullig LL, Samsa G, et al. Identifying cancer patients who alter care or lifestyle due to treatment-related financial distress. *Psychooncology* 2016;25:719–725.
82. DiMartino LD, Birken SA, Mayer DK. The relationship between cancer survivors' socioeconomic status and reports of follow-up care discussions with providers. *J Cancer Educ* 2017;32:749–755.
83. Khera N, Chow EJ, Leisenring WM, et al. Factors associated with adherence to preventive care practices among hematopoietic cell transplantation survivors. *Biol Blood Marrow Transplant* 2011;17:995–1003.
84. Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
85. Yeh ETH, Bickford CL. Cardiovascular complications of cancer therapy: incidence, pathogenesis, diagnosis, and management. *J Am Coll Cardiol* 2009;53:2231–2247.
86. Peng L, Ye X, Zhou Y, et al. Meta-analysis of incidence and risk of peripheral neuropathy associated with intravenous bortezomib. *Support Care Cancer* 2015;23:2813–2824.
87. Peng L, Bu Z, Ye X, et al. Incidence and risk of peripheral neuropathy with nab-paclitaxel in patients with cancer: a meta-analysis. *Eur J Cancer Care [Br]* 2017;26:e12407.
88. Gordon LG, Merolini KMD, Lowe A, et al. A systematic review of financial toxicity among cancer survivors: we can't pay the co-pay. *Patient* 2017;10:295–309.
89. Norton EC, Dowd BE. Log odds and the interpretation of logit models. *Health Serv Res* 2018;53:859–878.
90. Norton EC, Dowd BE, Maciejewski ML. Odds ratios—current best practice and use. *JAMA* 2018;320:84–85.
91. de Souza JA, Kung S, O'Connor J, et al. Determinants of patient-centered financial stress in patients with locally advanced head and neck cancer. *J Oncol Pract* 2017;13:e310–318.
92. Gupta D, Lis CG, Grutsch JF. Perceived cancer-related financial difficulty: implications for patient satisfaction with quality of life in advanced cancer. *Support Care Cancer* 2007;15:1051–1056.

See JNCCN.org for supplemental online content.

Supplemental online content for:

Financial Burdens of Cancer Treatment: A Systematic Review of Risk Factors and Outcomes

Grace L. Smith, MD, PhD, MPH; Maria A. Lopez-Olivo, MD, PhD; Pragati G. Advani, MD, DrPH; Matthew S. Ning, MD; Yimin Geng, MS, MSLIS; Sharon H. Giordano, MD, MPH; and Robert J. Volk, PhD

J Natl Compr Canc Netw 2019;17(10):1184–1192

eFigure 1: Search Strategy for MEDLINE (Ovid)

eTable 1: Quality Assessment: Assessing Risk of Bias of Individual Studies

eTable 2: Study and Patient Characteristics and Measures of Financial Toxicity Burdens

eTable 3: Strength of Evidence Assessment Using GRADE Criteria

eTable 4: Risk Factors of Financial Toxicity Burdens

eTable 5: Financial Toxicity Burdens and Health Outcomes

eAppendix 1: Supplementary Results

1	exp *Health Expenditures/
2	exp *Health Care Costs/
3	exp *"Cost of Illness"/
4	exp Financing, Personal/
5	*Fees/ and Charges/sn [Statistics & Numerical Data]
6	(Economic adj (bankruptcy or Insolvency or hardship or burden or stress or distress or difficult*)).ti,ab,kf.
7	((("out of pocket" or cost or cost* or expenditure* or expense or expenses or spending*) adj15 cancer).tw,kf.
8	treatment costs.ti,ab.
9	Financial.ti,ab,kf.
10	or/1-9
11	exp *Neoplasms/ or Neoplasms/ec [Economics]
12	(cancer* or neoplas* or tumor* or leuk?emia* or lymphoma* or malignan* or oncolog* or "Cell Transplantation").ti,kf.
13	11 or 12
14	limit 13 to english language
15	10 and 14
16	(cancer and treatment and decision*).ti. and 15
17	exp Health Care Costs/sn [Statistics & Numerical Data]
18	exp *"Cost Sharing"/
19	exp *Financing, Personal/ec, sn [Economics, Statistics & Numerical Data]
20	exp *Health Expenditures/ec, sn [Economics, Statistics & Numerical Data]
21	6 or 9 or 17 or 18 or 19 or 20
22	15 and 21
23	22 and 12
24	((medical or health) adj expenditure*).ti.
25	Financial.ti.
26	out-of-pocket.ti.
27	(Economic adj (bankruptcy or Insolvency or hardship or burden or stress or distress or difficult*)).ti.
28	or/24-27
29	13 and 28
30	limit 29 to english language
31	*Health Behavior/
32	exp patient compliance/ or patient dropouts/ or exp treatment refusal/
33	*Stress, Psychological/
34	exp *Adaptation, Psychological/
35	exp self efficacy/ or "self efficacy".tw,kf.
36	exp *Anxiety/
37	(coping or adaptive or adaptation).tw,kf.
38	(distress or distresses or stress or stresses).tw,kf.
39	(patient* adj3 ("non compliance" or "non-adherence" or noncompliance or nonadherence)).tw,kf.
40	exp "Patient Acceptance of Health Care"/
41	exp Withholding Treatment/
42	*Medication Adherence/sn [Statistics & Numerical Data]
43	(cancer and treatment and decision*).ti.
44	*Decision Making/
45	((delay* or forgo*) adj3 (care or treatment)).tw,kf.

(continued on next page)

eFigure 1. Search strategy for MEDLINE (Ovid).

```

46 | or/31-45
47 | 23 and 46
48 | remove duplicates from 47
49 | exp Survival Rate/
50 | exp *Survival Analysis/
51 | survival.ti.
52 | exp *Mortality/sn, td [Statistics & Numerical Data, Trends]
53 | exp *"Quality of Life"/
54 | "Quality of Life".ti,kf.
55 | or/49-54
56 | 23 and 55
57 | remove duplicates from 56
58 | 57 not 48
59 | exp *insurance, health/
60 | *Insurance Coverage/
61 | exp *Income/
62 | Insurance Carriers/
63 | exp *Socioeconomic Factors/
64 | exp *Demography/
65 | exp "Social Determinants of Health"/
66 | exp *Neoplasms/ec [Economics]
67 | exp *Drug Therapy/ec [Economics]
68 | exp *Antineoplastic Agents/ec [Economics]
69 | exp *Specialties, Surgical/ec [Economics]
70 | Therapeutics/ec [Economics]
71 | exp *Comorbidity/
72 | exp *Radiotherapy/ec [Economics]
73 | exp *Neoplasm Recurrence, Local/ec [Economics]
74 | Recurrence/ and Neoplasms/ and 12
75 | ec.fs.
76 | 74 and 75
77 | or/59-73,76
78 | 23 and 77
79 | 48 or 57 or 78
80 | 16 or 79 or 30
81 | remove duplicates from 80
82 | or/1-7,28
83 | "financial relationships to disclose".ab.
84 | Financial Disclosures.ab.
85 | financial conflicts of interest.ab.
86 | financial conflicts of interests.ab.
87 | financial interests.ab.
88 | financial interest.ab.
89 | 83 or 84 or 85 or 86 or 87 or 88
90 | 81 and 89
91 | 81 not 90
92 | 82 and 90
93 | 91 or 92

```

eFigure 1. Search strategy for MEDLINE (Ovid). (cont.)

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies

Reference	Criteria														Global rating
	1. Was the research question or objective in this article clearly stated?	2. Was the study population clearly specified and defined?	3. Was the participation rate of eligible persons at least 50%?	4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?	5. Were a sample size justification, power description, or variance and effect estimates provided?	6. For the analyses in this article, was the exposure of interest measured prior to the outcomes being measured?	7. Was the time frame sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?	8. For exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcome (eg, categories of exposure or exposure measured as a continuous variable)?	9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	10. Was the exposure assessed more than once over time?	11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	12. Were the outcome assessors blinded to the exposure status of participants?	13. Was loss to follow-up after the exposure baseline $\leq 20\%$?	14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposures and outcomes?	
1	Yes	Yes	NR	Yes	No	Yes	NA	NA	Yes	No	Yes	NR	NA	No	Fair
2	Yes	Yes	Yes	Yes	Yes	Yes	NA	NA	Yes	No	Yes	Yes	NA	Yes	Good
3	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	NR	Yes	Yes	Good
4	Yes	Yes	Yes	Yes	Yes	NA	NA	Yes	Yes	No	Yes	NR	NA	Yes	Good
5	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
6	Yes	Yes	NR	Yes	Yes	NA	NA	NA	Yes	No	Yes	NR	NA	Yes	Fair
7	Yes	Yes	NR	Yes	Yes	CD	Yes	Yes	Yes	No	Yes	NR	Yes	Yes	Good
8	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
9	Yes	Yes	NR	Yes	No	CD	Yes	NA	Yes	No	Yes	NR	NA	Yes	Fair
10	Yes	Yes	NR	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
11	Yes	Yes	CD	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
12	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	No	No	Yes	NR	NA	Yes	Fair
13	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
14	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	No	No	Yes	NR	NA	Yes	Fair
15	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	No	Yes	Yes	NR	NR	Yes	Fair
16	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	NR	No	Yes	Fair
17	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
18	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
19	Yes	Yes	NR	Yes	No	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
20	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
21	Yes	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
22	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NA	Yes	No	Yes	NR	NA	Yes	Good
23	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good

(continued on next page)

Abbreviations: CD, cannot determine; NA, not applicable; NR, not reported.

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies (cont.)

Reference	Criteria														Global rating
	1. Was the research question or objective in this article clearly stated?	2. Was the study population clearly specified and defined?	3. Was the participation rate of eligible persons at least 50%?	4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?	5. Were a sample size justification, power description, or variance and effect estimates provided?	6. For the analyses in this article, were exposures of interest measured prior to the outcomes being measured?	7. Was the time frame sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?	8. For exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcome (eg, categories of exposure or exposure measured as a continuous variable)?	9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	10. Was the exposure assessed more than once over time?	11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	12. Were the outcome assessors blinded to the exposure status of participants?	13. Was loss to follow-up after baseline and outcomes? ≤20%?	14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposures and outcomes?	
24	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
25	Yes	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
26	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
27	Yes	Yes	CD	Yes	No	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
28	Yes	Yes	NR	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
29	Yes	Yes	Yes	Yes	Yes	NA	Yes	CD	Yes	No	Yes	NR	NA	Yes	Good
30	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
31	No	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
32	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
33	Yes	Yes	CD	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
34	Yes	Yes	NR	Yes	Yes	NA	Yes	NA	Yes	No	Yes	No	NR	Yes	Good
35	Yes	Yes	CD	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
36	Yes	Yes	NR	Yes	Yes	NA	Yes	Yes	Yes	Yes	Yes	NR	NR	Yes	Good
37	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	No	CD	NA	Yes	Fair
38	Yes	No	NR	NR	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
39	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
40	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
41	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
42	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	NR	No	Yes	Good
43	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	NR	No	Yes	Good
44	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good
45	Yes	Yes	NA	No	No	No	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Fair
46	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes	Good

(continued on next page)

Abbreviations: CD, cannot determine; NA, not applicable; NR, not reported.

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies (cont.)

Reference	Criteria													Global rating
	1. Was the research question or objective in this article clearly stated?	2. Was the study population clearly specified and defined?	3. Was the participation rate of eligible persons at least 50%?	4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?	5. Were a sample size justification, power description, or variance and effect estimates provided?	6. For the analyses in this article, was the exposure of interest measured prior to the outcomes being measured?	7. Was the time frame sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?	8. For exposures that vary in amount or level, did the study examine different levels of the exposure as related to the outcome (eg, categories of exposure or exposure measured as a continuous variable)?	9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	10. Was the exposure assessed more than once over time?	11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	12. Were the outcome assessors blinded to the exposure status of participants?	13. Was loss to follow-up after baseline $\leq 20\%$?	14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposures and outcomes?
47	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
48	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	No
49	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NR	No	No
50	Yes	Yes	CD	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	NR	NA	Yes
51	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
52	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	No	NR	NA	No
53	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
54	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
55	Yes	Yes	Yes	No	No	Yes	Yes	Yes	Yes	No	Yes	NR	NA	Yes
56	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
57	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NR	No	Yes
58	Yes	Yes	No	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
59	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	NR	Yes	Yes
60	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
61	Yes	Yes	Yes	Yes	No	NA	Yes	Yes	Yes	No	Yes	NR	NA	No
62	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	NR	NA	Yes
63	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
64	Yes	Yes	Yes	Yes	NR	Yes	Yes	Yes	Yes	Yes	Yes	NR	Yes	Yes
65	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
66	Yes	Yes	No	No	NR	No	Yes	Yes	Yes	No	Yes	NR	NA	No
67	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
68	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
69	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes

Abbreviations: CD, cannot determine; NA, not applicable; NR, not reported.

(continued on next page)

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies (cont.)

	Criteria													
	4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?	3. Was the participation rate of eligible persons at least 50%?	2. Was the study population clearly specified and defined?	1. Was the research question or objective in this article clearly stated?	5. Were a sample size justification, power description, or variance and effect estimates provided?	6. For the analyses in this article, was the exposure measured prior to the outcomes being measured?	7. Was the time frame sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?	8. For exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcome (eg, categories of exposure or exposure measured as a continuous variable)?	9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	10. Was the exposure assessed more than once over time?	11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?	12. Were the outcome assessors blinded to the exposure status of participants?	13. Was loss to follow-up after the exposure $\leq 20\%$?	14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposures and outcomes?
Reference	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
70	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
71	Yes	No	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes
72	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	No	NA	Yes
73	Yes	Yes	Yes	Yes	No	NA	Yes	Yes	Yes	No	Yes	NR	NA	No
74	Yes	Yes	Yes	Yes	Yes	NA	Yes	Yes	Yes	No	Yes	NR	NA	Yes

Abbreviations: CD, cannot determine; NA, not applicable; NR, not reported.

(references on next page)

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies (cont.)

References

- Shankaran V, Leahy T, Steelquist J, et al. Pilot feasibility study of an oncology financial navigation program. *J Oncol Pract* 2018;14:e122–129.
- Wheeler SB, Spencer JC, Pinheiro LC, et al. Financial impact of breast cancer in black versus white women. *J Clin Oncol* 2018;36:1695–1701.
- de Souza JA, Kung S, O'Connor J, et al. Determinants of patient-centered financial stress in patients with locally advanced head and neck cancer. *J Oncol Pract* 2017;13:e310–318.
- de Souza JA, Yap BJ, Wroblewski K, et al. Measuring financial toxicity as a clinically relevant patient-reported outcome: the validation of the Comprehensive Score for financial Toxicity (COST). *Cancer* 2017;123:476–484.
- Farias AJ, Du XL. Association between out-of-pocket costs, race/ethnicity, and adjuvant endocrine therapy adherence among Medicare patients with breast cancer. *J Clin Oncol* 2017;35:86–95.
- Meeker CR, Wong YN, Egleston BL, et al. Distress and financial distress in adults with cancer: an age-based analysis. *J Natl Compr Canc Netw* 2017;15:1224–1233.
- Pisu M, Azuero A, Benz R, et al. Out-of-pocket costs and burden among rural breast cancer survivors. *Cancer Med* 2017;6:572–581.
- Abel GA, Albelda R, Khera N, et al. Financial hardship and patient-reported outcomes after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016;22:1504–1510.
- Banegas MP, Guy GP Jr, de Moor JS, et al. For working-age cancer survivors, medical debt and bankruptcy create financial hardships. *Health Aff (Millwood)* 2016;35:54–61.
- Chongpison Y, Hornbrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
- Delgado-Guay MO, Chisholm G, Williams J, et al. Frequency, intensity, and correlates of spiritual pain in advanced cancer patients assessed in a supportive/palliative care clinic. *Palliat Support Care* 2016;14:341–348.
- DiMartino LD, Birken SA, Mayer DK. The relationship between cancer survivors' socioeconomic status and reports of follow-up care discussions with providers. *J Cancer Educ* 2017;32:749–755.
- Farias AJ, Hansen RN, Zeliadt SB, et al. The association between out-of-pocket costs and adherence to adjuvant endocrine therapy among newly diagnosed breast cancer patients. *Am J Clin Oncol* 2018;41:708–715.
- Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
- Keim-Malpass J, Levine B, Danhauer SC, et al. Work-related perceptions and quality of life among breast cancer survivors. *Psychooncology* 2016;25:873–876.
- Lathan CS, Cronin A, Tucker-Seeley R, et al. Association of financial strain with symptom burden and quality of life for patients with lung or colorectal cancer. *J Clin Oncol* 2016;34:1732–1740.
- Lee M, Khan MM. Gender differences in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:384–393.
- Lee M, Salloum RG. Racial and ethnic disparities in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:534–544.
- Meeker CR, Geynisman DM, Egleston BL, et al. Relationships among financial distress, emotional distress, and overall distress in insured patients with cancer. *J Oncol Pract* 2016;12:e755–764.
- Narang AK, Nicholas LH. Out-of-pocket spending and financial burden among Medicare beneficiaries with cancer. *JAMA Oncol* 2017;3:757–765.
- Nipp RD, Zullig LL, Samsa G, et al. Identifying cancer patients who alter care or lifestyle due to treatment-related financial distress. *Psychooncology* 2016;25:719–725.
- Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
- Whitney RL, Bell JF, Reed SC, et al. Predictors of financial difficulties and work modifications among cancer survivors in the United States. *J Cancer Surviv* 2016;10:241–250.
- Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
- Yabroff KR, Dowling EC, Guy GP Jr, et al. Financial hardship associated with cancer in the United States: findings from a population-based sample of adult cancer survivors. *J Clin Oncol* 2016;34:259–267.
- Zheng Z, Yabroff KR, Guy GP Jr, et al. Annual medical expenditure and productivity loss among colorectal, female breast, and prostate cancer survivors in the United States. *J Natl Cancer Inst* 2015;108:djv382.
- Callahan C, Brintzenhofesoc K. Financial quality of life for patients with cancer: an exploratory study. *J Psychosoc Oncol* 2015;33:377–394.
- Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
- Guy GP Jr, Yabroff KR, Ekwueme DU, et al. Healthcare expenditure burden among non-elderly cancer survivors, 2008–2012. *Am J Prev Med* 2015;49(Suppl 5):S489–497.
- Huntington SF, Weiss BM, Vogl DT, et al. Financial toxicity in insured patients with multiple myeloma: a cross-sectional pilot study. *Lancet Haematol* 2015;2:e408–416.
- Jones WC, Parry C, Devine S, et al. Prevalence and predictors of distress in posttreatment adult leukemia and lymphoma survivors. *J Psychosoc Oncol* 2015;33:124–141.
- Palmer NR, Weaver KE, Hauser SP, et al. Disparities in barriers to follow-up care between African American and White breast cancer survivors. *Support Care Cancer* 2015;23:3201–3209.
- Pisu M, Kenzik KM, Oster RA, et al. Economic hardship of minority and non-minority cancer survivors 1 year after diagnosis: another long-term effect of cancer? *Cancer* 2015;121:1257–1264.
- Tucker-Seeley RD, Abel GA, Uno H, et al. Financial hardship and the intensity of medical care received near death. *Psychooncology* 2015;24:572–578.
- Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
- Bender CM, Gentry AL, Brufsky AM, et al. Influence of patient and treatment factors on adherence to adjuvant endocrine therapy in breast cancer. *Oncol Nurs Forum* 2014;41:274–285.
- Bestvina CM, Zullig LL, Rushing C, et al. Patient-oncologist cost communication, financial distress, and medication adherence. *J Oncol Pract* 2014;10:162–167.
- de Souza JA, Yap BJ, Hlubocky FJ, et al. The development of a financial toxicity patient-reported outcome in cancer: the COST measure. *Cancer* 2014;120:3245–3253.
- Dusetzina SB, Winn AN, Abel GA, et al. Cost sharing and adherence to tyrosine kinase inhibitors for patients with chronic myeloid leukemia. *J Clin Oncol* 2014;32:306–311.
- Ekwueme DU, Yabroff KR, Guy GP Jr, et al. Medical costs and productivity losses of cancer survivors—United States, 2008–2011. *MMWR Morb Mortal Wkly Rep* 2014;63:505–510.
- Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
- Jaggi R, Hawley ST, Abraham P, et al. Impact of adjuvant chemotherapy on long-term employment of survivors of early-stage breast cancer. *Cancer* 2014;120:1854–1862.
- Jagsi R, Pottow JAE, Griffith KA, et al. Long-term financial burden of breast cancer: experiences of a diverse cohort of survivors identified through population-based registries. *J Clin Oncol* 2014;32:1269–1276.

eTable 1. Quality Assessment: Assessing Risk of Bias of Individual Studies (cont.)

44. Khera N, Chang YH, Hashmi S, et al. Financial burden in recipients of allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2014;20:1375–1381.
45. Kircher SM, Johansen ME, Nimeiri HS, et al. Impact of Medicare Part D on out-of-pocket drug costs and medical use for patients with cancer. *Cancer* 2014;120:3378–3384.
46. Martin MY, Fouad MN, Oster RA, et al. What do cancer patients worry about when making decisions about treatment? Variation across racial/ethnic groups. *Support Care Cancer* 2014;22:233–244.
47. Regenbogen SE, Veenstra CM, Hawley ST, et al. The personal financial burden of complications after colorectal cancer surgery. *Cancer* 2014;120:3074–3081.
48. Veenstra CM, Regenbogen SE, Hawley ST, et al. A composite measure of personal financial burden among patients with stage III colorectal cancer. *Med Care* 2014;52:957–962.
49. Blinder V, Patil S, Eberle C, et al. Early predictors of not returning to work in low-income breast cancer survivors: a 5-year longitudinal study. *Breast Cancer Res Treat* 2013;140:407–416.
50. Davidoff AJ, Erten M, Shaffer T, et al. Out-of-pocket health care expenditure burden for Medicare beneficiaries with cancer. *Cancer* 2013;119:1257–1265.
51. Dowling EC, Chawla N, Forsythe LP, et al. Lost productivity and burden of illness in cancer survivors with and without other chronic conditions. *Cancer* 2013;119:3393–3401.
52. Goodwin JA, Coleman EA, Sullivan E, et al. Personal financial effects of multiple myeloma and its treatment. *Cancer Nurs* 2013;36:301–308.
53. Hamilton JG, Wu LM, Austin JE, et al. Economic survivorship stress is associated with poor health-related quality of life among distressed survivors of hematopoietic stem cell transplantation. *Psychooncology* 2013;22:911–921.
54. Kent EE, Forsythe LP, Yabroff KR, et al. Are survivors who report cancer-related financial problems more likely to forgo or delay medical care? *Cancer* 2013;119:3710–3717.
55. Ramsey S, Blough D, Kirchhoff A, et al. Washington State cancer patients found to be at greater risk for bankruptcy than people without a cancer diagnosis. *Health Aff (Millwood)* 2013;32:1143–1152.
56. Wan Y, Gao X, Mehta S, et al. Indirect costs associated with metastatic breast cancer. *J Med Econ* 2013;16:1169–1178.
57. Zafar SY, Peppercorn JM, Schrag D, et al. The financial toxicity of cancer treatment: a pilot study assessing out-of-pocket expenses and the insured cancer patient's experience. *Oncologist* 2013;18:381–390.
58. Zullig LL, Peppercorn JM, Schrag D, et al. Financial distress, use of cost-coping strategies, and adherence to prescription medication among patients with cancer. *J Oncol Pract* 2013;9(Suppl 6):60s–63s.
59. Meneses K, Azuero A, Hassey L, et al. Does economic burden influence quality of life in breast cancer survivors? *Gynecol Oncol* 2012;124:437–443.
60. Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
61. Bernard DSM, Farr SL, Fang Z. National estimates of out-of-pocket health care expenditure burdens among nonelderly adults with cancer: 2001 to 2008. *J Clin Oncol* 2011;29:2821–2826.
62. Streeter SB, Schwartzberg L, Husain N, et al. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *Am J Manag Care* 2011;17(Suppl 5):SP38–44.
63. Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
64. de Moor JS, Partridge AH, Winer EP, et al. The role of socioeconomic status in adjustment after ductal carcinoma in situ. *Cancer* 2010;116:1218–1225.
65. Khera N, Chow EJ, Leisenring WM, et al. Factors associated with adherence to preventive care practices among hematopoietic cell transplantation survivors. *Biol Blood Marrow Transplant* 2011;17:995–1003.
66. Markman M, Luce R. Impact of the cost of cancer treatment: an internet-based survey. *J Oncol Pract* 2010;6:69–73.
67. Weaver KE, Rowland JH, Bellizzi KM, et al. Forgoing medical care because of cost: assessing disparities in healthcare access among cancer survivors living in the United States. *Cancer* 2010;116:3493–3504.
68. Pezzin LE, O'Neil MB, Nattlinger AB. The economic consequences of breast cancer adjuvant hormonal treatments. *J Gen Intern Med* 2009;24(Suppl 2):S446–450.
69. Eli K, Xie B, Wells A, et al. Economic stress among low-income women with cancer: effects on quality of life. *Cancer* 2008;112:616–625.
70. Gupta D, Lis CG, Grutsch JF. Perceived cancer-related financial difficulty: implications for patient satisfaction with quality of life in advanced cancer. *Support Care Cancer* 2007;15:1051–1056.
71. Francoeur RB. Cumulative financial stress and strain in palliative radiation outpatients: the role of age and disability. *Acta Oncol* 2005;44:369–381.
72. Langa KM, Fendrick AM, Chernew ME, et al. Out-of-pocket health-care expenditures among older Americans with cancer. *Value Health* 2004;7:186–194.
73. Smith JC, Johnson JT, Cognetti DM, et al. Quality of life, functional outcome, and costs of early glottic cancer. *Laryngoscope* 2003;113:68–76.
74. Houts PS, Harvey HA, Simmonds MA, et al. Characteristics of patients at risk for financial burden because of cancer and its treatment. *J Psychosoc Oncol* 1985;3:15–22.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Shankaran et al, 2018 ¹	Cohort-P	Seattle Cancer Care Alliance (n=23)	<ul style="list-style-type: none"> • Age ≥ 18 y • English-speaking • Diagnosis of nonmetastatic cancer • ≤ 3 y since treatment • All sites 	<ul style="list-style-type: none"> • Mean age, 60 y (range, 32–77 y) • 40% female • 85% white, 5% black, 5% Asian • 50% commercial insurance, 25% Medicare, 25% Medicaid 	<ul style="list-style-type: none"> • Financial hardship and burden • Hardship: debt during treatment (55%), decreased income during treatment (55%), loans during treatment (30%) • Burden: "I feel that my cancer diagnosis and treatment has been a financial burden to my household"; "I feel anxious or worried about how I will manage the costs related to my cancer care"; 48% high anxiety, 42% high burden
Wheeler et al, 2018 ²	Cohort-P	Carolina Breast Cancer Study survey (n=2,494)	<ul style="list-style-type: none"> • Diagnosis of incident breast cancer 	<ul style="list-style-type: none"> • Mean age, 52 y • 51% white, 49% black • 74% private insurance, 6% Medicare, 14% Medicaid, 55% uninsured 	<ul style="list-style-type: none"> • Financial impact • Modified framework of financial impact, based on self-reported survey data at 5 mo (baseline) and 25 mo (follow-up) postdiagnosis, negative impact on OOP and indirect costs (lost insurance, lost job, financial barrier to healthcare, transportation barrier; 48%)
de Souza et al, 2017 ³	Cohort-L	University of Chicago Medicine Multidisciplinary Clinic (n=73)	<ul style="list-style-type: none"> • Diagnosis of stage III–IVb head and neck cancer • Receiving chemoradiation • ECOG 0–1 	<ul style="list-style-type: none"> • Median age, 60 y (range, 26–79 y) • 22% female • 74% white • 55% private insurance, 26% Medicare, 19% Medicaid 	<ul style="list-style-type: none"> • OOP costs • Financial burden: OOP costs relative to household income • Cost-coping: selling possessions/property, using personal savings, borrowing money and taking credit card loans, having family members work more hours to pay for cancer care; 69% used at least one cost-coping strategy
de Souza et al, 2017 ⁴	CS	University of Chicago Medicine, North Shore University (n=233)	<ul style="list-style-type: none"> • Age ≥ 18 y • Diagnosis of stage IV cancer • All sites 	<ul style="list-style-type: none"> • Median age, 59 y (range, 27–88 y) • 58% female • 66% white, 23% black, 6% Hispanic, 4% Asian/Native American • 62% private/employer-based insurance, 31% Medicare, 6% Medicaid 	<ul style="list-style-type: none"> • Financial toxicity • COST tool • Median COST score, 23 (range, 0–44; mean $\pm$ SD, 22.23 $\pm$ 11.89); higher score = lower financial toxicity
Farias & Du, 2017 ⁵	CS	SEER-Medicare database (2007–2009) with Medicare Part D claims (n=8,688)	<ul style="list-style-type: none"> • Women aged ≥ 65 y • Diagnosis of stage I–III hormone receptor-positive breast cancer • Indicated for aromatase inhibitor therapy • ≥ 12 mo in Part A + B Medicare 	<ul style="list-style-type: none"> • Mean age, 75 $\pm$ 7 y (range, 65–103 y) • 83% non-Hispanic white, 6% black, 6% Hispanic, 5% Asian • 56% stage I, 34% stage II, 10% stage III • 60% BCS, 38% mastectomy, 21% chemotherapy, 60% radiation 	<ul style="list-style-type: none"> • Mean OOP costs • 30-d supply of medication amount paid at the time prescription was filled

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCOWFU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Meeker et al, 2017 ⁶	CS	Convenience sample from NCI-designated CCC outpatient clinics (n=119)	<ul style="list-style-type: none"> • Age ≥ 18 y • Diagnosis of cancer, at any point in treatment • All sites 	<ul style="list-style-type: none"> • Median age, 62 y (range, 22–87 y) • 52% female • 84% white, 11% black • 100% insured 	<ul style="list-style-type: none"> • In-Charge Financial Distress/Financial Well-Being Scale • 8-item scale from 1 (overwhelming financial distress) to 10 (highest level of financial well-being); 22% with any financial distress
Pisu et al, 2017 ⁷	Cohort-L	Rural Breast Cancer Survivor trial (n=432)	<ul style="list-style-type: none"> • Age ≥ 21 y • Diagnosis of stage 0–III breast cancer • ≤ 3 y since primary treatment • Resident of rural Florida 	<ul style="list-style-type: none"> • 4% aged 35–45 y, 49% aged 46–64 y, 48% aged 65–90 y • 6% minority race/ethnicity • 94% insured • 66% hormonal therapy, 42% surgery, chemotherapy, and radiation 	<ul style="list-style-type: none"> • OOP costs • Work and finances inventory questionnaire • Costs for medical care, counseling and health maintenance, side-effect management, home maintenance, and other • Measured at baseline and 3, 6, 9, and 12 mo • Mean OOP monthly costs were higher at baseline compared with 12 mo (mean ratio, 1.37; 95% CI, 1.13–1.66; $P=.007$); no difference between 3, 6, 9, and 12 mo
Abel et al, 2016 ⁸	CS	Dana-Farber Cancer Institute, Mayo Clinic Arizona, Roswell Park Cancer Institute (n=325)	<ul style="list-style-type: none"> • Age ≥ 18 y • English-speaking • ≥ 150 d post-HCT 	<ul style="list-style-type: none"> • 49% aged ≤ 60 y, 51% aged >60 y • 40% female • 92% white, 8% nonwhite • 62% employer-based insurance, 30% government insurance, 8% self-insured • 53% received autologous, 47% received allogeneic HCT • 32% multiple myeloma, 24% NHL, 17% AML 	<ul style="list-style-type: none"> • Financial distress at 6 mo from transplant • Hardship: "How satisfied are you with your family's present financial situation?" (49% unsatisfied); "How difficult is it to meet monthly payments?" (42% faced difficulties); "How do your family's finances work out at the end of the month?" (19% did not have enough money); any hardship, 56%; all 3 hardships, 15% • Difficulty: with temporary relocation, 19%; transit, gas, parking, 41%; changes at home, such as cleaning and special foods, 36%; any difficulty, 51%
Banegas et al, 2016 ⁹	Cohort-R	2012 LIVESTRONG Survey (n=4,719)	<ul style="list-style-type: none"> • Age 18–64 y • Diagnosis of cancer, at any point in treatment • All sites 	<ul style="list-style-type: none"> • 32% aged 18–44 y, 34% aged 45–54 y, 34% aged 55–64 y • 67% female • 88% non-Hispanic white, 2% black, 6% Hispanic • 85% private insurance, 8% public insurance, 3% uninsured 	<ul style="list-style-type: none"> • Financial hardship • Borrowed money or went into debt (34%), went into bankruptcy (3%), worried about paying bills (64%), made financial sacrifices (40%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for Financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Chongpison et al, 2016 ¹⁰	Cohort-P	Kaiser Permanente, North California or Oregon/Southwest Washington, 2010–2011 (n=576)	<ul style="list-style-type: none"> ≥5 y from diagnosis of rectal cancer as of 2010 Received a permanent ostomy or an anastomosis ± a temporary stoma 	<ul style="list-style-type: none"> 41% female 83% white, 4% black, 8% Asian, 6% other races; 94% non-Hispanic ethnicity, 6% Hispanic 	<ul style="list-style-type: none"> Perceived financial burden The City of Hope Quality of Life Colorectal Cancer scale: “How much financial burden currently resulted from your illness and treatment?” Mean financial burden score, 2±2.8 on a scale of 0–10; 22% perceived current financial burden as moderate to high (≥4 points)
Delgado-Guay et al, 2016 ¹¹	CS	MD Anderson Cancer Center, October 2012–January 2013 (n=292)	<ul style="list-style-type: none"> Diagnosis of advanced cancer All sites 	<ul style="list-style-type: none"> Median age, 61 y (range, 22–92 y) 47% female 65% white, 16% black, 12% Hispanic, 5% Asian 53% metastasis, 41% local/recurrent disease, 6% no evidence of active disease 	<ul style="list-style-type: none"> Financial distress Edmonton Symptom Assessment Scale Distress/Suffering secondary to financial issues
DiMartino et al, 2017 ¹²	CS	2011 MEPS, ECSS (n=1,320)	<ul style="list-style-type: none"> Age ≥18 y All sites 	<ul style="list-style-type: none"> 23% aged <55 y, 24% aged 55–64 y, 27% aged 65–74 y, 26% aged ≥75 y 56% female 90% non-Hispanic white, 10% minority 72% private insurance, 24% public insurance, 4% uninsured 	<ul style="list-style-type: none"> Financial hardship Ever unable to cover share of costs of medical visits due to cancer (12%)
Farias et al, 2018 ¹³	CS	Truven Health Analytics MarketScan commercial claims and encounters database, January 2008–December 2010 (n=6,863)	<ul style="list-style-type: none"> Age ≤64 y Diagnosis of EBC Breast cancer surgery ≤12 mo ≥1 prescription for AI or tamoxifen 	<ul style="list-style-type: none"> 8% aged 18–39 y, 33% aged 40–49 y, 41% aged 50–59 y, 17% aged 60–64 y 18% HMO, 60% PPO, 22% other insurance 33% mastectomy, 67% BCS 	<ul style="list-style-type: none"> OOP costs 30-d supply of medications including copayments (accounted for 89% toward OOP), deductibles (6%), and coinsurance (5%); mean, \$17.10 Other OOP costs included inpatient/outpatient services and other pharmacy medications (mean, \$176.50)
Kale & Carroll, 2016 ¹⁴	CS	2011 MEPS (n=380)	<ul style="list-style-type: none"> Diagnosis of cancer All sites 	<ul style="list-style-type: none"> 72% aged <65 y 56% female 88% non-Hispanic white, 5% Hispanic, 5% black 71% private insurance, 25% public insurance, 92% any insurance 	<ul style="list-style-type: none"> Cancer-related financial burden Problem: loans, debt, bankruptcy; financial sacrifices; worry about paying large medical bills; inability to cover cost of medical care visits 29% had ≥1 problem, 2% had all 4 problems
Keim-Malpass et al, 2016 ¹⁵	CS	Memorial Sloan Kettering Cancer Center and University of Texas–Southwestern Center for Breast Care, April 2003–February 2006 (n=258)	<ul style="list-style-type: none"> Age ≥18 y Diagnosis of stage I–III breast cancer within ≤8 mo Fluent in English 	<ul style="list-style-type: none"> Mean age, 49.9±9.62 y 95% white/Caucasian, 5% nonwhite 50% stage I, 41% stage II, 9% stage III 74% chemotherapy 	<ul style="list-style-type: none"> Workplace perception Scale from 1 (not at all) to 5 (very much) Supportive work environment (mean, 3.92±1.21); ability to keep up work responsibilities (mean, 4.51±0.81) Financial problems: financial worries (mean, 2.10±1.22); difficulty paying for basics (16%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCC, basal cell carcinoma; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for Financial Toxicity, CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHL, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Lathan et al, 2016 ¹⁶	Cohort-P	CanCORS Consortium, 2003–2005 (2,434 patients with lung cancer, 2,909 with CRC)	<ul style="list-style-type: none"> Patients or surrogates of patients from CanCORS study 4–12 mo from diagnosis 	<ul style="list-style-type: none"> Age <59 y at diagnosis: 27% (lung cancer) and 38% (CRC) 44% female both lung cancer and CRC Patients who were white: 74% (lung cancer) and 65% (CRC) 	<ul style="list-style-type: none"> Patient-reported financial reserves “if you lost all of your current sources of income and had to live off savings, how long could you continue to live at your current address and standard of living?” (<1 mo, 1–2 mo, 3–6 mo, 7–12 mo, >1 y, do not know) 40% patients lung cancer and 33% with CRC had financial reserves (<2 mo)
Lee & Khan, 2016 ¹⁷	CS	NHIS, 2006–2013 (n=15,159)	<ul style="list-style-type: none"> Age ≥20 y All sites 	<ul style="list-style-type: none"> 8% aged <40 y, 39% aged 40–64 y, 35% aged 65–79 y, 18% aged ≥80 y 62% female 77% non-Hispanic white, 12% black, 8% Hispanic, 3% Asian 28% private insurance, 52% Medicare, 4% Medicaid, 9% government insurance, 7% uninsured 	<ul style="list-style-type: none"> CRN “During the past 12 mo, was there a time when you did not take your prescribed medication because of cost? Do not include OTC medications”; 11% reported CRN
Lee & Salloum, 2016 ¹⁸	CS	NHIS, 2006–2013 (n=10,998)	<ul style="list-style-type: none"> Age ≥18 y All sites At least 1 y from diagnosis 	<ul style="list-style-type: none"> 10% aged <40 y, 47% aged 40–64 y, 35% aged 65–79 y, 8% aged ≥80 y 58% female 79% non-Hispanic white, 12% black, 9% Hispanic 34% private insurance, 5% Medicaid, 43% Medicare, 10% government insurance, 8% uninsured 	<ul style="list-style-type: none"> CRN “During the past 12 mo, was there a time when you did not take your prescribed medication because of cost? Do not include OTC medications”; 13% reported CRN
Meeker et al, 2016 ¹⁹	CS	Fox Chase Cancer Center, September 2013–April 2014 (n=120)	<ul style="list-style-type: none"> Age >18 y Received anticancer medications All sites Insured 	<ul style="list-style-type: none"> Median age, 62 y (range, 22–87 y) 52% female 84% white, 11% black, 3% Asian, 1% Hispanic, 1% other 68% surgery, 67% chemotherapy, 39% radiation, 16% targeted therapy, 15% hormone therapy 	<ul style="list-style-type: none"> Financial concerns InCharge Financial Distress/Financial Well-Being Scale Concerns: need to continue to work to pay for treatment (40%), paying for treatment (43%), paying other bills (41%) Most frequent assistant needs: knowing whom to ask for help (29%), identifying sources of financial assistance (26%), understanding the insurance policy (24%), and accessing community resources (21%) Financial distress: InCharge scale; 29% had high to overwhelming financial distress Overall distress: score of ≥4 (65%), ≥8 (28%) Emotional distress: ≥1 emotional symptom (66%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWFU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Narang & Nicholas, 2017 ²⁰	CS	Health and Retirement Study, July 2002–December 2012 (n=1,409)	<ul style="list-style-type: none"> • Age >65 y • New cancer diagnosis • All sites 	<ul style="list-style-type: none"> • Median age, 73 y (interquartile range, 69–79 y) • 46% female • 91% white, 8% black • 8% Medicaid, 8% VA, 18% Medicare HMO, 28% employer-sponsored, 25% Medigap, 13% uninsured 	<ul style="list-style-type: none"> • Financial burden • Calculated as OOP expenses divided by total household income
Nipp et al, 2016 ²¹	CS	Health Well Foundation (financial assistance program), June 2010–May 2011 (n=174)	<ul style="list-style-type: none"> • Age ≥21 y • On active chemotherapy or hormonal therapy for solid tumor malignancy • All sites 	<ul style="list-style-type: none"> • Median age, 67 y • 96% female • 83% Caucasian • 69% Medicare, 3% Medicaid, 16% employer-provided, 20% personal insurance, 35% personal supplemental, 20% other • 85% breast cancer, 4% CRC • 69% metastasis 	<ul style="list-style-type: none"> • Financial burden • Care altering coping: did not fill prescription (28%), took less medication (22%), missed a test (10%), missed a procedure (8%), missed an appointment (6%) • Lifestyle altering coping: reduced spending on leisure (78%), reduced spending on basics (57%), borrowed money to pay for medications (54%), used savings (50%), sold possessions (18%), family worked more (15%)
Ramsey et al, 2016 ²²	Cohort-R	CSS of Western Washington linked with federal bankruptcy records, 1995–2009 (n=231,596 [original], n=4,728 [propensity-matched])	<ul style="list-style-type: none"> • Age ≥21 y • Diagnosis of primary cancer • All sites 	<ul style="list-style-type: none"> • Mean age, 53 y • 46% female • 86% white, 14% nonwhite 	<ul style="list-style-type: none"> • Financial insolvency • Filed for bankruptcy after cancer diagnosis (2%)
Whitney et al, 2016 ²³	CS	2011 MEPS (n=1,209)	<ul style="list-style-type: none"> • Age ≥18 y • Completed ECSS survey • All sites 	<ul style="list-style-type: none"> • 46% aged <65 y, 54% aged ≥65 y • 58% female • 84% non-Hispanic white, 16% minority • 4% uninsured, 27% public insurance, 68% private insurance 	<ul style="list-style-type: none"> • Self-reported financial concerns: financial difficulty: Debt, bankruptcy, inability to cover medical costs (18%); financial worry: Financial sacrifices or worry about bills (28%) • 33% ≥1 of above concerns • Self-reported work concerns: long-term work modification: Changing to a less demanding job, delaying, or taking early retirement (15%); short-term work modification: extended/unpaid leave from work (39%) • 44% ≥1 of above concerns

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWFU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBs, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHL, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Wong et al, 2016 ²⁴	CS	NCI cancer centers in Ohio, Pennsylvania, Illinois, and Michigan, April 2010–September 2012 (n=1,233)	<ul style="list-style-type: none"> • Diagnosis of cancer • All sites 	<ul style="list-style-type: none"> • 73% aged ≤65 y, 27% aged >65 y • 58% female • 88% white, 12% nonwhite • 42% metastasis 	<ul style="list-style-type: none"> • Financial concerns • a = cost-related burden; b + c = uncertainty; a + b + c = total score (range, 0–15); higher score indicates greater concerns • (a) How much of a burden on you is the cost of your medical care? (22% major or extreme burden); (b) I am afraid that my health insurance will not pay for a clinical trial (40% agreed); (c) I am worried that I would not be able to afford the cost of treatment for a clinical trial (30% agreed)
Yabroff et al, 2016 ²⁵	CS	2011 MEPS, ECSS (n=1,202)	<ul style="list-style-type: none"> • Age ≥18 y • All sites 	<ul style="list-style-type: none"> • 53% aged ≥65 y • 61% female • 76% non-Hispanic white, 24% all other 	<ul style="list-style-type: none"> • Financial hardship • 20% material hardship (loans, debt, bankruptcy, inability to cover medical costs, financial sacrifices) (20%); 23% psychological hardship (worry about paying large medical bills)
Zheng et al, 2015 ²⁶	CS	MEPS household component, 2008–2012 (n=3,278)	<ul style="list-style-type: none"> • Age ≥18 y • CRC, female breast cancer, and male prostate cancer survivors vs adults without cancer 	<ul style="list-style-type: none"> • 55% CRC, 57% female • 72%–87% non-Hispanic white • 16% CRC, 48% breast cancer, 36% prostate cancer among all cancer survivors 	<ul style="list-style-type: none"> • Economic burden • Medical expenses: for inpatient/outpatient, home healthcare, dental care, vision care, prescriptions; examined by source of payment (OOP, insurance) and by type of service (ambulatory, inpatient) • Productivity losses: employment disability, missed days at work, days spent in bed, reason for not working
Callahan & Brintzenhofesoc, 2015 ²⁷	CS	University of Maryland, Baltimore, and The Catholic University of America, May 2010–October 2011 (n=90)	<ul style="list-style-type: none"> • Diagnosis of cancer • All sites 	<ul style="list-style-type: none"> • 60% female • 35% African American, 53% Caucasian, 6% multiracial, 7% other race • 83% insured 	<ul style="list-style-type: none"> • Financial quality of life • Ability to manage all current and future financial obligations related to cancer care, within the context of sound healthcare decision-making • Socioeconomic Well-Being Scale
Delgado-Guay et al, 2015 ²⁸	CS	CCC, GPH in Texas (n=149)	<ul style="list-style-type: none"> • Age >18 y • Normal cognition • Diagnosis of advanced breast, colon, lung, or prostate cancer 	<ul style="list-style-type: none"> • Mean age, 60 y • 50% female • 41% white, 36% black, 23% Hispanic • 81% metastasis 	<ul style="list-style-type: none"> • Objective financial burden: % of family income spent on OOP expenses on healthcare during the last month, without considering health insurance • Subjective financial distress: distress due to financial burden within the past month (concerns, difficulties, worries)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBs, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasm; MEPS, Medical Expenditure Panel Survey; NHL, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Guy et al, 2015 ²⁹	CS	2008–2012 MEPS (n=4,271)	<ul style="list-style-type: none"> Age 18–64 y Cancer survivors (all sites) vs adults without cancer 	<ul style="list-style-type: none"> 65% aged 50–64 y 65% female 81% non-Hispanic white, 8% black, 7% Hispanic 74% private insurance, 15% public insurance, 11% uninsured 	<ul style="list-style-type: none"> OOP expense burden % OOP costs relative to family income High OOP = expenses ≥20% of family income (4%)
Huntington et al, 2015 ³⁰	CS	Tertiary academic medical center in Philadelphia, PA, August 2014–January 2015 (n=100)	<ul style="list-style-type: none"> Diagnosis of multiple myeloma ≥3 mo ongoing treatment 	<ul style="list-style-type: none"> Mean age, 64 y 53% female 71% white, 26% black 43% commercial insurance, 39% Medicare fee-for-service with supplement, 10% Medicare HMO 	<ul style="list-style-type: none"> Financial toxicity COST tool and patient self-report Mean COST score, 23±11.1; score ≤23 = higher financial burden Self-reported significant financial burden (19%) Self-reported higher than expected treatment costs (59%)
Jones et al, 2015 ³¹	CS	Colorado Central Cancer Registry, May 2002–August 2006 (n=477)	<ul style="list-style-type: none"> Age 18–85 y, resident of Colorado Diagnoses of leukemia or lymphoma Completed treatment within past 3–48 mo 	<ul style="list-style-type: none"> 15% aged 18–39 y, 53% aged 40–64 y, 31% aged 65–85 y 46% female 91% white, 5% Hispanic, 2% black 60% NHL, 19% Hodgkin lymphoma, 5% AML, 2% ALL, 6% CML, 8% CLL 94% chemotherapy, 5% bone marrow transplant 	<ul style="list-style-type: none"> Perceived financial burden “How much financial burden have you incurred because of illness and treatment?” (0 [none]–10 [a great deal]) Mean perceived financial burden, 4.2±3.6
Palmer et al, 2015 ³²	CS	NCI-CCCFWU, January 2009–November 2011 (n=191)	<ul style="list-style-type: none"> Diagnosis of first primary breast cancer, nonmetastasis Completed treatment within past 6–24 mo 	<ul style="list-style-type: none"> 22% aged <50 y, 35% aged 50–59 y, 27% aged 60–69 y, 16% aged ≥70 y 84% white, 16% black 61% private insurance, 8% public insurance, 25% both private and public insurance, 6% uninsured 	<ul style="list-style-type: none"> Cost-related barriers to follow-up care OOP medical costs are too high (32%); insurance does not cover it (22%); cannot afford other costs to receive care (eg, gas/travel or lost wages) (25%)
Pisu et al, 2015 ³³	CS	CanCORS Consortium after 2003 (n=3,342 [1,364 with lung cancer, 2,068 with CRC])	<ul style="list-style-type: none"> Diagnosis of lung cancer or CRC in past 4–12 mo 	<ul style="list-style-type: none"> For lung and CRC patients, respectively: <ul style="list-style-type: none"> Age <54 y at diagnosis: 13%, 24% Female: 49%, 46% White: 83%, 76%; black: 12%, 15%; Hispanic: 5%, 8% Uninsured: 1%, 3%; no prescription coverage: 14%, 10% Stage I: 42%, 26% 	<ul style="list-style-type: none"> Self-reported economic hardship Any economic hardship (53% lung cancer, 46% CRC) Difficulty living on current income (extremely difficult: 4% lung cancer, 2% CRC); anticipating hardship (a great deal: 4% lung cancer, 4% CRC); anticipating reductions in living (a great deal: 6% lung cancer, 5% CRC)
Tucker-Seeley et al, 2015 ³⁴	CS	Multisite, September 2002–February 2008 (n=281)	<ul style="list-style-type: none"> Age ≥20 y Advanced cancer with metastasis All sites 	<ul style="list-style-type: none"> Mean age, 59 y 46% female 66% white, 34% black 63% insured, 37% uninsured 	<ul style="list-style-type: none"> Financial hardship Household had to use all or most of savings because of family member's illness (29%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWFU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasm; MEPS, Medical Expenditure Panel Survey; NHL, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Zafar et al, 2015 ³⁵	CS	CanCORS Consortium (n=1,000)	<ul style="list-style-type: none"> Diagnosis of lung cancer or CRC Survivors/Disease-free or presence of recurrent/advanced disease 	<ul style="list-style-type: none"> Median age, 64 y (range, 52–82 y) 47% female 76% white, 24% nonwhite 81% private insurance, 50% Medicare, 4% Medicaid, 12% VA/military, 2% other 27% lung cancer, 73% CRC 89% cancer-free, 11% recurrent 	<ul style="list-style-type: none"> Financial burden Difficulty living on household income (48%) Quality of insurance coverage
Bender et al, 2014 ³⁶	Cohort-L	Comprehensive Breast Program, University of Pittsburgh Cancer Institute, December 2008–March 2010 (n=91)	<ul style="list-style-type: none"> Age 18–75 y Hormone-positive EBC (stage I, II, or IIIa) Received endocrine therapy Completed ≥8 y education, fluent in English 	<ul style="list-style-type: none"> Mean age, 56.7 ± 9.7 y 97% Caucasian 59% stage I, 41% stage II/III 23% chemotherapy 	<ul style="list-style-type: none"> Financial hardship OOP costs (COIN instrument) ($\bar{x}$ = 307.5 ± 734.2) Measure of economic hardship across 4 domains: financial strain ($\bar{x}$ = 2.3 ± 0.99), inability to make ends meet ($\bar{x}$ = 6.9 ± 0.98), not enough money for necessities ($\bar{x}$ = 12.8 ± 5.5), cutbacks and adjustments ($\bar{x}$ = 17.4 ± 0.95)
Bestvina et al, 2014 ³⁷	CS	Duke Cancer Institute and affiliated clinics, November 2012–June 2013 (n=300)	<ul style="list-style-type: none"> Diagnosis of solid tumor malignancy All sites Had insurance ≥1 mo anticancer therapy 	<ul style="list-style-type: none"> Median age, 60 y 48% female 76% white, 24% nonwhite 56% private insurance, 44% nonprivate insurance 	<ul style="list-style-type: none"> Financial burden Objective financial burden: if patients discussed OOP costs with their oncologist Subjective financial distress: InCharge financial distress/well-being scale 16% had high/overwhelming financial distress, 39% had higher than expected financial burden
de Souza et al, 2014 ³⁸	Item and exploratory factor analysis (COST tool)	University of Chicago, January 2012–August 2013 (n=155)	<ul style="list-style-type: none"> Diagnosis of advanced cancer Receiving treatment ≥3 mo All sites 	<ul style="list-style-type: none"> Median age, 59.5 y (range, 24–84 y) 45% female 74% non-Hispanic white, 18% black, 6% Hispanic, 2% other 62% private or employer insurance, 32% Medicare, 4% Medicaid 	<ul style="list-style-type: none"> Financial toxicity COST tool Themes: resources, personal affect, financial coping, family support Median COST score, 21 (range, 3–44; mean, 22.5 ± 11.3) Higher score = lower financial toxicity
Dusetzina et al, 2014 ³⁹	Cohort-R	Truven Health MarketScan database, 2002–2011 (n=1,541)	<ul style="list-style-type: none"> Age 18–64 y Diagnosis of CML Have insurance from 3 mo before and 6 mo after initiation of imatinib 	<ul style="list-style-type: none"> Mean age, 48.8 y (SD, 11.0) 44% female 60% PPO insurance 	<ul style="list-style-type: none"> Prescription copayment Required for a 30-d supply of imatinib 88.5% paid any copayment (excluding deductible) for this supply

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute myeloid leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWFO, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHL, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Ekwueme et al, 2014 ⁴⁰	CS	2008–2011 MEPS and 2011 MEPS Cancer Survivorship Survey (n=6,722 and n=1,202, respectively)	<ul style="list-style-type: none"> Cancer survivors (all sites) vs adults without cancer 	<ul style="list-style-type: none"> 18% aged ≥80 y 58% female 85% white, 7% black, 5% Hispanic, 3% other 75% private insurance (age <65 y), 55%–63% Medicare and private insurance (age ≥65 y) 	<ul style="list-style-type: none"> Lost work productivity Employment disability (unable to work because of illness/injury); missed worked days, days spent in bed
Fenn et al, 2014 ⁴¹	CS	NHIS 2010 data (n=2,108)	<ul style="list-style-type: none"> Diagnosis of cancer All sites 	<ul style="list-style-type: none"> 53% aged 18–64 y, 47% aged ≥65 y 56% female 90% white, 7% black, 2% Asian 51% Medicare, 37% private insurance 	<ul style="list-style-type: none"> Financial problems "To what degree has cancer caused financial problems for you and your family?": 9% reported "a lot," 12% reported "some," 10% reported "a little," 70% reported "not at all"
Jagsi et al, 2014 ⁴²	Cohort-R	SEER data for Los Angeles, CA, and Detroit, MI, survey, June 2005–February 2007 (n=746)	<ul style="list-style-type: none"> Age 20–64 y Diagnosis of stage 0–III nonrecurrent breast cancer 	<ul style="list-style-type: none"> 26% aged <46 y, 42% aged 46–55 y, 33% aged ≥56 y 42% white, 18% black, 38% Latina 19% stage 0, 33% stage I, 37% stage II, 12% stage III 61% chemotherapy, 61% BCS, 69% radiation 	<ul style="list-style-type: none"> Long-term work loss among those working at the time of diagnosis: 34% with 4-year unemployment; of those, 50% reported that it was important for them to work and 31% were actively seeking work
Jagsi et al, 2014 ⁴³	Cohort-R	SEER data for Los Angeles, CA, and Detroit, MI, survey, June 2005–February 2007 (n=1,502)	<ul style="list-style-type: none"> Age 20–79 y Diagnosis of stage 0–III breast cancer 	<ul style="list-style-type: none"> 21% aged <46 y, 27% aged ≥65 y 43% white, 17% black, 20% English-speaking Latina, 20% Spanish-speaking 18% stage 0, 34% stage I, 33% stage II, 12% stage III 52% chemotherapy, 62% BCS, 67% radiation 	<ul style="list-style-type: none"> Financial impact Financial decline: worse off financially since diagnosis (33% any decline) OOP (17% >\$5,000) Medical debt 4 y after diagnosis (12%) Decline in employment status (15%)
Khera et al, 2014 ⁴⁴	CS	Mayo Clinic in Arizona, Rochester, or Jacksonville, January 2006–June 2012 (n=268)	<ul style="list-style-type: none"> Received allogeneic HCT 	<ul style="list-style-type: none"> Median age, 55 y (range, 20–71 y) 49% female 92% non-Hispanic white, 5% nonwhite, 3% Hispanic 59% private insurance, 26% Medicare + supplement, 7% Medicare only 35% AML, 21% MDS/MPN, 17% NHL/CLL, 14% ALL 	<ul style="list-style-type: none"> Subjective financial burden: overall 80%; being sick has hurt me financially (73% agreed); have to pay more than I can afford (33% agreed); medicines are too expensive (30% agreed); I have enough money to take care of my health (32% disagreed); I can pay my medical bills (22% disagreed); I can make enough money to pay for my healthcare (35% disagreed) Objective financial burden: overall 47%; household income decreased by more than 50% (26% reported yes); had to refinance/second mortgage home (8%); had to sell home (1%); declared bankruptcy (3%); withdrew money from retirement (25%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCQWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Kircher et al, 2014 ⁴⁵	CS	MEPS data, 2002–2005 and 2007–2010 (n=6,607 [1,878 near-elderly, 4,729 with Medicare])	<ul style="list-style-type: none"> • Age ≥55 y • Diagnosis of cancer • All sites 	<ul style="list-style-type: none"> • In Medicare group: mean age, 75.3 y; 46% female; 94% white, 4% black; 62% private insurance, 38% public insurance • In near-elderly group: mean age, 59.1 y; 53% female; 92% white, 5% black; 91% private insurance, 1% public insurance 	<ul style="list-style-type: none"> • OOP cost burden • Direct payment for all prescription drugs
Martin et al, 2014 ⁴⁶	CS	CanCORS Consortium, 2003 (n=5,044)	<ul style="list-style-type: none"> • Diagnosis of lung cancer or CRC • ≈4 mo from diagnosis 	<ul style="list-style-type: none"> • 79% aged >55 y • 47% female • 74% white • 97% insured 	<ul style="list-style-type: none"> • Worry • Treatment side effects (75%); cost of treatment (40%); time away from family (50%); time away from work (52%); transportation to treatment (22%)
Regenbogen et al, 2014 ⁴⁷	CS	SEER registry of metropolitan Detroit, MI, and of the state of Georgia, August 2011–March 2013 (n=937)	<ul style="list-style-type: none"> • Age ≥21 y • Pathologic stage III colon or rectal cancer • 3–12 mo postsurgery 	<ul style="list-style-type: none"> • 16% aged <50 y, 37% aged 50–64 y, 23% aged 65–74 y, 24% aged ≥75 y • 46% female • 72% white, 22% black • 42% private insurance, 45% Medicare, 4% Medicaid • 85% had chemotherapy 	<ul style="list-style-type: none"> • Personal financial burden: global financial burden (illness impacted my finances) (62%); financial worry measured using a Likert scale dichotomized later into “low” vs “high” • Composite financial burden score range, 0–7; higher scores = increased financial burden; 62% endorsed no measures of financial burden
Veenstra et al, 2014 ⁴⁸	CS	SEER registry of metropolitan Detroit, MI, and of the state of Georgia, August 2011–March 2013 (n=956)	<ul style="list-style-type: none"> • Age ≥21 y • Pathologic stage III colon or rectal cancer • 3–12 mo postsurgery 	<ul style="list-style-type: none"> • 17% aged <50 y, 37% aged 50–64 y, 23% aged 65–74 y, 24% aged >75 y • 46% female • 71% white, 23% black • 38% Medicare, 33% employer insurance, 3% Medicaid 	<ul style="list-style-type: none"> • Personal financial burden: global financial burden (illness impacted my finances) (61%); financial worry measured using a Likert scale dichotomized later into “low” vs “high” • Mean financial burden score was 1.72±1.83 (range, 0–6; higher scores denoted increased financial burden)
Blinder et al, 2013 ⁴⁹	Cohort-L	California's BCCTP (n=274)	<ul style="list-style-type: none"> • >6 mo from diagnosis of stage 0–III breast cancer • Curative treatment • Not undergoing treatment • Enrolled in BCCTP (family income ≤200% poverty level) • English-/Spanish-speaking 	<ul style="list-style-type: none"> • Median age, 49 y (range, 26–85 y) • 53% Latina, 33% non-Latina, 14% other • 10% DCIS, 31% stage I, 44% stage II, 14% stage III • 39% mastectomy, 61% BCS, 16% reconstruction, 70% chemotherapy, 69% endocrine therapy, 73% radiation 	<ul style="list-style-type: none"> • Financial status • Perceived adequacy of financial resources • 62% reported inadequacy of financial resources to meet needs

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasm; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Davidoff et al, 2013 ⁵⁰	CS	MCBS, 1997–2007 (n=1,868)	<ul style="list-style-type: none"> • Diagnosis of cancer • All sites 	<ul style="list-style-type: none"> • 6% aged <65 y • 49% female • 82% white, 9% black, 6% Hispanic • 37% private/employer insurance • 36% surgery, 22% radiation, 31% antineoplastic 	<ul style="list-style-type: none"> • Total OOP spending • High spending was >20% of income • Total OOP/income: mean, 23.9 (median, 10) for patients with cancer • OOP/income of >10% among 50% of patients and >20% among 28% of patients
Dowling et al, 2013 ⁵¹	CS	MEPS survey, 2008–2010 (n=4,960)	<ul style="list-style-type: none"> • Age ≥18 y • Patients with cancer compared with adults without cancer • All sites 	<ul style="list-style-type: none"> • 7% aged 18–39 y, 27% aged 40–59 y, 47% aged 60–79 y, 18% aged ≥80 y • 59% female • 85% non-Hispanic white, 7% black, 5% Hispanic, 3% other/multiple 	<ul style="list-style-type: none"> • Lost productivity • Lacked employment in past 12 mo (for age <65 y) (25%); limitation at work, housework, or school (12%); inability to do any activities (7%); experienced cognitive limitations (6%)
Goodwin et al, 2013 ⁵²	CS	Myeloma Institute for Research and Therapy (n=1,015)	<ul style="list-style-type: none"> • Diagnosis of multiple myeloma 	<ul style="list-style-type: none"> • Mean age, 61 y (SD, 9.26) • 39% female • 90% white, 6% black, 1% Asian, 1% American Indian/Alaskan Native • 3% Hispanic/Latino 	<ul style="list-style-type: none"> • Financial effects of multiple myeloma: unemployment, disability, health/medical/life insurance, retirement or OOP expenses • Burden of treatment costs/financial hardship: abrupt stoppage of insurance, inability to obtain other insurance, income spent on deductibles, copayments, prescriptions, healthcare, temporary travel, lodging (42% somewhat to very much of a burden) • OOP expenses: 36% of income spent in first 12 mo of treatment and 28% during past 12 mo prior to survey
Hamilton et al, 2013 ⁵³	CS	Mount Sinai Medical Center and Hackensack University Medical Center (n=181)	<ul style="list-style-type: none"> • Age ≥18 y • HCT within past 9–36 mo • English-speaking • With telephone access • Not currently relapsed • ≥ moderately distressed 	<ul style="list-style-type: none"> • 55% female • 87% white/non-Hispanic • 99.4% had insurance during transplant with 59% receiving full coverage and 36% >half coverage of transplant costs 	<ul style="list-style-type: none"> • Economic survivorship stressors • Financial (reducing vacations, leisure activities, spending on food/clothing, delaying purchases); possible range of score: 0–8.1; mean, 1.81±1.85 • Employment (pay cut, loss of income, disability, leave of absence); possible range of score: 0–38; mean, 6.28±6.87; • Insurance (time spent dealing with insurance company and insurance denying a payment); possible range of score: 0–20; mean, 3.43±4.68

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Kent et al, 2013 ⁵⁴	CS	2010 NHIS data with multistage sampling (n = 1,556)	<ul style="list-style-type: none"> • Diagnosis of cancer at age >21 y • All sites 	<ul style="list-style-type: none"> • 51% aged 40–64 y, 30% aged ≥65 y • 58% female • 83% non-Hispanic white, 8% black, 6% Hispanic 	<ul style="list-style-type: none"> • Cancer-related financial problems • “To what degree has cancer caused financial problems for you and your family?” • 32% reported some cancer-related financial problems
Ramsey et al, 2013 ⁵⁵	Cohort-R	SEER data, western Washington State linked with U.S. Bankruptcy Court, January 1995–December 2009 (n = 197,840)	<ul style="list-style-type: none"> • Age ≥21 y and <90 y • Diagnosis of cancer • Not an in-situ stage • All sites 	<ul style="list-style-type: none"> • 7% aged <40 y, 49% aged 40–64 y, 44% aged ≥65 y • 50% female • 88% white, 12% nonwhite • 50% localized cancer, 22% regional cancer, 24% distant cancer, 4% unstaged cancer 	<ul style="list-style-type: none"> • Filing for bankruptcy (overall 2.2%) • US Bankruptcy Code Chapter 7: debtors liquidate assets (bank accounts, investments, cars, homes) to pay creditors to get discharged from debts excluding child or spousal support and income tax (of those overall, 83% under this code). • U.S. Bankruptcy Code Chapter 13: Debtors file for repayment plan to pay back debts and retain ownership of assets (of those overall, 17% under this code).
Wan et al, 2013 ⁵⁶	Cohort-R	Marketscan commercial claims and encounters and health and productivity management databases (n = 571)	<ul style="list-style-type: none"> • Age 18–64 y • Diagnosis of breast cancer compared with adults without cancer 	<ul style="list-style-type: none"> • Mean age, 49 ± 7 y • 47%–51% POS insurance, 8%–14% HMO, 36%–38% PPO • 76% EBC, 24% MBC 	<ul style="list-style-type: none"> • Indirect costs • Paid time off including absenteeism (or sick leave) • Short-term disability associated costs
Zafar et al, 2013 ⁵⁷	Cohort-L	Health Well Foundation and Duke University Medical Center, June 2010–May 2011 (n = 254)	<ul style="list-style-type: none"> • Diagnosis of solid tumor • Receiving chemotherapy or endocrine therapy • All sites 	<ul style="list-style-type: none"> • Mean age, 64 y • 86% female • 80% white, 13% black, 1% Asian • 59% Medicare, 81% drug coverage • 71% breast cancer diagnosis • 67% ≥12 mo chemotherapy 	<ul style="list-style-type: none"> • Financial burden • Objective financial burden (OOP expenses) • Subjective financial burden (patient self-report): 93% reported any burden, 33% reported significant burden, 9% reported catastrophic cancer-related financial burden from OOP expenses
Zullig et al, 2013 ⁵⁸	CS	Health Well Foundation and Duke University Medical Center, June 2010–May 2011 (n = 164)	<ul style="list-style-type: none"> • Age ≥21 y • Diagnosis of solid tumor • Receiving chemotherapy or endocrine therapy • All sites 	<ul style="list-style-type: none"> • Mean age, 63–70 y • 97% female • 83% white, 10% black, 4% other • 78% prescription drug coverage, 73% Medicare/Medicaid, 16% employer insurance, 20% purchased insurance personally • 85% breast cancer, 4% CRC, 10% other • 72% ≥12 mo chemotherapy 	<ul style="list-style-type: none"> • Subjective financial distress • 5-point Likert scale: “not a financial burden at all” to “catastrophic financial burden” • 47% reported significant or catastrophic financial distress

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for Financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Meneses et al, 2012 ⁵⁹	Cohort-L	BCEI study (n=132)	<ul style="list-style-type: none"> • Age ≥21 y • Stage 0–II breast cancer • <2 y from diagnosis • >1 mo since initial treatment • No recurrence or metastasis 	<ul style="list-style-type: none"> • 58% aged 46–65 y, 22% aged 66–83 y • 81% Caucasian • 57% stage I 	<ul style="list-style-type: none"> • Economic burden • BCEI survey (at 3 mo, 6 mo) • Work changes in motivation, productivity, quality, quantity): 17%–38% • Financial hardship (changes in income, applying for unemployment benefits, finding second jobs, selling property, borrowing money, using savings, declaring bankruptcy, changing economic lifestyle, missing bill payments, increase in insurance premiums, or reaching coverage limits): 2%–46%, most common change in income • Increased OOP expenses
Shankaran et al, 2012 ⁶⁰	CS	CSS subset of SEER, January 2008–March 2010 (n=284)	<ul style="list-style-type: none"> • Age ≥18 y • Stage III colon cancer 	<ul style="list-style-type: none"> • Mean age, 60 y (range, 22–86 y) • 49% female • 84% white 	<ul style="list-style-type: none"> • Perceived financial burden • Associated with employment, income, insurance, copayments, deductibles, monthly expenses. Self-report history of selling, refinancing, second mortgage, debt, borrowing, or ≥20% income decline. • 38% reported ≥1 treatment-related financial burden; 23% reported debt
Bernard et al, 2011 ⁶¹	CS	MEPS—household component data, 2001–2008 (n=4,110)	<ul style="list-style-type: none"> • Diagnosis of cancer • Receiving treatment • All sites 	<ul style="list-style-type: none"> • 49% aged 55–64 y • 62% female 	<ul style="list-style-type: none"> • OOP healthcare expenditure burden • High burden = health-related spending ≥20% of income (13%)
Streeter et al, 2011 ⁶²	CS	Wolters Kluwer Dynamic Lifecycle Database, 2007–2009 (n=10,508)	<ul style="list-style-type: none"> • Initiated with oral oncolytic therapy • All sites 	<ul style="list-style-type: none"> • 49% aged 41–65 y, 37% aged 65–80 y, 12% aged >81 y • 53% female • 84% commercial insurance, 17% Medicare 	<ul style="list-style-type: none"> • Cost burden • Associated with insurance plan, cost-sharing requirements, income, concurrent prescription activity • 73% had cost sharing \$0–\$100; 16% had cost sharing >\$500
Neugut et al, 2011 ⁶³	Cohort-R	Claims data from Medco Health Solutions, January 2007–December 2008 (n=22,160)	<ul style="list-style-type: none"> • Age ≥50 y • EBC • Lumpectomy or mastectomy within 12 mo of AI initiation 	<ul style="list-style-type: none"> • 24% aged 50–59 y, 35% aged 60–69 y, 28% aged 70–79 y, 13% aged ≥80 y • 90% white/other, 5% black, 3% Hispanic 	<ul style="list-style-type: none"> • Financial factors • Copayments (the amount paid by a subscriber for a 90-d mail-order prescription) ≥\$90 (30%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCGWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
de Moor et al, 2010 ⁶⁴	CS	Dana-Farber Cancer Institute/Harvard Cancer Center Rapid Case Registry (n=487)	<ul style="list-style-type: none"> Newly diagnosed DCIS ≤3 mo from surgery Fluent in English/Spanish 	<ul style="list-style-type: none"> Mean age, 54.4 ± 11.02 y 94% white 30% mastectomy, 50% radiation 	<ul style="list-style-type: none"> Financial status “How would you describe your household's financial situation right now?” measured in level of difficulty experienced with paying bills 70% high financial status (have money for special things), 18% medium status (little spare money to buy extra or special things), 12% low status (able to pay bills but need to cut back on other things or difficulty paying bills no matter what)
Khara et al, 2011 ⁶⁵	CS	Fred Hutchinson Cancer Research Center (n=1,549)	<ul style="list-style-type: none"> Age > 18 y Survived ≥2 y after HCT All sites 	<ul style="list-style-type: none"> Median age, 54.9 y (range, 18.2–81.8 y) 49% female 95% white, 2% Hispanic/Latino 28% chronic leukemia, 23% acute leukemia, 18% lymphoma 	<ul style="list-style-type: none"> Concerns about medical costs Coverage, lifetime cap, bankruptcy 25% worried that expenses will reach the limit and insurance will not pay, 1% reached the limit, 3% filed for bankruptcy
Markman & Luce, 2010 ⁶⁶	CS	NexCura cancer decision-support program by American Cancer Society (n=1,767)	<ul style="list-style-type: none"> Diagnosis of cancer of breast, colon, lung, or prostate 	<ul style="list-style-type: none"> 2% aged >60 y 58% female 22% Medicare, 25% HMO insurance, 4% Medicaid 	<ul style="list-style-type: none"> Financial impact of cancer treatment Self-reported financial (income, OOP) and subjective (distress-associated) impact 64% of all patients had “any distress”; 19% of all patients and 39% of those with <\$40,000/y income reported “large amount of distress” with financial costs (none, small amount, large amount)
Weaver et al, 2010 ⁶⁷	CS	NHIS, 2003–2006 (n=6,602)	<ul style="list-style-type: none"> Age ≥ 18 y History of cancer All sites 	<ul style="list-style-type: none"> 48% aged 18–64 y, 52% aged ≥65 y 64% female 82% white, 10% black, 8% Hispanic 63% private insurance, 30% public insurance, 7% uninsured 	<ul style="list-style-type: none"> Cost-driven healthcare access Ever had to forgo, delay, or not adhere to recommended treatment due to cost
Pezzin et al, 2009 ⁶⁸	CS	Population-based cohort of elderly patients with breast cancer (n=1,890)	<ul style="list-style-type: none"> Age ≥ 65 y Received adjuvant hormone therapy in first 12 mo post-breast surgery in 2003 	<ul style="list-style-type: none"> 34% aged 65–69 y, 33% aged 70–74 y, 23% aged 75–79 y, 11% aged ≥80 y 90% white, 3% black, 4% Hispanic 45% had full prescription drug coverage 	<ul style="list-style-type: none"> Health-related financial problems Self-reported difficulty in paying medical bills, including breast cancer-specific prescriptions in past 12 mo (“somewhat” and “very difficult” = financial problems) Any financial difficulty (31%–37%)

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCC, basaloid carcinoma; BCCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCB, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Ell et al, 2008 ⁶⁹	Cohort-L	Outpatient oncology clinic associated with University of Southern California (n=487)	<ul style="list-style-type: none"> Breast or gynecologic cancer Oversampled Hispanic women with low income Participants in IMPAACT trial In active treatment/follow-up 	<ul style="list-style-type: none"> 53% aged >50 y 76% Latina, 12% Asian, 7% non-Hispanic white, 6% black 40% Medi-Cal/Medicare, 32% government care program, 24% uninsured 49% breast cancer, 51% gynecologic cancer, 30% stage III/IV/recurrent 	<ul style="list-style-type: none"> Economic stress: at baseline, 6 mo, and 12 mo Unemployment (baseline 70%); cost concern about hospitalization, chemotherapy, radiation, medicines, or inability to pay for prescriptions (68%); wages lost for sick time or attending a medical appointment (47%) Financial stress: at 12 mo (overall 49%) Yes if economic situation worsened over past year, not satisfied with current economic situation, or not hopeful for better economic situation in future
Gupta et al, 2007 ⁷⁰	CS	Cancer Treatment Centers of America patient case series, April 2001–November 2004 (n=954)	<ul style="list-style-type: none"> Diagnosis of cancer All sites 	<ul style="list-style-type: none"> Median age, 56 y (range, 20–90 y) 61% female 26% breast cancer, 19% CRC, 16% lung cancer, 6% pancreatic cancer, 5% prostate cancer, 29% other cancer 9% stage I, 19% stage II, 20% stage III, 31% stage IV, 21% stage unknown 	<ul style="list-style-type: none"> Perceived financial difficulty OLO-C30 tool: “Has your physical condition or medical treatment caused you financial difficulties?” 31% “not at all”; 38% “a little”; 16% “quite a bit”; 15% “very much”
Francoeur, 2005 ⁷¹	CS	Northeastern United States, 1992–1993 (n=268)	<ul style="list-style-type: none"> Age ≥ 30 y Diagnosis of recurrent cancer Not receiving treatment Not terminally ill, expected to live ≥ 1 y All sites 	<ul style="list-style-type: none"> Mean age, 62 y (range, 30–90 y) 50% female 88% Caucasian 22% breast cancer, 20% lung cancer, 14% head and neck cancer, 10% gynecologic cancer, 9% prostate cancer 61% surgery, 31% radiation, 8% surgery + radiation, 34% chemotherapy 	<ul style="list-style-type: none"> Objective Family Financial Stress Index OOP expenses adjusted for family size and income; objective financial stress due to medical bills; extent of wages lost; diversity of resources tapped due to illness; change in employment; overall financial trend Subjective patient perceptions of financial strain Difficulty paying bills, inadequacy of insurance, worry about expenses
Langa et al, 2004 ⁷²	CS	National Health and Retirement study, 1995 (n=988)	<ul style="list-style-type: none"> Patient with active or prior cancer compared with adults without cancer All sites 	<ul style="list-style-type: none"> Mean age, 80 y 43%–56% female 94% white, 5% black 99% Medicare 	<ul style="list-style-type: none"> OOP expenditures Hospital and nursing home stay; outpatient services such as doctor visits, dental visits, or surgery; home care, community special services; prescriptions Adjusted mean total OOP expenditures for: cancer, no active treatment: \$1,560; and cancer, active treatment: \$2,100

(continued on next page)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCOWFU, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasm; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

Reference	Study Design	Setting/Data Source (Subjects, n)	Study Inclusion	Participant Demographic	Financial Toxicity Burden: Definition, Tool/Measure, Prevalence
Smith et al, 2003 ⁷³	Cohort-R	Academic tertiary referral center (n=101)	<ul style="list-style-type: none"> • Diagnosis of glottis cancer • Lesion ranging from Tis to invasive T1 	<ul style="list-style-type: none"> • 73% surgery, 27% radiation • 22% Tis, 78% T1 lesion 	<ul style="list-style-type: none"> • Nontangible costs of treatment • Hidden-cost questionnaire • Travel time, travel distance, treatment time, amount of work missed by patient and friends of family members
Houts et al, 1985 ⁷⁴	CS	9 medical practices affiliated with Central Pennsylvania Oncology Group (n=185)	<ul style="list-style-type: none"> • Diagnosis of cancer • >6 mo from diagnosis • All sites 	<ul style="list-style-type: none"> • Median age, 57 y (range, 16–76 y) • 70% female • 40% breast cancer, 37% other cancer • 36% curative treatment, 64% palliation 	<ul style="list-style-type: none"> • Financial burden • medical costs; OOP expenses (transportation, lodging, food, nonmedical healing, nonprescription drugs, clothing, telephone calls, miscellaneous); wages lost; total costs; % income lost • Of total financial burden incurred, 50% from wages lost, 25% OOP expenses, and 25% included medical expenses from being uninsured • Mean % of family income lost: 16%; median, 7% (range, 0%–134%)

(*references on next page*)

Abbreviations: AI, aromatase inhibitor; ALL, acute lymphoblastic leukemia; AML, acute myeloid leukemia; BCCTP, Breast and Cervical Cancer Treatment Program; BCEI, Breast Cancer Education Intervention; BCS, breast-conserving surgery; CanCORS, (NCI) Cancer Care Outcomes Research and Surveillance; CCC, comprehensive cancer center; CCCWFO, Comprehensive Cancer Care—Wake Forest University; CLL, chronic lymphocytic leukemia; CML, chronic myeloid leukemia; Cohort-L, cohort-longitudinal; Cohort-P, cohort-prospective; Cohort-R, cohort-retrospective; COIN, Collection of Indirect and Nonmedical Direct Costs, modified for monthly OOP costs; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related medication nonadherence; CS, cross-sectional; CSS, Cancer Surveillance System; DCIS, ductal carcinoma in situ; EBC, early-stage breast cancer; ECSS, Experiences With Cancer Survivorship Supplement; GPH, general public hospital; HCT, hematopoietic cell transplantation; HMO, health maintenance organization; MBC, metastatic breast cancer; MCBBS, Medicare Current Beneficiary Survey; MDS/MPN, myelodysplastic syndromes/myeloproliferative neoplasms; MEPS, Medical Expenditure Panel Survey; NHIS, National Health Interview Survey; NHL, non-Hodgkin's lymphoma; OOP, out-of-pocket; POS, point of service (insurance); PPO, preferred provider organization; VA, Veterans Administration.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

References

- Shankaran V, Leahy T, Steelquist J, et al. Pilot feasibility study of an oncology financial navigation program. *J Oncol Pract* 2018;14:e122–129.
- Wheeler SB, Spencer JC, Pinheiro LC, et al. Financial impact of breast cancer in black versus white women. *J Clin Oncol* 2018;36:1695–1701.
- de Souza JA, Kung S, O'Connor J, et al. Determinants of patient-centered financial stress in patients with locally advanced head and neck cancer. *J Oncol Pract* 2017;13:e310–318.
- de Souza JA, Yap BJ, Wroblewski K, et al. Measuring financial toxicity as a clinically relevant patient-reported outcome: the validation of the Comprehensive Score for financial Toxicity (COST). *Cancer* 2017;123:476–484.
- Farias AJ, Du XL. Association between out-of-pocket costs, race/ethnicity, and adjuvant endocrine therapy adherence among Medicare patients with breast cancer. *J Clin Oncol* 2017;35:86–95.
- Meeker CR, Wong YN, Egleston BL, et al. Distress and financial distress in adults with cancer: an age-based analysis. *J Natl Compr Canc Netw* 2017;15:1224–1233.
- Pisu M, Azuero A, Benz R, et al. Out-of-pocket costs and burden among rural breast cancer survivors. *Cancer Med* 2017;6:572–581.
- Abel GA, Albelda R, Khera N, et al. Financial hardship and patient-reported outcomes after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016;22:1504–1510.
- Banegas MP, Guy GP Jr, de Moor JS, et al. For working-age cancer survivors, medical debt and bankruptcy create financial hardships. *Health Aff (Millwood)* 2016;35:54–61.
- Chongpison Y, Hornbrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
- Delgado-Guay MO, Chisholm G, Williams J, et al. Frequency, intensity, and correlates of spiritual pain in advanced cancer patients assessed in a supportive/palliative care clinic. *Palliat Support Care* 2016;14:341–348.
- DiMartino LD, Birken SA, Mayer DK. The relationship between cancer survivors' socioeconomic status and reports of follow-up care discussions with providers. *J Cancer Educ* 2017;32:749–755.
- Farias AJ, Hansen RN, Zeliadt SB, et al. The association between out-of-pocket costs and adherence to adjuvant endocrine therapy among newly diagnosed breast cancer patients. *Am J Clin Oncol* 2018;41:708–715.
- Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
- Keim-Malpass J, Levine B, Danhauer SC, et al. Work-related perceptions and quality of life among breast cancer survivors. *Psychooncology* 2016;25:873–876.
- Lathan CS, Cronin A, Tucker-Seeley R, et al. Association of financial strain with symptom burden and quality of life for patients with lung or colorectal cancer. *J Clin Oncol* 2016;34:1732–1740.
- Lee M, Khan MM. Gender differences in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:384–393.
- Lee M, Salloom RG. Racial and ethnic disparities in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:534–544.
- Meeker CR, Geynisman DM, Egleston BL, et al. Relationships among financial distress, emotional distress, and overall distress in insured patients with cancer. *J Oncol Pract* 2016;12:e755–764.
- Narang AK, Nicholas LH. Out-of-pocket spending and financial burden among Medicare beneficiaries with cancer. *JAMA Oncol* 2017;3:757–765.
- Nipp RD, Zullig LL, Samsa G, et al. Identifying cancer patients who alter care or lifestyle due to treatment-related financial distress. *Psychooncology* 2016;25:719–725.
- Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
- Whitney RL, Bell JF, Reed SC, et al. Predictors of financial difficulties and work modifications among cancer survivors in the United States. *J Cancer Surviv* 2016;10:241–250.
- Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
- Yabroff KR, Dowling EC, Guy GP Jr, et al. Financial hardship associated with cancer in the United States: findings from a population-based sample of adult cancer survivors. *J Clin Oncol* 2016;34:259–267.
- Zheng Z, Yabroff KR, Guy GP Jr, et al. Annual medical expenditure and productivity loss among colorectal, female breast, and prostate cancer survivors in the United States. *J Natl Cancer Inst* 2015;108:djv382.
- Callahan C, Brintzenhofesoc K. Financial quality of life for patients with cancer: an exploratory study. *J Psychosoc Oncol* 2015;33:377–394.
- Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
- Guy GP Jr, Yabroff KR, Ekwueme DU, et al. Healthcare expenditure burden among non-elderly cancer survivors, 2008–2012. *Am J Prev Med* 2015;49(Suppl 5):S489–497.
- Huntington SF, Weiss BM, Vogt DT, et al. Financial toxicity in insured patients with multiple myeloma: a cross-sectional pilot study. *Lancet Haematol* 2015;2:e408–416.
- Jones WC, Parry C, Devine S, et al. Prevalence and predictors of distress in posttreatment adult leukemia and lymphoma survivors. *J Psychosoc Oncol* 2015;33:124–141.
- Palmer NR, Weaver KE, Hauser SP, et al. Disparities in barriers to follow-up care between African American and White breast cancer survivors. *Support Care Cancer* 2015;23:3201–3209.
- Pisu M, Kenzik KM, Oster RA, et al. Economic hardship of minority and non-minority cancer survivors 1 year after diagnosis: another long-term effect of cancer? *Cancer* 2015;121:1257–1264.
- Tucker-Seeley RD, Abel GA, Uno H, et al. Financial hardship and the intensity of medical care received near death. *Psychooncology* 2015;24:572–578.
- Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
- Bender CM, Gentry AL, Brufsky AM, et al. Influence of patient and treatment factors on adherence to adjuvant endocrine therapy in breast cancer. *Oncol Nurs Forum* 2014;41:274–285.
- Bestvina CM, Zullig LL, Rushing C, et al. Patient-oncologist cost communication, financial distress, and medication adherence. *J Oncol Pract* 2014;10:162–167.
- de Souza JA, Yap BJ, Hlubocky FJ, et al. The development of a financial toxicity patient-reported outcome in cancer: the COST measure. *Cancer* 2014;120:3245–3253.
- Dusetzina SB, Winn AN, Abel GA, et al. Cost sharing and adherence to tyrosine kinase inhibitors for patients with chronic myeloid leukemia. *J Clin Oncol* 2014;32:306–311.
- Ekwueme DU, Yabroff KR, Guy GP Jr, et al. Medical costs and productivity losses of cancer survivors—United States, 2008–2011. *MMWR Morb Mortal Wkly Rep* 2014;63:505–510.
- Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
- Jagsi R, Hawley ST, Abrahamse P, et al. Impact of adjuvant chemotherapy on long-term employment of survivors of early-stage breast cancer. *Cancer* 2014;120:1854–1862.
- Jagsi R, Pottow JAE, Griffith KA, et al. Long-term financial burden of breast cancer: experiences of a diverse cohort of survivors identified through population-based registries. *J Clin Oncol* 2014;32:1269–1276.
- Khera N, Chang YH, Hashmi S, et al. Financial burden in recipients of allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2014;20:1375–1381.

eTable 2. Study and Patient Characteristics and Measures of Financial Toxicity Burdens (cont.)

45. Kircher SM, Johansen ME, Nimeiri HS, et al. Impact of Medicare Part D on out-of-pocket drug costs and medical use for patients with cancer. *Cancer* 2014;120:3378–3384.
46. Martin MY, Fouad MN, Oster RA, et al. What do cancer patients worry about when making decisions about treatment? Variation across racial/ethnic groups. *Support Care Cancer* 2014;22:233–244.
47. Regenberg SE, Veenstra CM, Hawley ST, et al. The personal financial burden of complications after colorectal cancer surgery. *Cancer* 2014;120:3074–3081.
48. Veenstra CM, Regenberg SE, Hawley ST, et al. A composite measure of personal financial burden among patients with stage III colorectal cancer. *Med Care* 2014;52:957–962.
49. Blinder V, Patil S, Eberle C, et al. Early predictors of not returning to work in low-income breast cancer survivors: a 5-year longitudinal study. *Breast Cancer Res Treat* 2013;140:407–416.
50. Davidoff AJ, Erten M, Shaffer T, et al. Out-of-pocket health care expenditure burden for Medicare beneficiaries with cancer. *Cancer* 2013;119:1257–1265.
51. Dowling EC, Chawla N, Forsythe LP, et al. Lost productivity and burden of illness in cancer survivors with and without other chronic conditions. *Cancer* 2013;119:3393–3401.
52. Goodwin JA, Coleman EA, Sullivan E, et al. Personal financial effects of multiple myeloma and its treatment. *Cancer Nurs* 2013;36:301–308.
53. Hamilton JG, Wu LM, Austin JE, et al. Economic survivorship stress is associated with poor health-related quality of life among distressed survivors of hematopoietic stem cell transplantation. *Psychooncology* 2013;22:911–921.
54. Kent EE, Forsythe LP, Yabroff KR, et al. Are survivors who report cancer-related financial problems more likely to forgo or delay medical care? *Cancer* 2013;119:3710–3717.
55. Ramsey S, Blough D, Kirchhoff A, et al. Washington State cancer patients found to be at greater risk for bankruptcy than people without a cancer diagnosis. *Health Aff (Millwood)* 2013;32:1143–1152.
56. Wan Y, Gao X, Mehta S, et al. Indirect costs associated with metastatic breast cancer. *J Med Econ* 2013;16:1169–1178.
57. Zafar SY, Peppercorn JM, Schrag D, et al. The financial toxicity of cancer treatment: a pilot study assessing out-of-pocket expenses and the insured cancer patient's experience. *Oncologist* 2013;18:381–390.
58. Zullig LL, Peppercorn JM, Schrag D, et al. Financial distress, use of cost-coping strategies, and adherence to prescription medication among patients with cancer. *J Oncol Pract* 2013;9(Suppl 6):60s–63s.
59. Meneses K, Azuero A, Hassey L, et al. Does economic burden influence quality of life in breast cancer survivors? *Gynecol Oncol* 2012;124:437–443.
60. Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
61. Bernard DSM, Farr SL, Fang Z. National estimates of out-of-pocket health care expenditure burdens among nonelderly adults with cancer: 2001 to 2008. *J Clin Oncol* 2011;29:2821–2826.
62. Streeter SB, Schwartzberg L, Husain N, et al. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *Am J Manag Care* 2011;17(Suppl 5):SP38–44.
63. Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
64. de Moor JS, Partridge AH, Winer EP, et al. The role of socioeconomic status in adjustment after ductal carcinoma in situ. *Cancer* 2010;116:1218–1225.
65. Khera N, Chow EJ, Leisenring WM, et al. Factors associated with adherence to preventive care practices among hematopoietic cell transplantation survivors. *Biol Blood Marrow Transplant* 2011;17:995–1003.
66. Markman M, Luce R. Impact of the cost of cancer treatment: an internet-based survey. *J Oncol Pract* 2010;6:69–73.
67. Weaver KE, Rowland JH, Bellizzi KM, et al. Forgoing medical care because of cost: assessing disparities in healthcare access among cancer survivors living in the United States. *Cancer* 2010;116:3493–3504.
68. Pezzin LE, O'Neil MB, Nattinger AB. The economic consequences of breast cancer adjuvant hormonal treatments. *J Gen Intern Med* 2009;24(Suppl 2):S446–450.
69. Ell K, Xie B, Wells A, et al. Economic stress among low-income women with cancer: effects on quality of life. *Cancer* 2008;112:616–625.
70. Gupta D, Lis CG, Grutsch JF. Perceived cancer-related financial difficulty: implications for patient satisfaction with quality of life in advanced cancer. *Support Care Cancer* 2007;15:1051–1056.
71. Francoeur RB. Cumulative financial stress and strain in palliative radiation outpatients: the role of age and disability. *Acta Oncol* 2005;44:369–381.
72. Langa KM, Fendrick AM, Chermew ME, et al. Out-of-pocket health-care expenditures among older Americans with cancer. *Value Health* 2004;7:186–194.
73. Smith JC, Johnson JT, Cognetti DM, et al. Quality of life, functional outcome, and costs of early glottic cancer. *Laryngoscope* 2003;113:68–76.
74. Houts PS, Harvey HA, Simmonds MA, et al. Characteristics of patients at risk for financial burden because of cancer and its treatment. *J Psychosoc Oncol* 1985;3:15–22.

eTable 3. Strength of Evidence Assessment Using GRADE Criteria

Analysis	Assessment	Other Details
Sample-weighted prevalence of financial burden		
Limitations in study design	Downgraded	8 studies judged “fair” in risk of bias
Inconsistency of result	Downgraded	Because of substantial heterogeneity
Indirectness of evidence	Downgraded	Because of substantial heterogeneity
Imprecision	Not downgraded	
Publications of bias	Not downgraded	Not present
General assessment	Very low	
Association between insurance status and financial burdens		
Limitations in study design	Not downgraded	
Inconsistency of result	Downgraded	Because of substantial heterogeneity
Indirectness of evidence	Downgraded	Because of substantial heterogeneity
Imprecision	Not downgraded	
Publications of bias	Not downgraded	Cannot be assessed
General assessment	Low	
Association between financial burdens and cancer medication adherence		
Limitations in study design	Not downgraded	
Inconsistency of result	Downgraded	Because of substantial heterogeneity
Indirectness of evidence	Downgraded	Differences in populations and outcomes
Imprecision	Downgraded	Large confidence intervals
Publications of bias	Not downgraded	Cannot be assessed
General assessment	Very low	

Abbreviation: GRADE, Grading of Recommendations Assessment, Development, and Evaluation criteria.

eTable 4. Risk Factors of Financial Toxicity Burdens

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Wheeler et al, 2018 ¹ • P, I	Adjusted logistic regression • Financial impact score	<ul style="list-style-type: none"> Income: adjusted risk differences for adverse financial impact; compared with income >\$50,000/y, patients with income \$15,000–\$29,999/y had ↑ financial impact score (worse impact) (ASD, 13.93 [SE, 3.33]) and those with income \$30,000–\$49,999 had ↑ financial impact score (ASD, 9.279 [SE, 2.86]; $P < .05$) Insurance: compared with privately insured patients, those with no insurance and Medicaid had ↑ financial impact score (worse impact) (ASD, 28.45 [SE, 4.74] and 21.11 [SE, 3.75]; $P < .001$) 	<ul style="list-style-type: none"> Age: older age had ↓ financial impact score (less impact) (ASD, –0.70 [SE, 0.10] per y); $P < .001$ Race/Ethnicity: black patients had ↑ financial impact score (worse impact) (ASD, 5.42 [SE, 2.16]) compared with white patients; $P < .05$ Marital status: separated/divorced patients had ↑ financial impact score (worse impact) (ASD, 6.35 [SE, 2.67]) compared with married patients; $P < .05$ 	<ul style="list-style-type: none"> Chemotherapy: ↑ financial impact score (worse impact) (ASD, 9.59 [SE, 2.53]) compared with no chemotherapy; $P < .05$ Cancer stage: stage IV had ↑ financial impact score (worse impact) (ASD, 14.09 [SE, 6.22]) compared with stage I; $P < .05$
de Souza et al, 2017 ² • P	Adjusted logistic regression • COST score (↑ COST score = lower financial toxicity)	<ul style="list-style-type: none"> Household income: high-income patients had ↑ COST score; patients with >800% of FPL had 12-point ↑ mean COST score vs those with ≤200% of FPL ($P = .003$) Employment: unemployed patients had ↓ COST score ($\beta = 11.68$; 95% CI, 4.43–18.44; $P < .001$) 	<ul style="list-style-type: none"> Race/Ethnicity: white patients had ↑ COST scores vs Hispanic ($\beta = -2.41$; 95% CI, –8.66–3.83), black ($\beta = -5.41$; 95% CI, –9.60 to –0.67), and Asian and Native American patients ($\beta = -9.85$; 95% CI, –19.82 to 0.12; $P = .04$) 	<ul style="list-style-type: none"> Number of in-patient admissions: patients with ≥ 3 vs ≤ 2 hospital admissions had ↓ COST scores ($\beta = -5.52$; 95% CI, –9.87 to –1.16; $P = .01$) Psychological distress: Patients with high distress had ↓ COST scores ($\beta = -0.34$; 95% CI, –0.56 to –0.12; $P = .003$)
Meeker et al, 2017 ³ • P	Adjusted linear regression • Financial distress score	<ul style="list-style-type: none"> Income: income $\geq \\$75,000/y$ had ↑ financial score (less distress) vs income <\$25,000/y ($\beta = 1.88$; 95% CI, 0.27–3.49; $P = .02$) 	<ul style="list-style-type: none"> Age: increasing age ≥ 65 y had ↑ financial score (less distress) compared with age <50 y ($\beta = 1.52$; 95% CI, 0.03–3.01; $P = .05$) 	
Pisu et al, 2017 ⁴ • D	Repeated-measures model with generalized linear mixed method • ↑ OOP cost	<ul style="list-style-type: none"> Employment: part-time, unemployed, retired, or on disability had ↑ OOP burden vs full-time employed, $P = .036$ Income decrease since end of treatment: patients with income decrease since treatment had ↑ OOP costs (mean ratio, 1.42; 95% CI, 1.11–1.81; $P = .005$) and ↑ OOP cost burden (mean ratio, 1.66; 95% CI, 1.21–2.29; $P = .002$) 	<ul style="list-style-type: none"> Age: age ≥ 65 y had ↓ OOP cost burden (mean ratio, 0.68; 95% CI, 0.50–0.92; $P = .01$) 	<ul style="list-style-type: none"> Time since treatment: patients within 12 mo of treatment (mean ratio, 1.55; 95% CI, 1.10–2.19; $P = .04$) had ↑ OOP cost compared with those within ≥ 24 mo of treatment Use of supportive services: patients using support services had ↑ OOP costs (mean ratio, 1.35; 95% CI, 1.03–1.76; $P = .03$) and ↑ OOP burden (mean ratio, 1.46; 95% CI, 1.05–2.03; $P = .03$)
Abel et al, 2016 ⁵ • P, I	Adjusted logistic regression • Financial hardship	<ul style="list-style-type: none"> Monthly income: low-income (OR, 10.43; 95% CI, 4.28–25.4; $P < .001$) and middle-income patients (OR, 2.92; 95% CI, 1.47–5.78; $P = .002$) experienced any of the 3 hardships; low-income patients (OR, 5.45; 95% CI, 1.43–20.7; $P = .01$) more likely to experience all 3 hardships than middle- or high-income patients. 	<ul style="list-style-type: none"> Marital status: married patients less likely to experience all hardships (OR, 0.39; 95% CI, 0.18–0.88; $P = .02$) 	<ul style="list-style-type: none"> Treatment type: patients with transplant-related costs more likely to experience any hardship (OR, 6.86; 95% CI, 3.82–12.3; $P < .001$) and all 3 hardships (OR, 3.84; 95% CI, 1.57–9.36; $P = .003$)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

(continued on next page)

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Banegas et al, 2016 ⁶ • I	Adjusted logistic regression • Borrow, go into debt, bankruptcy	<ul style="list-style-type: none"> Income: lower-income patients more likely to borrow/go into debt (\$40,000–\$80,000/y: OR, 2.46; $P<.001$; <\$40,000/y: OR, 3.52; $P<.001$) and file for bankruptcy (\$40,000–\$80,000/y: OR, 2.80; $P<.001$; <\$40,000/y: OR, 4.50; $P<.001$) compared with >\$80,000/y Employment status: unemployed (OR, 1.28; $P<.05$) more likely and retired (OR, 0.51; $P<.001$) less likely to borrow/go into debt; unemployed patients (OR, 1.74; $P<.05$) more likely to file for bankruptcy Insurance: publicly insured (OR, 1.95; $P<.001$) and uninsured patients (OR, 2.46; $P<.001$) more likely to borrow/go into debt; publicly insured patients (OR, 1.82; $P<.05$) more likely to file for bankruptcy 	<ul style="list-style-type: none"> Age: younger patients more likely to borrow money/go into debt (aged 45–54 y: OR, 1.66; aged 18–44 y: OR, 2.07; $P<.001$) and file for bankruptcy (aged 45–54 y: OR, 1.86; aged 18–44 y: OR, 1.81; $P<.05$) compared with those aged 55–64 y Sex and marital status: unmarried males (OR, 1.37; $P<.05$) and unmarried females (OR, 1.56; $P<.001$) more likely to borrow money/go into debt Race/Ethnicity: Hispanic patients (OR, 1.41; $P<.05$) more likely to borrow/go into debt than non-Hispanic patients 	<ul style="list-style-type: none"> Number of cancer diagnoses: patients with ≥ 2 more likely to borrow/go into debt (OR, 1.49; $P<.001$) and file for bankruptcy (OR, 2.00; $P<.001$) Time since last treatment: ≥ 6 y (OR, 0.82; $P<.05$) and 3–5 y (OR, 1.71; $P<.05$) less likely to borrow/go into debt compared with <1 y; ≥ 6 y (OR, 1.65; $P<.05$) more likely to file for bankruptcy
Chongpison et al, 2016 ⁷ • P	Mean and SD • Financial burden	<ul style="list-style-type: none"> Education: patients with lower education (high school or less) (mean, 2.41 ± 3.06) had $\uparrow$ financial burden score Household income: patients with lower income ($\leq \\$30,000/y$) (mean, 3.11 ± 3.34) had $\uparrow$ burden Employment: full-time or part-time (mean, 2.09 ± 2.83), unemployed (mean, 5.27 ± 3.77), and patients on disability (mean, 5.33 ± 3.06) had $\uparrow$ burden compared with retired and nonworking patients (mean, 1.69 ± 2.55) 		<ul style="list-style-type: none"> Type of surgery: patients with permanent (mean, 2.56 ± 3.01) and temporary ostomy (mean, 2.28 ± 2.74) had $\uparrow$ burden score than those with anastomosis (mean, 1.63 ± 2.62) Recall of depression after surgery: patients with depression (mean, 3.22 ± 3.34) had $\uparrow$ burden than those with no depression (mean, 1.35 ± 2.18)
Kale & Carroll, 2016 ⁸ • P, I	Adjusted logistic regression • Financial burden	<ul style="list-style-type: none"> Insurance: patients with no insurance during treatment (OR, 1.99; CI, 1.33–2.98) had $\uparrow$ burden compared with insured patients 	<ul style="list-style-type: none"> Age: younger patients (OR, 1.87; 95% CI, 1.23–2.83) had $\uparrow$ burden Sex: female patients (OR, 1.43; 95% CI, 1.00–2.04) had $\uparrow$ burden Race/Ethnicity: Hispanic (OR, 1.68; 95% CI, 1.02–2.76) and black patients (OR, 1.09; 95% CI, 0.51–2.18) had $\uparrow$ burden than non-Hispanic white patients 	<ul style="list-style-type: none"> Cancer type: patients with short-survival type cancer (OR, 3.28; 95% CI, 1.43–7.51) had $\uparrow$ burden than patients with breast cancer Time since last treatment: > 1 y (OR, 2.55; 95% CI, 1.68–3.86) had $\uparrow$ burden than ≥ 5 y Time since diagnosis: 1–<3 previous y (OR, 1.69; 95% CI, 1.14–2.50) had $\uparrow$ burden than ≥ 5 y

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Lathan et al, 2016 ⁹ • I	Unadjusted • Financial reserves	<ul style="list-style-type: none"> Insurance: patients reporting <2 mo financial reserves more likely to be insured by Medicaid or have no insurance than patients reporting >2 mo financial reserves Supplemental insurance: patients without had ↑ mean OOP expenses and financial burden, with a spending mean of 23.7% of household income on OOP expenses 		
Narang & Nicholas, 2017 ¹⁰ • D	Adjusted logistic regression • OOP expenses			<ul style="list-style-type: none"> Age: younger patients (mean, aged 52.6 vs 63.9 y) had ↑ odds of bankruptcy vs no bankruptcy Sex: females (55%) had ↑ odds bankruptcy Race: nonwhite patients (14% vs 11%) had ↑ odds bankruptcy
Ramsey et al, 2016 ¹¹ • I	Adjusted logistic regression • Bankruptcy in original cohort			<ul style="list-style-type: none"> Health status: patients in active treatment with good/fair/poor health status had ↑ financial difficulty (OR, 3.8; 95% CI, 1.0–14.2) and ↑ work modification (OR, 10.4; 95% CI, 2.0–54.8) compared with patients with excellent/very good health status; among patients age <65 y, those with good/fair/poor health status had ↑ financial difficulty (OR, 2.4; 95% CI, 1.4–4.1) and ↑ work modification (OR, 4.1; 95% CI, 1.6–10.2) compared with those with excellent/very good health
Whitney et al, 2016 ¹² • P, I	Adjusted logistic regression • Financial difficulty • Work modification	<ul style="list-style-type: none"> Income: patients in active treatment with income <200% of FPL (OR, 15.7; 95% CI, 2.6–95.2) or 200 to <400% of FPL (OR, 8.2; 95% CI, 1.3–52.4) had ↑ financial difficulty compared with income >400% FPL Residence: patients in nonmetropolitan area (OR, 6.4; 95% CI, 1.6–25.0) had ↑ financial difficulty compared with those in metropolitan area Education: patients aged <65 y with 12–16 y education (OR, 1.9; 95% CI, 1.1–3.6) had ↑ financial difficulty compared with ≥16 y education Insurance: patients aged <65 y and uninsured (OR, 3.5; 95% CI, 1.3–9.3) or publicly insured (OR, 9.0; 95% CI, 3.3–24.4) had ↑ work modification 	<ul style="list-style-type: none"> Race/Ethnicity: nonwhite patients (OR, 8.0; 95% CI, 2.2–28.4) had ↑ financial difficulty compared with white patients Marital status: patients aged <65 y and married (OR, 2.2; 95% CI, 1.0–4.7) had ↑ work modification compared with unmarried patients 	
Wong et al, 2016 ¹³ • P	One-way ANOVA or Kruskal-Wallis test • Cost concern	<ul style="list-style-type: none"> Education: patients with <high school (mean, 9.66±2.44; P=.01) had ↑ mean of total cost concerns Employment: unemployed (mean, 9.33±2.62; P<.001) had ↑ mean of total cost concerns 	<ul style="list-style-type: none"> Age: patients aged ≤65 y (mean, 8.86±2.67; P<.001) had ↑ mean of total cost concerns Sex: females (mean, 8.77±2.68; P=.04) had ↑ mean of total cost concerns Marital status: unmarried patients (mean, 8.97±2.62; P=.01) had ↑ mean of total cost concerns 	

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Yabroff et al, 2016 ¹⁴ • P, I	Adjusted logistic regression • 1. Material financial hardship • 2. Psychologic financial hardship	Patients aged 18–64: • Employment: employed but took extended leave or switched to part time had ↑ material hardship ($P<.001$) • Income and time since treatment: lower family income meant ↑ psychological hardship ($P=.019$) • Insurance: uninsured had ↑ psychological hardship ($P=.043$)	• Age: younger patients (aged 18–54 y vs aged 55–64 y vs aged 65–74 y vs aged ≥75 y) had ↑ material hardship ($P=.009$) • Age: in ≥65 y age group, those aged 65–74 had ↑ psychological hardship ($P=.012$) • Race/Ethnicity: nonwhite had ↑ material hardship ($P=.008$) • Sex: females had ↑ material hardship ($P=.034$)	• Time since treatment: shorter time interval since treatment had ↑ material ($P=.014$) and psychological ($P=.024$) hardship
Zheng et al, 2015 ¹⁵ • I, D	Frequencies • Employment disability and productivity loss		• Age: nonelderly CRC and breast cancer survivors had ↑ employment disability (13.6%, $P<0.001$; 4.8%, $P=.001$) and productivity loss at work (7.2 d, $P<0.001$; 3.3 d, $P=.002$) and at home (4.5 d, $P<.001$; 3.3 d, $P=.003$)	
Callahan & Brintzenhofesoc, 2015 ¹⁶ • P	Adjusted regression • Financial QoL	• Health insurance adequacy: patients with health insurance adequacy had ↑ financial QoL ($\beta=.259$; $P=.002$) • Financial stress: patients with lower financial stress had ↑ financial QoL ($\beta=.370$; $P<.001$) • Barriers to care: patients with lower barriers to care had ↑ financial QoL ($\beta=-.329$; $P=.003$)		
Delgado-Guay et al, 2015 ¹⁷ • P, I	Frequencies • Financial distress			• Cancer care facility: financial distress ↑ in GPH vs CCC • Financial distress present in 86% (95% CI, 76%–93%) served at CCC vs 90% (95% CI, 81%–96%) served at GPH ($P=.45$) • Median intensity score of financial distress, CCC = 4 vs GPH = 8, $P=.0003$.
Guy et al, 2015 ¹⁸ • D	Adjusted logistic regression • OOP burden	• Income: patients who were ear-poor/low-income (100%–200% FPL) had ↑ OOP burden compared with noncancer individuals; $P=.002$ • Employment: full-time employed had ↑ OOP burden compared with noncancer individuals; $P<.05$	• Age: patients aged 50–64 y had ↑ OOP burden compared with noncancer individuals; $P=.02$	
Huntington et al, 2015 ¹⁹ • P	Adjusted regression analyses • Financial burden (COST score)	• Income: patients with lower income, \$40,000–\$79,999/y ($\beta=7.8$; 95% CI, 2.7–12.9; $P=.0031$) vs ≥\$80,000 ($\beta=11.8$; 95% CI, 7.1–16.4; $P<.0001$) had ↑ financial burden	• Marital status: nonmarried patients ($\beta=5.6$; 95% CI, 1.5–9.6; $P=.007$) had ↑ financial burden • Age: younger patients ($\beta=0.36$; 95% CI, 0.15–0.56; $P<.001$) had ↑ financial burden	• Time since diagnosis: patients with longer duration since diagnosis ($\beta=-4.8$; 95% CI, -9.3 to -0.2; $P=.042$) had ↑ financial burden

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

(continued on next page)

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Jones et al, 2015 ²⁰ • P	ANOVA • Perceived financial burden	<ul style="list-style-type: none"> Income: increased income had ↓ financial burden: <\$25,000/y (mean, 5.9±3.9), \$25,000–\$50,000/y (mean, 5.2±3.7), \$51,000–\$75,000/y (mean, 4.1±3.5), \$76,000–\$100,000/y (mean, 3.4±3.2), >\$100,000/y (mean, 2.8±3.0); $P<.001$ 	<ul style="list-style-type: none"> Age: increased age had ↓ financial burden, $P<.001$; age 18–39 y (mean, 6.1±3.5), age 40–64 y (mean, 4.7±3.6), age >65 y (mean, 2.5±3.1) Marital status: married patients had ↓ financial burden (mean, 3.9; SD, 3.6) vs unmarried patients (mean, 4.8±3.8); $P=.024$ Race: black patients had ↑ OOP vs white patients (52% vs 28%; $P=.01$) 	<ul style="list-style-type: none"> Time since treatment: patients with longer time since treatment had ↓ financial burden ($P=.038$); patients still receiving “maintenance” treatment (mean, 4.7±3.0) vs 3–24 mo (mean, 4.6±3.8) and 25–48 mo (mean, 3.8±3.5) from any treatment
Palmer et al, 2015 ²¹ • P, I	Chi-square test • OOP costs			
Pisu et al, 2015 ²² • P	Adjusted logistic regression • Economic hardship	<ul style="list-style-type: none"> No drug coverage: ↑ risk of economic hardship among patients with lung cancer (OR, 1.47; 95% CI, 1.01–2.14) and CRC (OR, 1.48; 95% CI, 1.02–2.13) Income: ≥\$20,000/y had ↓ risk of economic hardship 	<ul style="list-style-type: none"> Race/Ethnicity: among patients with CRC, black patients had ↑ risk of economic hardship vs white patients (OR, 1.69; 95% CI, 1.25–2.29) Number of individuals supported: patients with ≥3 dependents had ↑ risk of economic hardship among patients with lung cancer (OR, 1.80; 95% CI, 1.13–2.85) and with CRC (OR, 1.59; 95% CI, 1.11–2.26) Age: among patients with lung cancer, those aged ≥65 y had ↓ risk of economic hardship; those with CRC aged ≥55 y had ↓ risk of economic hardship 	<ul style="list-style-type: none"> Intensity of end-of-life care received: patients with ↑ financial hardship received more intensive end-of-life care (OR, 3.05; 95% CI, 1.22–7.62)
Tucker-Seeley et al, 2015 ²³ • I	Adjusted logistic regression • Financial hardship			
Zafar et al, 2015 ²⁴ • P	Adjusted path models • Financial difficulty	<ul style="list-style-type: none"> Income: patients with lower income (<\$20,000/y) (OR, 0.61; $P<.001$) had ↑ financial difficulty 	<ul style="list-style-type: none"> Age: younger (OR, 0.65 per 10 y older, $P<.001$) had ↑ financial difficulty 	
Fenn et al, 2014 ²⁵ • P	Bivariate analyses • Financial problems	<ul style="list-style-type: none"> ↓ Education: had ↑ financial problems ($P=.002$) ↓ Income: had ↑ financial problems ($P<.001$) Insurance: Medicare had ↓ financial problems ($P<.001$) 	<ul style="list-style-type: none"> Age: had ↑ financial problems ($P<.001$) Sex: females had ↑ financial problems ($P=.016$) Race: nonwhite patients had ↑ financial problems ($P=.001$) 	<ul style="list-style-type: none"> Type of cancer: patients with thyroid (30.6%), ovarian (25.3%), and lung cancer (23.7%) had ↑ financial problems ($P<.001$)
Jagsi et al, 2014 ²⁶ • I	Adjusted logistic regression • Excess unemployment	<ul style="list-style-type: none"> Employment support: patients with no support (OR, 1.33; 95% CI, 1.08–1.67; $P=.011$) had excess unemployment 	<ul style="list-style-type: none"> Age: older patients (>56 vs <46 y; OR, 1.42; 95% CI, 1.03–1.95; $P=.031$) had excess unemployment 	<ul style="list-style-type: none"> Comorbidity: patients with ≥2 comorbidities (OR, 2.16; 95% CI, 1.59–2.94; $P<.001$) had excess unemployment Chemotherapy: patients who received chemotherapy (OR, 1.42; 95% CI, 1.03–1.98; $P=.038$) had excess unemployment

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Jagsi et al, 2014 ²⁷ • P, I, D	Adjusted logistic regression • Financial decline	<ul style="list-style-type: none"> Income: patients with <\$50,000 vs >\$50,000 had ↑ financial decline (OR, 1.77; P=.034) Employment: patients working part-time had ↑ financial decline vs those not working (OR, 1.88; P=.019) Patients with reductions in work hours vs no reductions had ↑ financial decline (OR, 1.68; P=.04) Prescription insurance: partial (OR, 3.67; P=.001) and no coverage (OR, 2.70; P=.01) vs complete coverage had ↑ financial decline 	<ul style="list-style-type: none"> Age: younger patients (all categories <65 vs ≥65 y; P<.001) had ↑ financial decline Race: Spanish-speaking Latina vs white patients (OR, 2.76; P=.006) had ↑ financial decline 	<ul style="list-style-type: none"> Chemotherapy: patients who received chemotherapy (OR, 2.14; P=.002) had ↑ financial decline Recurrent breast cancer: patients with recurrent disease (OR, 2.44; P=.017) had ↑ financial decline
Khera et al, 2014 ²⁸ • P, I	Adjusted logistic regression • Objective financial burden	<ul style="list-style-type: none"> Employment: compared with retired, employed patients had ↓ (OR, 0.32; 95% CI, 0.18–0.50; P=.0004) 	<ul style="list-style-type: none"> Age (per 5-y increase): younger patients (OR, 0.80; 95% CI, 0.7–0.97; P=.0009) had ↑ financial burden 	<ul style="list-style-type: none"> Current physical and mental functioning: patients with lower score (<40) on PCS-12 (OR, 3.02; 95% CI, 1.6–5.9; P=.001) and MCS-12 (OR, 3.50; 95% CI, 1.4–8.5; P=.006) had ↑ financial burden
Martin et al, 2014 ²⁹ • P	Adjusted logistic regression • Worry about treatment costs	<ul style="list-style-type: none"> Patients with higher income (≥\$20,000/y), higher education (≥high school), insurance (treatment being covered), and savings (≥1 y) had significant ↓ worry; all P<.05 	<ul style="list-style-type: none"> Race: Asian vs white: those with ≥2 dependents vs 1 dependent to support had ↑ worry, P<.05 Age: older patients (all categories aged ≥60 y vs aged <54 y) had ↑ worry, P<.05 	
Regenbogen et al, 2014 ³⁰ • I	Poisson regression • Personal financial burden (positive EE = ↑ burden)	<ul style="list-style-type: none"> Income: patients with lower income had ↑ financial burden. <\$20,000/y (EE, 0.41; 95% CI, 0.19–0.63; P<.001); \$20,000–\$49,000/y (EE, 0.65; 95% CI, 0.46–0.83; P<.001); \$50,000–\$89,000/y (EE, 0.32; 95% CI, 0.15–0.50; P<.001) vs ≥\$90,000/y 	<ul style="list-style-type: none"> Age: younger patients had ↑ financial burden compared with those aged <50 y, 50–64 y (EE, –0.22; 95% CI, –0.36 to 0.10; P<.001); 65–74 y (EE, –0.68; 95% CI, –0.83 to –0.52; P<.001); ≥75 y (EE, 1.00; 95% CI, –1.19 to –0.80; P<.001) 	<ul style="list-style-type: none"> Postoperative complications: yes (EE, 0.31; 95% CI, 0.20–0.42; P<.001) had ↑ financial burden Chemotherapy: yes (EE, 0.54; 95% CI, 0.32–0.75; P<.01) had ↑ financial burden
Veenstra et al, 2014 ³¹ • P	t test or ANOVA • Personal financial burden	<ul style="list-style-type: none"> Income: patients with lower income (<\$20,000/y (mean ± SE, 1.70 ± 0.14); \$20,000–\$49,000/y (mean ± SE, 2.31 ± 0.12; P<.001) had ↑ financial burden Insurance: uninsured patients (mean ± SE, 2.67 ± 0.22; P<.001) had ↑ burden Employment: unemployed (mean ± SE, 2.57 ± 0.28) or disabled patients (mean ± SE, 2.79 ± 0.07; P<.001) had ↑ burden 	<ul style="list-style-type: none"> Age: younger patients (<50 y: mean ± SE, 2.51 ± 0.16; 50–64 y: mean ± SE, 2.13 ± 0.10; P<.001) had ↑ burden Race: black patients (mean ± SE, 1.92 ± 0.13) and other/unknown patients (mean ± SE, 2.16 ± 0.26; P=.022) had ↑ burden Marital status: married/partnered patients (mean ± SE, 1.85 ± 0.08; P=.012) had ↑ burden 	<ul style="list-style-type: none"> Chemotherapy: patients who received chemotherapy (Mean ± SE, 1.88 ± 0.06; P<.001) had ↑ burden

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

Table 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Davidoff et al, 2013 ³² • D	Generalized linear models • OOP/Income >20%	• Insurance and income: patients with supplemental insurance and higher income had ↓ OOP spending	• Age: older patients had ↓ OOP spending	• Assets, comorbidity, and type of treatment: those with assets, comorbidity, and cancer-directed radiation or antineoplastic therapy had ↑ OOP spending
Dowling et al, 2013 ³³ • I	Adjusted logistic regression • Productivity loss			• Site/Type of cancer at diagnosis: Patients with short survival cancer or multiple cancers had ↑ productivity loss ($P<.05$), were more likely to accomplish less ($P<.0001$), and be limited in the kind of work or activities engaged in because of physical health ($P<.0001$)
Goodwin et al, 2013 ³⁴ • I, D	t test • Mean % of income spent on OOP expenses • Financial burden			• Chemotherapy: patients receiving chemotherapy had ↑ mean % of income spent for OOP expenses during first 12 mo of therapy (38%) vs those not receiving chemotherapy (31%) ($t_{6,13}=2.03$; 95% CI, 0.82–12.44; $P=.025$); those receiving chemotherapy at time of survey had ↑ financial burden (52%) vs those not receiving chemotherapy. • Duration of treatment: patients who received ≥ 4 y of treatment used ↓ % of income for OOP expenses vs those with <4 y of treatment
Kent et al, 2013 ³⁵ • P, I	Adjusted analyses • Financial problems		• Age: younger patients (<40 y and those aged ≥ 65 y) ($P=.0001$) • Race/Ethnicity: non-Hispanic black, Hispanic, and other patients had ↑ financial problems vs non-Hispanic white patients ($P=.006$)	• Time since diagnosis: patients within <2 y had ↑ financial problems vs a longer period ($P<.0001$) • Recurrence/Multiple cancer: patients with presence of either had ↑ financial problems vs otherwise ($P=.049$) • Chemotherapy/Radiation: patients with either treatment(s) ↑ financial problems vs otherwise ($P<.001$)
Ramsey et al, 2013 ³⁶ • I	Frequencies • Bankruptcy		All $P<.001$; frequencies (%) represent distribution in bankruptcy vs no-bankruptcy groups • Age: younger patients (<40 y, 19% vs 7%; 40–64 y, 62% vs 48%) had ↑ bankruptcy compared with aged ≥ 65 y • Sex: female (56% vs 50%) had ↑ bankruptcy • Race/Ethnicity: nonwhite patients (14% vs 12%) had ↑ bankruptcy	• Stage at diagnosis: patients with localized (59% vs 50%) or regional stage (25% vs 22%) disease had ↑ bankruptcy vs those with distant stage (13% vs 24%) • Cancer site: patients with breast cancer (23% vs 17%) had ↑ bankruptcy; those with prostate cancer (13% vs 17%) had ↑ bankruptcy

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Wan et al, 2013 ³⁷ • I	Adjusted generalized linear modeling and linear regression • Short-term disability costs	• Insurance: HMO vs non-HMO ($P < .05$) had ↑ short-term disability costs	• Age: older patients ($P = .002$) had ↑ short-term disability costs	<ul style="list-style-type: none"> Chemotherapy: patients receiving chemotherapy ($P < .001$) had ↑ short-term disability costs Early vs metastatic stage: patients with metastatic breast cancer had ↑ short-term disability days and costs at 12 mo postdiagnosis vs those with early-stage breast cancer (41.2 ± 61.4 vs 24.7 ± 44.7 d; $\\$6,166 \pm \\$9,194$ vs $\\$3,690 \pm \\$6,673$; both $P < .001$) Early vs metastatic stage: patients with metastatic breast cancer incurred overall vs those with early-stage breast cancer ($\\$3,953$ with 95% CI, $\\$3,072$–$\\$5,086$ vs $\\$2,676$ with 95% CI, $\\$2,360$–$\\$3,034$) at $P = .009$
Meneses et al, 2012 ³⁸ • P, I	Generalized linear mixed models for repeated measures • Economic burden			<ul style="list-style-type: none"> Time since treatment: economic burden ↓ with longer time interval since treatment; mean, 2.94 economic burden items at baseline (range, 0–11) vs 2.45 at 3 mo (range, 0–13), and 2.25 at 6 mo (range, 0–14); median, 2 at baseline and 1 at 3 and 6 mo
Shankaran et al, 2012 ³⁹ • P, I	Univariate analyses • Financial hardship	<ul style="list-style-type: none"> Annual income: $\leq \\$30,000$ vs $> \\$70,000$/y (OR, 1.7; 95% CI, 0.8–3.9); $\\$30,000$–$\\$50,000$/y (OR, 4.5; 95% CI, 2.2–9.1) had ↑ hardship Insurance: Medicaid vs private had ↑ hardship (OR, 2.9; 95% CI, 1.0–8.0); Medicare had ↑ hardship (OR, 0.3; 95% CI, 0.2–0.6) Employment: presence of work disability, leave of absence, or unemployed vs employed had ↑ hardship (OR, 5.6; 95% CI, 2.4–13.3) Retired had ↓ hardship (OR, 0.4; 95% CI, 0.2–0.8) 	<ul style="list-style-type: none"> Age: patients aged < 50 y vs ≥ 75 y had ↑ hardship (OR, 30.7; 95% CI, 6.6–143.9); 50–64 y (OR, 10.2; 95% CI, 2.3–44.6) Race: nonwhite vs white patients had ↑ hardship (OR, 3.2; 95% CI, 1.5–6.7) 	
Streeter et al, 2011 ⁴⁰ • D, I	Frequencies • OOP costs	<ul style="list-style-type: none"> Type of insurance: ↑ proportion of patients who paid $> \\$500$ OOP for first oral chemotherapy claim in Medicare group compared with commercially insured group (6% vs 11%; $P < .001$) 		
Neugut et al, 2011 ⁴¹ • D, I	Frequencies • Copayment		<ul style="list-style-type: none"> Age: median copayment higher among younger patients ($\\$50$) than older patients ($\\40) 	
Markman & Luce, 2010 ⁴² • P, D	Frequencies • OOP expenses			<ul style="list-style-type: none"> Cancer site: prostate 20% with OOP $> \\$5,000$ vs breast 41%, colon 46%, lung 45%

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Pezzin et al, 2009 ⁴³ • P	Adjusted analyses • Financial problems	<ul style="list-style-type: none"> • Drug cost coverage: patients with none (OR, 4.5; 95% CI, 3.3–5.9) or partial coverage (OR, 3.6; 95% CI, 2.8–4.5) had ↑ financial problems • Income: patients with lower income (<\$15,000 vs ≥\$50,000/y [OR, 1.7; 95% CI, 1.1–2.8], \$15,000–\$29,999/y [OR, 3.1; 95% CI, 2.3–4.3], or \$30,000–\$44,999/y [OR, 1.8; 95% CI, 1.3–2.5]) had ↑ financial problems 	<ul style="list-style-type: none"> • Marital status: widowed vs married patients (OR, 0.6; 95% CI, 0.5–0.8) had ↓ financial problems 	<ul style="list-style-type: none"> • Treatment modality: patients on aromatase inhibitor vs tamoxifen (OR, 1.4; 95% CI, 1.1–1.7) had ↑ financial problems
Francoeur, 2005 ⁴⁴ • P, I, D	Structural equation modeling • Disability days • Financial strain	<ul style="list-style-type: none"> • OFFS: patients experiencing OFFS reported higher subjective financial strain ($P<.05$), difficulty paying bills ($P<.005$), and inadequacy of their insurance to meet their needs ($P<.005$) 	<ul style="list-style-type: none"> • Age: among patients with same level of financial stress, older patients perceived less financial strain from difficulty paying bills than younger patients ($P<.005$) • Age and disability days: among patients reporting above-mean disability days, older patients perceived more financial strain about adequacy of insurance and finances in the future ($P<.01$) 	
Langa et al, 2004 ⁴⁵ • D	Regression analyses • OOP expenses	<ul style="list-style-type: none"> • Income status and currently receiving treatment: patients with low income spent 27% of income on OOP expenses compared with 5% of income by patients with high income with no cancer history 		
Houts et al, 1985 ⁴⁶ • P, I, D	Regression analyses • OOP costs • Wages lost	<ul style="list-style-type: none"> • Income: patients with low income spent ↑ % of income on medical expenses ($\beta = -.21$; $P<.01$) 	<ul style="list-style-type: none"> • Age: younger patients had ↑ wages lost ($\beta = -.07$; $P<.01$), total cost ($\beta = -.028$; $P<.01$), and ↑ income lost ($\beta = -.024$; $P<.01$) • Sex: females had ↑ wages lost ($\beta = -.020$; $P<.01$) and ↑ total costs incurred ($\beta = -.016$; $P<.05$) • Distance to treatment: patients residing at greater distance from treatment facility had ↑ OOP expenses ($\beta = 0.2$; $P<.01$) 	<ul style="list-style-type: none"> • Number of outpatient visits: patients with more visits had ↑ medical costs ($\beta = 0.22$; $P<.01$) and ↑ OOP expenses ($\beta = 0.14$; $P<.05$) • Number of days in hospital: patients with more days in hospital had ↑ OOP ($\beta = 0.42$; $P<.01$), ↑ total costs ($\beta = 0.16$; $P<.05$), and ↑ % income lost ($\beta = -.02$; $P<.01$)
Smith et al, 2003 ⁴⁷ • I	Unadjusted Wilcoxon rank sum test • Nontangible costs			<ul style="list-style-type: none"> • Type of treatment: patients receiving radiation had ↑ costs vs endoscopic excision: Median number of treatments (35 vs 2), median travel distance (660 vs 150 miles), median travel time (1,440 vs 180 min), median hr work missed (76 vs 24 hr); $P<.01$

(continued on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for Financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)

Reference • Burden Type	Model • Outcome	Socioeconomic	Demographic	Clinical
Lee & Khan, 2016 ⁴⁸ • I	Adjusted logistic regression • CRN	<ul style="list-style-type: none"> Sex and insurance status: women with private health insurance (OR, 2.72; 95% CI, 1.94–3.79; $P < .0001$) and uninsured women (OR, 20.63; 95% CI, 14.94–28.47) reported ↑ CRN than men with private health insurance 	<ul style="list-style-type: none"> Sex: women reported ↑ CRN (OR, 1.27; 95% CI, 1.06–1.53) compared with men 	
Lee & Salloum, 2016 ⁴⁹ • I	Adjusted logistic regression • CRN		<ul style="list-style-type: none"> Age and race: among patients aged 18–64 y, black (OR, 1.07; 95% CI, 0.83–1.37) and Hispanic patients (OR, 1.61; 95% CI, 1.23–2.10) had ↑ CRN; among patients aged ≥65 y, black (OR, 2.64; 95% CI, 1.73–4.01) and Hispanic patients (OR, 2.07; 95% CI, 1.32–3.24) had ↑ CRN 	
Palmer et al, 2015 ²¹ • P, I	Adjusted logistic regression • Cost-related barriers to survivorship care	<ul style="list-style-type: none"> Insurance: patients with public insurance (OR, 4.82; 95% CI, 1.14–20.43) and uninsured patients (OR, 31.01; 95% CI, 2.67–360.5) reported ↑ barriers than private or both private and public insurance 		

(references on next page)

Abbreviations: ASD, adjusted score difference; CCC, comprehensive cancer center; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; CRN, cost-related nonadherence; D, direct material burdens; EE, effect estimate; FPL, federal poverty level; GPH, general public hospital; HMO, health maintenance organization; I, indirect burdens; MCS, Mental Component Score; OFFS, objective family financial stress; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PCS, Physical Component Score; QoL, quality of life.

eTable 4. Risk Factors of Financial Toxicity Burdens (cont.)**References**

- Wheeler SB, Spencer JC, Pinheiro LC, et al. Financial impact of breast cancer in black versus white women. *J Clin Oncol* 2018;36:1695–1701.
- de Souza JA, Yap BJ, Wroblewski K, et al. Measuring financial toxicity as a clinically relevant patient-reported outcome: the validation of the Comprehensive Score for financial Toxicity (COST). *Cancer* 2017;123:476–484.
- Meeker CR, Wong YN, Egleston BL, et al. Distress and financial distress in adults with cancer: an age-based analysis. *J Natl Compr Canc Netw* 2017;15:1224–1233.
- Pisu M, Azuero A, Benz R, et al. Out-of-pocket costs and burden among rural breast cancer survivors. *Cancer Med* 2017;6:572–581.
- Abel GA, Albelda R, Khera N, et al. Financial hardship and patient-reported outcomes after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016;22:1504–1510.
- Banegas MP, Guy GP Jr, de Moor JS, et al. For working-age cancer survivors, medical debt and bankruptcy create financial hardships. *Health Aff (Millwood)* 2016;35:54–61.
- Chongpison Y, Hornbrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
- Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
- Lathan CS, Cronin A, Tucker-Seeley R, et al. Association of financial strain with symptom burden and quality of life for patients with lung or colorectal cancer. *J Clin Oncol* 2016;34:1732–1740.
- Narang AK, Nicholas LH. Out-of-pocket spending and financial burden among Medicare beneficiaries with cancer. *JAMA Oncol* 2017;3:757–765.
- Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
- Whitney RL, Bell JF, Reed SC, et al. Predictors of financial difficulties and work modifications among cancer survivors in the United States. *J Cancer Surviv* 2016;10:241–250.
- Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
- Yabroff KR, Dowling EC, Guy GP Jr, et al. Financial hardship associated with cancer in the United States: findings from a population-based sample of adult cancer survivors. *J Clin Oncol* 2016;34:259–267.
- Zheng Z, Yabroff KR, Guy GP Jr, et al. Annual medical expenditure and productivity loss among colorectal, female breast, and prostate cancer survivors in the United States. *J Natl Cancer Inst* 2015;108:djv382.
- Callahan C, Brintzenhofszoc K. Financial quality of life for patients with cancer: an exploratory study. *J Psychosoc Oncol* 2015;33:377–394.
- Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
- Guy GP Jr, Yabroff KR, Ekwueme DU, et al. Healthcare expenditure burden among non-elderly cancer survivors, 2008–2012. *Am J Prev Med* 2015;49(Suppl 5):S489–497.
- Huntington SF, Weiss BM, Vogl DT, et al. Financial toxicity in insured patients with multiple myeloma: a cross-sectional pilot study. *Lancet Haematol* 2015;2:e408–416.
- Jones WC, Parry C, Devine S, et al. Prevalence and predictors of distress in posttreatment adult leukemia and lymphoma survivors. *J Psychosoc Oncol* 2015;33:124–141.
- Palmer NR, Weaver KE, Hauser SP, et al. Disparities in barriers to follow-up care between African American and White breast cancer survivors. *Support Care Cancer* 2015;23:3201–3209.
- Pisu M, Kenzik KM, Oster RA, et al. Economic hardship of minority and non-minority cancer survivors 1 year after diagnosis: another long-term effect of cancer? *Cancer* 2015;121:1257–1264.
- Tucker-Seeley RD, Abel GA, Uno H, et al. Financial hardship and the intensity of medical care received near death. *Psychooncology* 2015;24:572–578.
- Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
- Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
- Jagsi R, Hawley ST, Abrahamse P, et al. Impact of adjuvant chemotherapy on long-term employment of survivors of early-stage breast cancer. *Cancer* 2014;120:1854–1862.
- Jagsi R, Pottow JAE, Griffith KA, et al. Long-term financial burden of breast cancer: experiences of a diverse cohort of survivors identified through population-based registries. *J Clin Oncol* 2014;32:1269–1276.
- Khera N, Chang YH, Hashmi S, et al. Financial burden in recipients of allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2014;20:1375–1381.
- Martin MY, Fouad MN, Oster RA, et al. What do cancer patients worry about when making decisions about treatment? Variation across racial/ethnic groups. *Support Care Cancer* 2014;22:233–244.
- Reegenbogen SE, Veenstra CM, Hawley ST, et al. The personal financial burden of complications after colorectal cancer surgery. *Cancer* 2014;120:3074–3081.
- Veenstra CM, Reegenbogen SE, Hawley ST, et al. A composite measure of personal financial burden among patients with stage III colorectal cancer. *Med Care* 2014;52:957–962.
- Davidoff AJ, Erten M, Shaffer T, et al. Out-of-pocket health care expenditure burden for Medicare beneficiaries with cancer. *Cancer* 2013;119:1257–1265.
- Dowling EC, Chawla N, Forsythe LP, et al. Lost productivity and burden of illness in cancer survivors with and without other chronic conditions. *Cancer* 2013;119:3393–3401.
- Goodwin JA, Coleman EA, Sullivan E, et al. Personal financial effects of multiple myeloma and its treatment. *Cancer Nurs* 2013;36:301–308.
- Kent EE, Forsythe LP, Yabroff KR, et al. Are survivors who report cancer-related financial problems more likely to forgo or delay medical care? *Cancer* 2013;119:3710–3717.
- Wan Y, Gao X, Mehta S, et al. Washington State cancer patients found to be at greater risk for bankruptcy than people without a cancer diagnosis. *Health Aff (Millwood)* 2013;32:1143–1152.
- Meneses K, Azuero A, Hassey L, et al. Does economic burden influence quality of life in breast cancer survivors? *Gynecol Oncol* 2012;124:437–443.
- Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
- Streeter SB, Schwartzberg L, Husain N, et al. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *Am J Manag Care* 2011;17(Suppl 5):SP38–44.
- Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
- Markman M, Luce R. Impact of the cost of cancer treatment: an internet-based survey. *J Oncol Pract* 2010;6:69–73.
- Pezzin LE, O'Neil MB, Nattinger AB. The economic consequences of breast cancer adjuvant hormonal treatments. *J Gen Intern Med* 2009;24(Suppl 2):S446–450.
- Francœur RB. Cumulative financial stress and strain in palliative radiation outpatients: the role of age and disability. *Acta Oncol* 2005;44:369–381.
- Langa KM, Fendrick AM, Chernew ME, et al. Out-of-pocket health-care expenditures among older Americans with cancer. *Value Health* 2004;7:186–194.
- Houts PS, Harvey HA, Simmonds MA, et al. Characteristics of patients at risk for financial burden because of cancer and its treatment. *J Psychosoc Oncol* 1985;3:15–22.
- Smith JC, Johnson JT, Cognetti DM, et al. Quality of life, functional outcome, and costs of early glottic cancer. *Laryngoscope* 2003;113:68–76.
- Lee M, Khan MM. Gender differences in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:384–393.
- Lee M, Salloum RG. Racial and ethnic disparities in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:534–544.

Table 5. Financial Toxicity Burdens and Health Outcomes

Reference • Burden Type	Statistical Analysis • Financial Toxicity Burden Measure	Dependent Variable and/or Outcome Measures • Association of Financial Toxicity Burden Measure With Outcome Measure(s)
Financial toxicity burden and adherence to cancer treatment		
Farias & Du, 2017 ¹ • D	Adjusted logistic regression • OOP costs	<ul style="list-style-type: none"> Adherence to adjuvant endocrine therapy (aromatase inhibitor or tamoxifen) ↑ OOP was associated with ↓ adherence Compared with OOP cost \$0–\$2.65 (reference), OR of adherence for OOP cost \$2.66–\$10.00: OR, 0.81; 95% CI, 0.71–0.94; OOP cost \$10.01–\$41.25: OR, 0.65, 95% CI, 0.56–0.76; OOP cost \$41.26–\$105.55: OR, 0.21, 95% CI, 0.18–0.24; OOP cost >\$105.55: OR, 0.31; 95% CI, 0.26–0.37
Farias et al, 2018 ² • D	Adjusted logistic regression • OOP costs	<ul style="list-style-type: none"> Adherence to adjuvant endocrine therapy (aromatase inhibitor or tamoxifen) ↑ OOP was associated with ↓ adherence Compared with OOP cost \$0–\$4.99 (reference), RR of adherence for OOP cost of \$5.00–\$9.99: RR, 1.01; 95% CI, 0.97–1.05; OOP cost \$10.00–\$14.99: RR, 0.93; 95% CI, 0.88–0.98; OOP cost \$15.00–\$19.99: RR, 0.92; 95% CI, 0.85–0.98; OOP cost ≥\$20.00: RR, 0.94; 95% CI, 0.89–0.99
Bender et al, 2014 ³ • P, I	Effect modifier in conditional regression • Financial hardship	<ul style="list-style-type: none"> Adherence to endocrine therapy ↑ financial hardship measured by OOP cost perception was effect modifier of association between negative mood and adherence to endocrine therapy ($P < .001$)
Bestvina et al, 2014 ⁴ • P, I	Adjusted logistic regression • Financial burden	<ul style="list-style-type: none"> Medication nonadherence Nonadherence ↑ among patients who had cost discussions (OR, 2.58; 95% CI, 1.14–5.85), experienced ↑ financial burden (OR, 2.89; 95% CI, 1.41–5.89) and experienced ↑ financial distress (OR, 1.64; 95% CI, 1.38–1.96)
Dusetzina et al, 2014 ⁵ • D	Generalized estimating equations • Copayment	<ul style="list-style-type: none"> Discontinuation: 17% of patients with 1 copayment discontinued therapy during first 180 days; with increased risk of discontinuation compared with patients with ↓ copayment (RR = 1.70; 95% CI, 1.30–2.22) Nonadherence: 30% of patients with 1 copayment were nonadherent to therapy; with increased risk of nonadherence (RR = 1.42; 95% CI, 1.19–1.69)
Jagsi et al, 2014 ⁶ • P, I	Frequency	<ul style="list-style-type: none"> Treatment nonadherence: In patients with breast cancer Frequency of going without medications (5%), taking less than recommended treatment (4%), and missing doctor's appointments (8%)
Zafar et al, 2013 ⁷ • D	Frequency	<ul style="list-style-type: none"> Healthcare noncompliance: in patients with cancer needing copayment assistance Frequency of taking less than prescribed medications (20%), filling part of prescription (19%), avoiding filling prescription (24%), avoiding procedures (7%) or tests (9%), spreading out chemotherapy/appointments (7%), and skipping appointments (4%)
Zullig et al, 2013 ⁸ • P, I	Frequency and adjusted logistic regression • Subjective financial distress	<ul style="list-style-type: none"> Prescription medication nonadherence Overall nonadherence 45%, of which 4% took medications prescribed for another person, 22% took less than prescribed amount, 25% filled only part of prescription, and 27% did not fill the prescription because of medication cost. In subset of nonadherent patients: 69% asked doctor for less expensive medicine, 66% reduced spending on basics like food or clothing, 26% purchased over-the-counter product, 58% shopped for lowest priced medication, 68% borrowed money or used credit card to pay, and 69% obtained medication samples. Although statistically significant in unadjusted analyses, in adjusted analysis, ↑ subjective financial distress not associated with medication nonadherence (OR, 1.60; 95% CI, 0.71–3.60; $P = .26$).
Shankaran et al, 2012 ⁹ • P, I	Adjusted logistic regression • Perceived financial burden	<ul style="list-style-type: none"> Economically motivated treatment nonadherence Nonadherence ↑ among younger <50 y ($P = .04$), low income ≤\$30,000/y ($P = .012$), and unemployed/disabled/leave of absence ($P = .05$)

(continued on next page)

Abbreviations: AMD, adjusted mean difference; BSI, Brief Symptom Inventory; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; D, direct material burdens; FACT-B, Functional Assessment of Cancer Therapy-Breast tool; FACT-G, Functional Assessment of Cancer Therapy-General tool; HR, hazard ratio; I, indirect burdens; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PHQ, Patient Health Questionnaire; QoL, quality of life; RR, risk ratio; SE, standard error.

eTable 5. Financial Toxicity Burdens and Health Outcomes (cont.)

Reference • Burden Type	Statistical Analysis • Financial Toxicity Burden Measure	Dependent Variable and/or Outcome Measures • Association of Financial Toxicity Burden Measure With Outcome Measure(s)
Financial toxicity burden and adherence to cancer treatment (cont.)		
Neugut et al, 2011 ¹⁰ • D, I	Adjusted logistic regression and Cox proportional hazard analyses • Copayment	<ul style="list-style-type: none"> Nonpersistence and nonadherence to recommended medications ↑ copayment associated with ↑ nonpersistence and ↑ nonadherence Among women of Medicare-eligible age, compared with copayment <\$30 (ref), OR of persistence for copayment of \$30–\$89.99: OR, 0.69; 95% CI, 0.62–0.75 Among women aged <65 y, compared with <\$30 (ref), copayment of >\$90: OR, 0.72; 95% CI, 0.65–0.80 Among women of Medicare-eligible age, compared with copayment <\$30 (ref), OR of adherence \$30–\$89.99: OR, 0.83; 95% CI, 0.72–0.96; >\$90: OR, 0.70; 95% CI, 0.60–0.82
Streeter et al, 2011 ¹¹ • D, I	Adjusted logistic regression • Cost-sharing	<ul style="list-style-type: none"> Abandonment of oral or intravenous cancer treatment ↑ cost sharing associated with ↑ abandonment Compared with cost sharing of ≤\$100 (ref), OR of abandonment for \$101–\$150: OR, 1.84; 95% CI, 1.23–2.75; \$151–\$200: OR, 1.51; 95% CI, 0.97–2.34; \$201–\$250: OR, 2.30; 95% CI, 1.31–4.04; \$251–\$350: OR, 2.31; 95% CI, 1.59–3.36; cost sharing \$351–\$500: OR, 3.28; 95% CI, 2.20–4.88; cost sharing >\$500: OR, 4.46; 95% CI, 3.80–5.22
Markman & Luce, 2010 ¹² • P, D	Frequency • Financial impact	<ul style="list-style-type: none"> Not to receive a recommended treatment 10% of patients decided not to receive a treatment because it was too expensive; 25% for patients with annual income <\$40,000
Pezzin et al, 2009 ¹³ • P	Adjusted logistic regression • Insurance prescription coverage	<ul style="list-style-type: none"> Making appropriate switch between adjuvant endocrine therapies (from tamoxifen to aromatase inhibitor) Compared with full drug coverage, women with no prescription drug coverage ↓ likelihood of making appropriate switch to aromatase inhibitor: OR, 0.7; 95% CI, 0.5–0.9; partial prescription drug coverage: OR, 0.8; 95% CI, 0.7–1.0 Women near poverty (income \$15,000–\$29,000/y) had ↓ likelihood of making appropriate switch: OR, 0.6; 95% CI, 0.4–0.9
Huntington et al, 2015 ¹⁴ • P	Fisher exact test • Financial burden (COST tool)	<ul style="list-style-type: none"> Patients with ↑ financial burden (COST score ≤23) more likely to: <ul style="list-style-type: none"> Delay initiation of myeloma treatment: 29% (P=.003) Fill only part of myeloma therapy prescription: 22% (P=.0077) Fill only part of noncancer medication: 16% (P=.0019) Stop myeloma therapy prescription: 22% (P=.0011) Stop noncancer medication: 20% (P=.0077) Refuse recommended test because of cost: 18% (P=.016) Skip clinic visits: 12% (P=.027)
Financial toxicity burden and adherence to general healthcare		
Nipp et al, 2016 ¹⁵ • I	Frequency	<ul style="list-style-type: none"> Use of healthcare-altering cost-coping strategies Overall, 39% used ≥1 cost-coping strategies, 28% did not fill prescriptions, 22% took less than recommended medications, 10% missed a medical test, 8% missed a procedure, and 6% missed an appointment.
Guy et al, 2015 ¹⁶ • D	Adjusted logistic regression • OOP burden among cancer survivors	<ul style="list-style-type: none"> Unable to obtain necessary medical care: patients with ↑ OOP burden had ↑ inability to obtain necessary medical care (19.2% vs 12.5%, P=.002) Delaying necessary medical care: patients with ↑ OOP burden ↑ delayed medical care (21.6% vs 13.8%, P=.002) Lower breast cancer screening: patients with ↑ OOP burden had ↓ screening rates among age-appropriate women (63.2% vs 75.9%, P=.02)
Khera et al, 2014 ¹⁷ • P, I	Frequency and adjusted logistic regression • Financial burden	<ul style="list-style-type: none"> Cost-related treatment nonadherence Overall, 21% reported not making a physician appointment or having a medical test performed due to cost Use of medical service, eg, physical therapy, deferred by 28% due to cost Younger patients (OR for per 5-y increase in age: 0.82; 95% CI, 0.6–0.9) and lower educational level (OR for ≤ high school vs 4-y degree: OR, 5.0; 95% CI, 1.7–15) more likely to exhibit deleterious health behaviors (including treatment nonadherence) due to cost concerns

(continued on next page)

Abbreviations: AMD, adjusted mean difference; BSI, Brief Symptom Inventory; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; D, direct material burdens; FACT-B, Functional Assessment of Cancer Therapy-Breast tool; FACT-G, Functional Assessment of Cancer Therapy-General tool; HR, hazard ratio; I, indirect burdens; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PHQ, Patient Health Questionnaire; QoL, quality of life; RR, risk ratio; SE, standard error.

eTable 5. Financial Toxicity Burdens and Health Outcomes (cont.)

Reference • Burden Type	Statistical Analysis • Financial Toxicity Burden Measure	Dependent Variable and/or Outcome Measures • Association of Financial Toxicity Burden Measure With Outcome Measure(s)
Financial toxicity burden and adherence to general healthcare (cont.)		
Kent et al, 2013 ¹⁸ • P, I	Frequency and adjusted logistic regression • Financial problems	<ul style="list-style-type: none"> • Delay/Forgo care • Patients reporting ↑ financial problems ↑ likely to report delaying medical care (18.5% vs 7.4%, $P < .0001$) or forgoing medical care (13.5% vs 5.1%; $P < .0001$), prescription medications (13.8% vs 7.7%, $P = .001$), dental care (20.0% vs 8.3%; $P < .0001$), eyeglasses (13.7% vs 5.9%; $P < .0001$), or mental health care (9.7% vs 7.1%; $P = .03$) compared with patients without financial problems
Khera et al, 2011 ¹⁹ • P, I	Frequency and ordinal regression analyses • Concerns about medical costs	<ul style="list-style-type: none"> • Adherence to preventive healthcare • ↑ cost concerns were associated with ↓ adherence to preventive health care: cutting back on prescribed medication (10%), not purchasing medication (11%), avoiding doctor's appointments (15%), not using physical therapy (13%), and not having a medical test performed (12%); 26% reported ≥ 1 behavior • ↑ concerns associated with ↓ adherence rates (OR, 1.48; 95% CI, 1.15–1.89)
Weaver et al, 2010 ²⁰ • I	Frequency	<ul style="list-style-type: none"> • Nonadherence to general healthcare • Forgo care (frequency, 8%), delay care (11%), forgo prescription medications (10%), forgo dental care (11%), forgo mental health care (3%)
Financial toxicity burden and healthcare decision-making		
DiMartino et al, 2017 ²¹ • I	Binary logit model • Financial hardship	<ul style="list-style-type: none"> • Having survivorship care planning discussions with provider • ↑ financial hardship associated with ↓ discussions with β coefficient, 0.74; SE, 0.30; mean marginal effect, 6.6%; $P < .05$
Wong et al, 2016 ²² • P	Adjusted linear regression • Financial concern	<ul style="list-style-type: none"> • Self-efficacy: patients with ↑ financial concern had ↓ self-efficacy in treatment-related decision-making ($P = .004$) • Preparation for decision-making: patients with ↑ financial concern had ↓ preparation for decision-making ($P < .001$) • Subjective distress: patients with ↑ financial concern had ↑ distress ($P < .001$) • Decisional conflict: patients with ↑ financial concern had ↑ decisional conflict ($P < .001$)
Overall survival		
Ramsey et al, 2016 ²³ • I	Proportional hazards regression • Bankruptcy	<ul style="list-style-type: none"> • Overall survival • All-cause mortality ↑ in patients who filed for bankruptcy vs patients who did not: HR, 1.79; 95% CI, 1.64–1.96
QoL		
de Souza et al, 2017 ²⁴ • P	Pearson correlation • Financial toxicity (COST tool)	<ul style="list-style-type: none"> • Health-related QoL • Measured using two scales: FACT-G and EORTC-QoL • Adjusted correlation between COST measure and both FACT-G and EORTC-QoL statistically significant • FACT-G $r = 0.42$ ($P < .001$); EORTC-QoL $r = 0.33$ ($P < .001$)
Abel et al, 2016 ²⁵ • P, I	Adjusted logistic regression • Financial hardship	<ul style="list-style-type: none"> • QoL below median: patients experiencing any hardship (OR, 2.90; 95% CI, 1.7–4.9) or all 3 hardships (OR, 2.16; 95% CI, 0.99–4.7) had ↑ odds of QoL below median. • Self-reported health below median: patients experiencing any hardship (OR, 2.18; 95% CI, 1.3–3.6) had ↑ odds of self-reported health below median. • Perceived stress above median: patients experiencing any hardship (OR, 2.08; 95% CI, 1.3–3.5) or all 3 hardships (OR, 3.14; 95% CI, 1.4–6.8) had ↑ odds of perceived stress above median.
Chongpison et al, 2016 ²⁶ • P	Adjusted linear regression • Financial burden	<ul style="list-style-type: none"> • Depression: survivors with self-reported depression (in past 30 days from survey) perceived ↑ financial burden (adjusted mean, 3.01; 95% CI, 2.57–3.45) compared with patients reporting no depression (adjusted mean, 1.67; 95% CI, 1.43–1.91)

(continued on next page)

Abbreviations: AMD, adjusted mean difference; BSI, Brief Symptom Inventory; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; D, direct material burdens; FACT-B, Functional Assessment of Cancer Therapy-Breast tool; FACT-G, Functional Assessment of Cancer Therapy-General tool; HR, hazard ratio; I, indirect burdens; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PHQ, Patient Health Questionnaire; QoL, quality of life; RR, risk ratio; SE, standard error.

eTable 5. Financial Toxicity Burdens and Health Outcomes (cont.)

Reference • Burden Type	Statistical Analysis • Financial Toxicity Burden Measure	Dependent Variable and/or Outcome Measures • Association of Financial Toxicity Burden Measure With Outcome Measure(s)	
		QoL (cont.)	
Delgado-Guay et al, 2016 ²⁷ • P	Adjusted logistic regression • Financial distress	• <u>Spiritual pain</u> : associated with ↑ financial distress (OR, 1.22; 95% CI, 1.11–1.34)	
	Adjusted linear regression • Financial burden	• Patients with ↑ financial burden had ↓ QoL based on Short-Form Health Survey • Physical component score: ↓ score (β, −2.45; 95% CI, −3.75 to −1.15) • Mental component score: ↓ score (β, −3.05; 95% CI, −4.42 to −1.67) • Depressed mood: ↑ depression (OR, 1.95; 95% CI, 1.29–2.95) • Cancer recurrence—related worries: ↑ worries (OR, 3.54; 95% CI, 2.65–4.72)	
Kale & Carroll, 2016 ²⁸ • P, I	Adjusted linear regression • Financial burden	• QoL based on FACT-B • ↑ ability to keep up with work responsibilities was associated with ↑ QoL scores (estimate, 6.70; SE, 1.26; $P < .001$) • ↑ financial worries were associated with ↓ QoL scores (estimate = −5.33; SE, 0.83; $P < .001$) • Difficulty paying for basics was associated with ↓ QoL (estimate = −7.01; SE, 2.83; $P = .01$)	
	Adjusted ordinal logistic regression • Financial reserve	• Symptom burden based on EORTC C30, pain based on Brief Pain Inventory, QoL based on EuroQoL-5 dimension scale for patients with lung cancer and CRC • Symptom burden: patients with ↓ financial reserves (<12 mo) reported ↑ symptom burden (lung cancer: AMD, 5.25; 95% CI, 3.29–7.04; CRC: AMD, 5.31; 95% CI, 3.58–7.04) • Pain experienced: patients with ↓ financial reserves (<12 mo) had ↑ pain (lung cancer: AMD, 5.03; 95% CI, 3.29–7.22; CRC: AMD, 3.45; 95% CI, 1.25–5.66) • QoL: patients with ↓ financial reserves (<12 mo) had ↓ QoL (lung cancer: AMD, 4.70; 95% CI, 2.82–6.58; CRC: AMD, 5.22; 95% CI, 3.61–6.82)	
Lathan et al, 2016 ³⁰ • I	Adjusted logistic regression • Financial reserve	• Overall distress: 65% reported significant distress scores (≥ 4), 28% had high scores (8–10) • Emotional distress: 66% had ≥ 1 emotional symptom; 49% reported worry, 38% nervousness, 37% fear, 30% sadness, 29% depression, 26% loss of interest • Financial distress associated with overall distress with emotional distress mediating this association; a 1-unit improvement in financial well-being led to a 0.553-unit decrease in overall distress ($P < .001$).	
	Frequency and structural equation modeling • Financial distress/well-being (lnCharge scale)	• QoL based on FACT-G • Financial distress inversely correlated with QoL ($r = -0.23$; $P = .0057$)	
Meeker et al, 2016 ³¹ • P	Spearman correlation • Financial distress	• QoL based on EuroQoL EQ-5D • ↑ financial distress was associated with ↓ QoL (adjusted $\beta = -0.06$ per burden category; $P < .001$)	
	Adjusted structural equation modeling • Financial distress	• Patients reporting “a lot” vs no financial problems had more poor physical health ($P < .001$) • More poor mental health ($P < .001$) • Poorer satisfaction with social activities and relationships ($P < .001$) • Overall QoL: Compared with patients with no financial problems, patients reporting “a lot” of financial problems had lower odds of QoL $\geq$ good (OR, 0.24; 95% CI, 0.14–0.40); reporting “some” financial problems: OR, 0.57; 95% CI, 0.33–0.99; reporting “a few” financial problems: OR, 0.67; 95% CI, 0.41–1.08)	
Delgado-Guay et al, 2015 ³² • P, I	Spearman correlation • Financial distress		
Zafar et al, 2015 ³³ • P	Adjusted structural equation modeling • Financial distress		
Fenn et al, 2014 ³⁴ • P	Adjusted logistic regression • Financial problems		

Abbreviations: AMD, adjusted mean difference; BSI, Brief Symptom Inventory; COST, Comprehensive Score for Financial Toxicity; CRC, colorectal cancer; D, direct material burdens; FACT-B, Functional Assessment of Cancer Therapy-Breast tool; FACT-G, Functional Assessment of Cancer Therapy-General tool; HR, hazard ratio; I, indirect burdens; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PHQ, Patient Health Questionnaire; QoL, quality of life; RR, risk ratio; SE, standard error.

(continued on next page)

eTable 5. Financial Toxicity Burdens and Health Outcomes (cont.)

Reference • Burden Type	Statistical Analysis • Financial Toxicity Burden Measure	Dependent Variable and/or Outcome Measures • Association of Financial Toxicity Burden Measure With Outcome Measure(s)	
		QoL (cont.)	
Meneses et al, 2012 ²⁵ • P, I	Structural equation modeling analyses • Economic events	• QoL: ↑ in economic events significantly associated with poorer QoL at all time points (baseline, 3-mo, and 6-mo follow-up)	
		• At 6-mo time point, proportion of survivors reporting change in motivation ($P=.016$), productivity ($P=.002$), quality of work ($P=.01$), days missed from work ($P<.001$) and sacrificing other things ($P=.001$) as various domains of QoL declined compared with baseline.	
Khara et al, 2011 ¹⁹ • P, I	Adjusted logistic regression • Cost concerns	• Physical and mental functioning	
		• ↑ Concerns about medical costs associated with ↓ physical (OR, 2.68; 95% CI, 2.03–3.53) and mental functioning (OR, 2.32; 95% CI, 1.65–3.27)	
de Moor et al, 2010 ³⁵ • P	Analysis of variance; adjusted linear regression • Financial status	• Women with better self-rated financial status experienced decline in depression and anxiety over time while women with medium or low financial status experienced increase in both over time	
		• Depression at 9-mo follow-up: ↑ financial status associated with ↓ depression ($F\text{-test}=9.84$; $df=2,349$; $P=.0001$)	
Ell et al, 2008 ³⁷ • P, I	Adjusted generalized estimating equations modeling • Cost concerns/worry of lost wages	• Anxiety at 9-mo follow-up: ↑ financial status associated with ↓ anxiety ($F\text{-test}=10.17$; $df=2,346$; $P=.0001$)	
		• QoL measured using FACT-G, PHQ, and BSI	
		• Functional well-being QoL: patients with medical cost concerns or worries of wages lost at baseline and financial stress at 12 mo time point had poorer functional well-being	
		• Emotional well-being QoL: patients with medical cost concerns or worries of wages lost at baseline and financial stress at 12 mo time point had poorer emotional well-being	
		• Physical well-being QoL: patients with medical cost concerns at baseline and financial stress at 12 mo time point had poorer physical well-being	
		• Social-family well-being QoL: patients with wage worries at baseline and financial stress at 12 mo time point had poorer social-family well-being	
		• Depression (PHQ-9) and anxiety (BSI-A) scales: patients with medical cost concerns or worries of wages lost had higher scores on both scales	
		• Major depression (PHQ-9≥10): patients with wage-related worries (OR, 2.12; 95% CI, 1.19–3.76) and financial stress (OR, 2.13; 95% CI, 1.14–3.99) more likely to meet the criteria for major depression	
Gupta et al, 2007 ³⁸ • P	Adjusted linear regression • Financial difficulty	• QoL: 10-unit ↑ in financial difficulty score associated with ↓ in QoL score (Ferrans and Powers Quality of Life Index); Overall QoL: Δ score, –0.40; 95% CI, –0.48 to –0.32; $P<.001$	
		• Health and physical: Δ score, –0.52; 95% CI, –0.64 to –0.41; $P<.001$	
		• Social and economic: Δ score, –0.36; 95% CI, –0.44 to –0.29; $P<.001$	
		• Psychological and spiritual: Δ score, –0.32; 95% CI, –0.44 to –0.21; $P<.001$	
		• Family: Δ score, –0.24; 95% CI, –0.33 to –0.14; $P<.001$	

Abbreviations: AMD, adjusted mean difference; BSI, Brief Symptom Inventory; COST, Comprehensive Score for financial Toxicity; CRC, colorectal cancer; D, direct material burdens; FACT-B, Functional Assessment of Cancer Therapy-Breast tool; FACT-G, Functional Assessment of Cancer Therapy-General tool; HR, hazard ratio; I, indirect burdens; OOP, out of pocket; OR, odds ratio; P, psychological burdens; PHQ, Patient Health Questionnaire; QoL, quality of life; RR, risk ratio; SE, standard error.

(references on next page)

eTable 5. Financial Toxicity Burdens and Health Outcomes (cont.)**References**

1. Farias AJ, Du XL. Association between out-of-pocket costs, race/ethnicity, and adjuvant endocrine therapy adherence among Medicare patients with breast cancer. *J Clin Oncol* 2017;35:86–95.
2. Farias AJ, Hansen RN, Zeliadt SB, et al. The association between out-of-pocket costs and adherence to adjuvant endocrine therapy among newly diagnosed breast cancer patients. *Am J Clin Oncol* 2018;41:708–715.
3. Bender CM, Gentry AL, Brufsky AM, et al. Influence of patient and treatment factors on adherence to adjuvant endocrine therapy in breast cancer. *Oncol Nurs Forum* 2014;41:274–285.
4. Bestvina CM, Zullig LL, Rushing C, et al. Patient-oncologist cost communication, financial distress, and medication adherence. *J Oncol Pract* 2014;10:162–167.
5. Dusetzina SB, Winn AN, Abel GA, et al. Cost sharing and adherence to tyrosine kinase inhibitors for patients with chronic myeloid leukemia. *J Clin Oncol* 2014;32:306–311.
6. Jaggi R, Pottow JAE, Griffith KA, et al. Long-term financial burden of breast cancer: experiences of a diverse cohort of survivors identified through population-based registries. *J Clin Oncol* 2014;32:1269–1276.
7. Zafar SY, Peppercorn JM, Schrag D, et al. The financial toxicity of cancer treatment: a pilot study assessing out-of-pocket expenses and the insured cancer patient's experience. *Oncologist* 2013;18:381–390.
8. Zullig LL, Peppercorn JM, Schrag D, et al. Financial distress, use of cost-coping strategies, and adherence to prescription medication among patients with cancer. *J Oncol Pract* 2013;9(Suppl 6):60s–63s.
9. Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
10. Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
11. Streeter SB, Schwartzberg L, Husain N, et al. Patient and plan characteristics affecting abandonment of oral oncolytic prescriptions. *Am J Manag Care* 2011;17(Suppl 5):SP38–44.
12. Markman M, Luce R. Impact of the cost of cancer treatment: an internet-based survey. *J Oncol Pract* 2010;6:69–73.
13. Pezzin LE, O'Neil MB, Nattinger AB. The economic consequences of breast cancer adjuvant hormonal treatments. *J Gen Intern Med* 2009;24(Suppl 2):S446–450.
14. Huntington SF, Weiss BM, Vogl DT, et al. Financial toxicity in insured patients with multiple myeloma: a cross-sectional pilot study. *Lancet Haematol* 2015;2:e408–416.
15. Nipp RD, Zullig LL, Samsa G, et al. Identifying cancer patients who alter care or lifestyle due to treatment-related financial distress. *Psychooncology* 2016;25:719–725.
16. Guy GP Jr, Yabroff KR, Ekwueme DU, et al. Healthcare expenditure burden among non-elderly cancer survivors, 2008–2012. *Am J Prev Med* 2015;49(Suppl 5):S489–497.
17. Khera N, Chang YH, Hashmi S, et al. Financial burden in recipients of allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2014;20:1375–1381.
18. Kent EE, Forsythe LP, Yabroff KR, et al. Are survivors who report cancer-related financial problems more likely to forgo or delay medical care? *Cancer* 2013;119:3710–3717.
19. Khera N, Chow EJ, Leisenring WM, et al. Factors associated with adherence to preventive care practices among hematopoietic cell transplantation survivors. *Biol Blood Marrow Transplant* 2011;17:995–1003.
20. Weaver KE, Rowland JH, Bellizzi KM, et al. Forgoing medical care because of cost: assessing disparities in healthcare access among cancer survivors living in the United States. *Cancer* 2010;116:3493–3504.
21. DiMartino LD, Birken SA, Mayer DK. The relationship between cancer survivors' socioeconomic status and reports of follow-up care discussions with providers. *J Cancer Educ* 2017;32:749–755.
22. Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
23. Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
24. de Souza JA, Yap BJ, Wroblewski K, et al. Measuring financial toxicity as a clinically relevant patient-reported outcome: the validation of the Comprehensive Score for financial Toxicity (COST). *Cancer* 2017;123:476–484.
25. Abel GA, Albelda R, Khera N, et al. Financial hardship and patient-reported outcomes after hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016;22:1504–1510.
26. Chongpison Y, Hombrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
27. Delgado-Guay MO, Chisholm G, Williams J, et al. Frequency, intensity, and correlates of spiritual pain in advanced cancer patients assessed in a supportive/palliative care clinic. *Palliat Support Care* 2016;14:341–348.
28. Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
29. Keim-Malpass J, Levine B, Danhauer SC, et al. Work-related perceptions and quality of life among breast cancer survivors. *Psychooncology* 2016;25:873–876.
30. Lathan CS, Cronin A, Tucker-Seeley R, et al. Association of financial strain with symptom burden and quality of life for patients with lung or colorectal cancer. *J Clin Oncol* 2016;34:1732–1740.
31. Meeker CR, Geynisman DM, Eggleston BL, et al. Relationships among financial distress, emotional distress, and overall distress in insured patients with cancer. *J Oncol Pract* 2016;12:e755–764.
32. Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
33. Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
34. Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
35. Meneses K, Azuero A, Hassey L, et al. Does economic burden influence quality of life in breast cancer survivors? *Gynecol Oncol* 2012;124:437–443.
36. de Moor JS, Partridge AH, Winer EP, et al. The role of socioeconomic status in adjustment after ductal carcinoma in situ. *Cancer* 2010;116:1218–1225.
37. Ell K, Xie B, Wells A, et al. Economic stress among low-income women with cancer: effects on quality of life. *Cancer* 2008;112:616–625.
38. Gupta D, Lis CG, Grutsch JF. Perceived cancer-related financial difficulty: implications for patient satisfaction with quality of life in advanced cancer. *Support Care Cancer* 2007;15:1051–1056.

eAppendix 1. Supplementary Results

Study Characteristics

Patients with cancer at any disease site (eg, breast, prostate, lung) were eligible in 38 studies (51%). Single-disease site-specific studies most commonly focused on breast cancer (n=14 studies, representing 46,033 study subjects) and hematologic cancers (n=7 studies, representing 7,005 study subjects). Assessments of the measure of financial toxicity represented a variety of time points across the cancer survivorship spectrum, from diagnosis to treatment to follow-up; 7 studies focused on financial toxicity within the first 12 months of cancer treatment,¹⁻⁷ and 6 studies were longitudinal in design.⁸⁻¹³ A total of 5 studies (74%) were cross-sectional in design; 3 studies (4%) focused on subjects aged ≥ 65 years,¹⁴⁻¹⁶ and 7 studies oversampled for racial/ethnic minorities or medically underserved patients.^{4,8,17-21} For sample-weighted prevalence estimates, subgroup analyses demonstrating the prevalence of financial burdens ranged from 39% to 64%. When data were grouped by the specific type of financial burden, prevalence was as follows:

- In 7 studies reporting psychological burdens, 39% (95% CI, 25%–54%).
- In 13 studies reporting psychological and indirect burdens, 50% (95% CI, 39%–61%).
- In 1 study reporting indirect and direct burdens, 42% (95% CI, 39%–45%).
- In 2 studies reporting psychological, indirect, and direct burdens, 59% (95% CI, 54%–63%).
- In 1 study reporting indirect burdens, 64% (95% CI, 62%–66%).
- In 1 study reporting psychological and direct burdens, 49% (95% CI, 41%–56%).

Therefore, by type of financial burden, no substantial differences in prevalence were observed (meta-regression $P=.24$).

Risk of Bias

There was a risk of recall bias within studies, particularly for studies with cross-sectional or retrospective cohort design in which a subject's financial toxicity was not determined prospectively but instead depended on recall and self-report of financial toxicity that the subject retrospectively attributed as a consequence of cancer treatment. Therefore, temporality of associations between financial toxicity and outcomes could not be confirmed. Not all financial toxicity tools/measures had reliability and validity testing, which could lead to misclassification bias. However, some financial toxicity measures were objective (eg, out-of-pocket [OOP] costs and copays determined by insurance claims records, or bankruptcy as determined by public records and not subject self-reporting) and/or validation-related (eg, the COST tool), and themes between findings from studies using validated measures did not differ categorically from studies using other financial toxicity measures. Selection bias may have occurred with underrepresentation of nonwhite subjects in studies.

Selection bias may have also occurred if nonparticipants systematically differed from participants (eg, if nonparticipants tended to have more severe disease or financial toxicity causing barriers to participation), though in the 56 evaluable studies the participation rate exceeded 50% in 86% of studies. Of the 6 longitudinal studies, attrition bias because of deaths from metastatic or advanced cancer could have impacted the generalizability of longitudinal findings to patients with more severe disease.

Measures of Financial Toxicity

Regarding existing measures of financial toxicity, no single concept, measure, or tool was consistently used across studies. Measures included validated items or tools (the Medical Expenditure Panel Survey²² and the COST tool²³), objective quantification or estimation of OOP costs, and subjective questions about the severity of treatment, financial burdens, and/or psychological stress (eg, "How much financial burden currently resulted from your illness and treatment?").²⁴

Additional Quantitative Analyses

Funnel plot and regression asymmetry tests to assess small-study bias were not feasible because of the small number of studies for quantitative synthesis. For the association between insurance status and odds of financial toxicity, on sensitivity analysis, greater odds were observed in patients with breast cancer aged >65 years receiving adjuvant hormonal treatments. Insurance was still associated with financial toxicity when removing the studies evaluating breast cancer (odds ratio [OR], 1.77; 95% CI, 1.38–2.25; $P=31\%$) or any other particular type of cancer (OR, 2.17; 95% CI, 1.63–2.89; $P=0\%$). For the association between financial toxicity and odds of medication nonadherence, the association was still significant after removing the study evaluating leukemia only (OR, 2.22; 95% CI, 1.25–3.96; $P=13\%$).

References

1. Kale HP, Carroll NV. Self-reported financial burden of cancer care and its effect on physical and mental health-related quality of life among US cancer survivors. *Cancer* 2016;122:283–289.
2. Lee M, Salloum RG. Racial and ethnic disparities in cost-related medication non-adherence among cancer survivors. *J Cancer Surviv* 2016;10:534–544.
3. Zafar SY, McNeil RB, Thomas CM, et al. Population-based assessment of cancer survivors' financial burden and quality of life: a prospective cohort study. *J Oncol Pract* 2015;11:145–150.
4. Wong YN, Schluchter MD, Albrecht TL, et al. Financial concerns about participation in clinical trials among patients with cancer. *J Clin Oncol* 2016;34:479–487.
5. Ramsey SD, Bansal A, Fedorenko CR, et al. Financial insolvency as a risk factor for early mortality among patients with cancer. *J Clin Oncol* 2016;34:980–986.
6. Delgado-Guay M, Ferrer J, Rieber AG, et al. Financial distress and its associations with physical and emotional symptoms and quality of life among advanced cancer patients. *Oncologist* 2015;20:1092–1098.
7. Peng L, Ye X, Zhou Y, et al. Meta-analysis of incidence and risk of peripheral neuropathy associated with intravenous bortezomib. *Support Care Cancer* 2015;23:2813–2824.
8. Shankaran V, Jolly S, Blough D, et al. Risk factors for financial hardship in patients receiving adjuvant chemotherapy for colon cancer: a population-based exploratory analysis. *J Clin Oncol* 2012;30:1608–1614.
9. Regenbogen SE, Veenstra CM, Hawley ST, et al. The personal financial burden of complications after colorectal cancer surgery. *Cancer* 2014;120:3074–3081.
10. Goodwin JA, Coleman EA, Sullivan E, et al. Personal financial effects of multiple myeloma and its treatment. *Cancer Nurs* 2013;36:301–308.
11. Meeker CR, Wong YN, Egleston BL, et al. Distress and financial distress in adults with cancer: an age-based analysis. *J Natl Compr Canc Netw* 2017;15:1224–1233.
12. Zheng Z, Yabroff KR, Guy GP Jr, et al. Annual medical expenditure and productivity loss among colorectal, female breast, and prostate cancer survivors in the United States. *J Natl Cancer Inst* 2015;108:djv382.
13. de Souza JA, Kung S, O'Connor J, et al. Determinants of patient-centered financial stress in patients with locally advanced head and neck cancer. *J Oncol Pract* 2017;13:e310–318.
14. Banegas MP, Guy GP Jr, de Moor JS, et al. For working-age cancer survivors, medical debt and bankruptcy create financial hardships. *Health Aff (Millwood)* 2016;35:54–61.
15. Martin MY, Fouad MN, Oster RA, et al. What do cancer patients worry about when making decisions about treatment? Variation across racial/ethnic groups. *Support Care Cancer* 2014;22:233–244.
16. Neugut AI, Subar M, Wilde ET, et al. Association between prescription co-payment amount and compliance with adjuvant hormonal therapy in women with early-stage breast cancer. *J Clin Oncol* 2011;29:2534–2542.
17. Fenn KM, Evans SB, McCorkle R, et al. Impact of financial burden of cancer on survivors' quality of life. *J Oncol Pract* 2014;10:332–338.
18. Wheeler SB, Spencer JC, Pinheiro LC, et al. Financial impact of breast cancer in black versus white women. *J Clin Oncol* 2018;36:1695–1701.
19. Farias AJ, Hansen RN, Zeliadt SB, et al. The association between out-of-pocket costs and adherence to adjuvant endocrine therapy among newly diagnosed breast cancer patients. *Am J Clin Oncol* 2018;41:708–715.
20. Shankaran V, Leahy T, Steelquist J, et al. Pilot feasibility study of an oncology financial navigation program. *J Oncol Pract* 2018;14:e122–129.
21. DiMartino LD, Birken SA, Mayer DK. The relationship between cancer survivors' socioeconomic status and reports of follow-up care discussions with providers. *J Cancer Educ* 2017;32:749–755.
22. Peng L, Bu Z, Ye X, et al. Incidence and risk of peripheral neuropathy with nab-paclitaxel in patients with cancer: a meta-analysis. *Eur J Cancer Care [Br]* 2017;26:e12407.
23. Chongpison Y, Hornbrook MC, Harris RB, et al. Self-reported depression and perceived financial burden among long-term rectal cancer survivors. *Psychooncology* 2016;25:1350–1356.
24. Wan Y, Gao X, Mehta S, et al. Indirect costs associated with metastatic breast cancer. *J Med Econ* 2013;16:1169–1178.