NCCN: Continuing Education

Target Audience: This activity is designed to meet the educational needs of oncologists, nurses, pharmacists, and other healthcare professionals who manage patients with cancer.

Accreditation Statements

In support of improving patient care, National Comprehensive Cancer Network (NCCN) is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medicine (ACCME): NCCN designates this journal-based CME activity for a maximum of 1.0 AMA PRA Category 1 Credit™. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Nursing (ANCC): NCCN designates this educational activity for a maximum of 1.0 contact hour.

Pharmacy (ACPE): NCCN designates this knowledge-based continuing education activity for 1.0 contact hour (0.1 CEUs) of continuing education credit. UAN: JA4008196-0000-20-012-H01-P

All clinicians completing this activity will be issued a certificate of participation. To participate in this journal CE activity: (1) review the educational content; (2) take the posttest with a 66% minimum passing score and complete the evaluation at <https://education.nccn.org/node/88590>; and (3) view/print certificate.

Pharmacists: You must complete the posttest and evaluation within 30 days of the activity. Continuing pharmacy education credit is reported to the CPE Monitor once you have completed the posttest and evaluation and claimed your credits. Before completing these requirements, be sure your NCCN profile has been updated with your NAPB e-profile ID and date of birth. Your credit cannot be reported without this information. If you have any questions, please e-mail education@nccn.org.

Release date: November 10, 2020; Expiration date: November 10, 2021

Learning Objectives:

Upon completion of this activity, participants will be able to:

- Integrate into professional practice the updates to the NCCN Guidelines for T-Cell Lymphomas
- Describe the rationale behind the decision-making process for developing the NCCN Guidelines for T-Cell Lymphomas

Disclosure of Relevant Financial Relationships

The NCCN staff listed below discloses no relevant financial relationships:

Kerrin M. Rosenthal, MA; Kimberly Callan, MS; Genevieve Emberger Hartzman, MA; Erin Hesler; Kristina M. Gregory, RN, MSN, OCN; Rashmi Kumar, PhD; Karen Kanefield; and Kathy Smith.

Individuals Who Provided Content Development and/or Authorship Assistance:

Steven M. Horwitz, MD, Panel Chair, has disclosed that he has received consulting fees from Astex Pharmaceuticals, Celgene Corporation, Janssen Pharmaceutica Products, LP, Kura Oncology, Inc., Kyowa Hakko Kirin Co., Ltd., ADCT, C4 Therapeutics, Curio, Myeloid Therapeutics, Verastem, Seattle Genetics, Inc., and Takeda Pharmaceuticals North America, Inc.; and that he has received grant/research support from Celgene Corporation, Daiichi-Sankyo Co., Forty Seven, Inc., ADCT, Aileron, Corvus, Trillium, Verastem, Portola Pharmaceuticals, Inc., and Seattle Genetics, Inc.

Stephen Ansell, MD, PhD, Panel Vice Chair, has disclosed that he has received grant/research support from Bristol-Myers Squibb, Seattle Genetics, Takeda Pharmaceuticals North America, Inc., Al Therapeutics, Inc., Regeneron Pharmaceuticals, Inc., Trillium Therapeutics Inc., Affimed, and ADC Therapeutics.

Stefan K. Barta, MD, MRCP, MS, Panel Member, has disclosed that he has received grant/research support from Bayer Healthcare, Celgene Corporation, Merck & Co., Inc., and Seattle Genetics, Inc.; consulting fees from Janssen Pharmaceutica Products, LP, and Monsanto; and honoraria from Mundipharma.

Ahmet Dogan, MD, PhD, Panel Member, has disclosed that he has served as a scientific advisor for AbbVie, Inc.; received consulting fees from EUSA Pharma, Seattle Genetics, Inc., and Takeda Pharmaceuticals North America, Inc.; received grant/research support from Genentech, Inc. and Takeda Pharmaceuticals North America, Inc.; and received honoraria from Physicians Education Resource.

Aaron M. Goodman, MD, Panel Member, has disclosed that he has received consulting fees from EUSA Pharma and Seattle Genetics, Inc.

Gaurav Goyal, MD, Panel Member, has disclosed that he has no relevant financial relationships.

Ahmad Halwani, MD, Panel Member, has disclosed that he has received grant/research support from AbbVie, Inc., Bayer HealthCare, Genentech, Inc., Immune Design, Pharmacycics, and Roche Laboratories, Inc.

Bradley M. Haverkos, MD, MPH, MS, Panel Member, has disclosed that he has received consulting fees from Viracta Therapeutics.

Deepa Jagadeesh, MD, MPH, Panel Member, has disclosed that she has served as a scientific advisor for Kyowa Hakko Kirin Co., Ltd., Atara Biotherapeutics, Verastem, and Seattle Genetics, Inc.; and has received grant/research support from MEI Pharma Inc., Debiopharm, Regeneron Pharmaceuticals, Inc., and Seattle Genetics, Inc.

Neha Mehta-Shah, MD, Panel Member, has disclosed that she receives consulting fees from Kyowa Hakko Kirin Co., Ltd., C4 Therapeutics, and Karyopharm Therapeutics Inc.; and that she has received grant/research support from Bristol-Myers Squibb, Celgene Corporation, Genentech/Roche, Innate Pharma S.A., Verastem, Inc., and Corvus Pharmaceuticals, Inc.

Barbara Pro, MD, Panel Member, has disclosed that she has received grant/research support from Verastem, Inc., Seattle Genetics, Inc., and Takeda Pharmaceuticals North America, Inc.; honoraria from Seattle Genetics, Inc., and Takeda Pharmaceuticals North America, Inc.; and consulting fees from Seattle Genetics, Inc.

Aaron Shaver, MD, PhD, Panel Member, has disclosed that he has no relevant financial relationships.

Lubomir Sokol, MD, PhD, Panel Member, has disclosed that he has received consulting fees from Kyowa Hakko Kirin Co., Ltd.

Basem M. William, MD, Panel Member, has disclosed that he has received consulting fees from Celgene Corporation, Kyowa Hakko Kirin Co., Ltd., Guidepoint Global, and Techspert; and grant/research support from Incyte Corporation and Dova.

Jasmine Zain, MD, Panel Member, has disclosed that she has received consulting fees from Verastem, Mundi Pharma, and Curio Biotech SA; and she is a speaker and has received consulting fees and grant/research support from Seattle Genetics, Inc.

Mary A. Dwyer, MS, CGC, Director, Guidelines Operations, NCCN, has disclosed that she has no relevant financial relationships.

Hema Sundar, PhD, Oncology Scientist/Senior Medical Writer, NCCN, has disclosed that she has no relevant financial relationships.

To view all of the conflicts of interest for the NCCN Guidelines panel, go to [NCCN.org/disclosures/guidelinepanellisting.aspx](https://www.nccn.org/disclosures/guidelinepanellisting.aspx).

This activity is supported by educational grants from AstraZeneca; Celgene Corporation; Coherus BioSciences; Genentech, a member of the Roche Group; and TESARO, a GSK Company. This activity is supported in part by an educational grant from Bayer Healthcare Pharmaceuticals. This activity is supported by an independent medical education grant from Bristol-Myers Squibb. This activity is supported by a medical education grant from Exelixis, Inc. This activity is supported by an independent educational grant from Merck & Co., Inc.

T-Cell Lymphomas, Version 1.2021

Featured Updates to the NCCN Guidelines

Steven M. Horwitz, MD^{1,*}; Stephen Ansell, MD, PhD^{2,*}; Weiyun Z. Ai, MD, PhD³; Jeffrey Barnes, MD, PhD⁴; Stefan K. Barta, MD, MRCP, MS^{5,*}; Mark W. Clemens, MD⁶; Ahmet Dogan, MD, PhD^{1,*}; Aaron M. Goodman, MD^{7,*}; Gaurav Goyal, MD^{8,*}; Joan Guitart, MD⁹; Ahmad Halwani, MD^{10,*}; Bradley M. Haverkos, MD, MPH, MS^{11,*}; Richard T. Hoppe, MD¹²; Eric Jacobsen, MD¹³; Deepa Jagadeesh, MD, MPH^{14,*}; Allison Jones, MD¹⁵; Youn H. Kim, MD¹²; Neha Mehta-Shah, MD^{16,*}; Elise A. Olsen, MD¹⁷; Barbara Pro, MD^{9,*}; Saurabh A. Rajguru, MD¹⁸; Sima Rozati, MD, PhD¹⁹; Jonathan Said, MD²⁰; Aaron Shaver, MD, PhD^{21,*}; Andrei Shustov, MD²²; Lubomir Sokol, MD, PhD^{23,*}; Pallawi Torka, MD²⁴; Carlos Torres-Cabala, MD⁶; Ryan Wilcox, MD, PhD²⁵; Basem M. William, MD^{26,*}; Jasmine Zain, MD^{27,*}; Mary A. Dwyer, MS, CGC^{28,*}; and Hema Sundar, PhD^{28,*}

ABSTRACT

Hepatosplenic T-cell lymphoma (HSTCL) is a rare subtype of T-cell lymphoma associated with an aggressive clinical course and a worse prognosis. HSTCL develops in the setting of chronic immune suppression or immune dysregulation in up to 20% of cases and is most often characterized by spleen, liver, and bone marrow involvement. Diagnosis and management of HSTCL pose significant challenges given the rarity of the disease along with the absence of lymphadenopathy and poor outcome with conventional chemotherapy regimens. These Guidelines Insights focus on the diagnosis and treatment of HSTCL as outlined in the NCCN Guidelines for T-Cell Lymphomas.

J Natl Compr Canc Netw 2020;18(11):1460–1467
doi: 10.6004/jnccn.2020.0053

¹Memorial Sloan Kettering Cancer Center; ²Mayo Clinic Cancer Center; ³UCSF Helen Diller Family Comprehensive Cancer Center; ⁴Massachusetts General Hospital Cancer Center; ⁵Abramson Cancer Center at the University of Pennsylvania; ⁶The University of Texas MD Anderson Cancer Center; ⁷UC San Diego Moores Cancer Center; ⁸O'Neal Comprehensive Cancer Center at UAB; ⁹Robert H. Lurie Comprehensive Cancer Center of Northwestern University; ¹⁰Huntsman Cancer Institute at the University of Utah; ¹¹University of Colorado Cancer Center; ¹²Stanford Cancer Institute; ¹³Dana-Farber/Brigham and Women's Cancer Center; ¹⁴Case Comprehensive Cancer Center/University Hospitals Seidman Cancer Center and Cleveland Clinic Taussig Cancer Institute; ¹⁵St. Jude Children's Research Hospital/The University of Tennessee Health Science Center; ¹⁶Siteman Cancer Center at Barnes-Jewish Hospital and Washington University School of Medicine; ¹⁷Duke Cancer Institute; ¹⁸University of Wisconsin Carbone Cancer Center; ¹⁹The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins; ²⁰UCLA Jonsson Comprehensive Cancer Center; ²¹Vanderbilt-Ingram Cancer Center; ²²Fred Hutchinson Cancer Research Center/Seattle Cancer Care Alliance; ²³Moffitt Cancer Center; ²⁴Roswell Park Comprehensive Cancer Center; ²⁵University of Michigan Rogel Cancer Center; ²⁶The Ohio State University Comprehensive Cancer Center - James Cancer Hospital and Solove Research Institute; ²⁷City of Hope National Medical Center; and ²⁸National Comprehensive Cancer Network.

*Provided content development and/or authorship assistance.

See JNCCN.org for supplemental online content.

NCCN CATEGORIES OF EVIDENCE AND CONSENSUS

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Clinical trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

PLEASE NOTE

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. **The NCCN Guidelines Insights highlight important changes in the NCCN Guidelines recommendations from previous versions. Colored markings in the algorithm show changes and the discussion aims to further the understanding of these changes by summarizing salient portions of the panel's discussion, including the literature reviewed.**

The NCCN Guidelines Insights do not represent the full NCCN Guidelines; further, the National Comprehensive Cancer Network[®] (NCCN[®]) makes no representations or warranties of any kind regarding their content, use, or application of the NCCN Guidelines and NCCN Guidelines Insights and disclaims any responsibility for their application or use in any way.

The complete and most recent version of these NCCN Guidelines is available free of charge at NCCN.org.

© National Comprehensive Cancer Network, Inc. 2020. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN.

DIAGNOSIS^{a,b}**ESSENTIAL:**

- Review of all slides with at least one paraffin block representative of the tumor should be done by a hematopathologist with expertise in the diagnosis of T-cell lymphomas. Rebiopsy if consult material is nondiagnostic.
- A core biopsy of bone marrow or liver is required for diagnosis.^c Bone marrow aspirate, FNA of liver, or peripheral blood studies may be helpful but are not alone sufficient for diagnosis.
- Adequate immunophenotyping to establish diagnosis^{d,e}
 - IHC panel may include CD20, CD3, CD10, Ki-67, CD5, CD30, CD2, CD4, CD8, CD7, CD56, EBER-ISH, TCRβ, TCRδ, TIA-1, or granzyme B
 - Cell surface marker analysis by flow cytometry may include kappa/lambda, CD45, CD3, CD5, CD19, CD10, CD20, CD30, CD4, CD8, CD7, CD2, TCRαβ, or TCRγδ

See Workup
(HSTCL-2)

USEFUL UNDER CERTAIN CIRCUMSTANCES:

- Molecular analysis to detect^d clonal *TCR* gene rearrangements or other assessment of clonality.^f
- Karyotype to establish clonality and investigate the presence of isochromosome 7q and trisomy 8.
- FISH analysis for isochromosome 7q and trisomy 8.
- Genomic analysis for *STAT3*, *STAT5B*, *PIK3CD*, *SETD2*, *INO80*, *TET3*, and *SMARCA2*.

^a It is preferred that treatment occur at centers with expertise in the management of this disease.

^b See Principles of Molecular Analysis in T-Cell Lymphomas (TCLYM-A).

^c If the results are equivocal, core biopsy of spleen or splenectomy could be considered in centers with expertise.

^d See Use of Immunophenotyping/Genetic Testing in Differential Diagnosis of Mature B-Cell and NK/T-Cell Neoplasms (See NCCN Guidelines for B-Cell Lymphomas).

^e Typical immunophenotype: CD3+, generally TCRδ+ and TCRβ- (GM3 positive, βF-1 negative), CD4 -, CD8+/-, CD56 +/-, CD5-.

^f Clonal *TCR* gene rearrangements alone are not sufficient for diagnosis, as these can also be seen in patients with non-malignant conditions. Results should be interpreted in the context of overall presentation. See Principles of Molecular Analysis in T-Cell Lymphomas (TCLYM-A).

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

HSTCL-1

Hepatosplenic T-Cell Lymphoma

Hepatosplenic T-cell lymphoma (HSTCL) is a rare lymphoproliferative disorder associated with an aggressive clinical course and a worse prognosis.^{1,2} HSTCL accounts for ≤2% of all cases of T-cell lymphoma diagnosed worldwide, and in up to 20% of cases develops in the setting of chronic immune suppression or immune dysregulation particularly inflammatory bowel disease (IBD), hematologic malignancies, and previous solid organ transplant.^{3,4} The concomitant use of tumor necrosis factor-α (TNF-α) inhibitors and thiopurine-based immunomodulators has been identified as a risk factor for developing HSTCL among patients with IBD.^{5,6}

HSTCL is most often characterized by spleen, liver, and bone marrow involvement. Lymphadenopathy is uncommon and patients frequently present with systemic symptoms, hepatosplenomegaly, cytopenias, and sometimes hemophagocytic lymphohistiocytosis (HLH).^{7,8} Clinical presentation is highly nonspecific and a high index of suspicion is required to make the diagnosis. In most cases, the neoplastic cells typically arise from lymphocytes having the surface expression of T-cell receptor δ (TCRδ) and TCRγδ.^{9,10} In rare cases, neoplastic cells may express TCRαβ.^{11–13} TCRγδ variant has a male predominance with a median age of 35 years, whereas TCRαβ variant occurs more commonly in

women aged >50 years.⁴ Both are considered immunophenotypic variants of the same disease and are managed the same way. These Guidelines Insights focus on the diagnosis and treatment of HSTCL as outlined in the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for T-Cell Lymphomas.

Diagnosis

Diagnosis of HSTCL is most frequently established by a core needle biopsy of bone marrow or liver with adequate immunophenotyping (either by immunohistochemistry or cell surface marker analysis by flow cytometry) as well as through molecular studies (see HSTCL-1, above).⁴ Splenectomy may be required in equivocal cases, and core needle biopsy of spleen could be considered in some cases at centers of excellence with expertise performing this procedure. It is common for several biopsies to be needed prior to making a definitive diagnosis, because biopsy results may be inconclusive. Examination of peripheral blood smear, bone marrow aspirate, and fine needle aspiration of liver may be helpful but are not solely sufficient for the diagnosis.

Interpretation of cytotoxic cells seen on the bone marrow biopsy specimen may be difficult, and additional liver biopsy may be helpful to confirm the diagnosis. Liver biopsy with adequate immunophenotyping should

WORKUP

ESSENTIAL:

- H&P exam; full skin exam; attention to node-bearing areas, including Waldeyer's ring; evaluation of size of liver and spleen, nasopharynx
- Performance status
- B symptoms
- CBC with differential
- Bone marrow biopsy ± aspirate
- LDH
- Comprehensive metabolic panel
- HLH workup (See TCLYM-B 2 of 3)
- Uric acid
- PET/CT scan⁹ and/or C/A/P CT with contrast of diagnostic quality
- Echocardiogram or MUGA scan if anthracycline-based regimen is indicated
- Pregnancy testing in women of childbearing age (if chemotherapy or RT planned)
- HLA typing

USEFUL IN SELECTED CASES:

- Neck CT with contrast
- Head CT or MRI with contrast
- HIV testing
- Hepatitis B and C testing
- Consider quantitative EBV PCR
- Discussion of fertility issues and sperm banking
- Assessment of HTLV-1/2^h by serology or other methods as clinically indicated

See First-Line Therapy
(HSTCL-3)

⁹ Patients with T-cell lymphomas often have extranodal disease, which may be inadequately imaged by CT. PET scan may be preferred in these instances.

^h See map for prevalence of HTLV-1/2 by geographic region. HTLV-1/2 has been described in patients in non-endemic areas.

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

HSTCL-2

be reviewed by a hematopathologist.¹⁴ HSTCL is typically characterized by the following immunophenotypes: CD2+, CD3+, CD4-, CD5-, CD8±, CD56±, TCRγδ+, TIA1+, TdT, and granzyme B-.¹⁵ An immunohistochemistry panel to evaluate for HSTCL typically includes CD20, CD3, CD10, Ki-67, CD5, CD30, CD2, CD4, CD8, CD7, CD56, EBER-ISH, TCRβ, TCRδ, TIA-1, and granzyme B. Cell surface marker analysis by flow cytometry often includes kappa/lambda, CD45, CD3, CD5, CD19, CD10, CD20, CD30, CD4, CD8, CD7, CD2, TCRδ, TCRαβ, and TCRγδ.¹⁴

Molecular analysis or other assessment of clonality can be used to detect clonal *TCR* gene rearrangements (see TCLYM-A, page 1466). Identification of *TCR* gene rearrangement on molecular analysis reflects the clonality of the T cells. However, the molecular clonality studies cannot be used to define the T-cell subtype (αβ vs γδ) because *TCR*β and *TCR*γ gene rearrangements may be seen in both αβ and γδ HSTCL.¹²

Isochromosome 7q and trisomy 8 are the most common chromosomal abnormalities in HSTCL.^{16–19} Isochromosome 7q and ring chromosome 7 are associated with loss of 7p and amplification of 7q, resulting in altered expressions of several oncogenes located on chromosome 7 (*CHN2*, *ABCB1*, and *PPP1R9A*).²⁰ Gene expression profiling studies have identified distinct

molecular signatures that distinguish HSTCL from other T-cell lymphomas.^{21–23} In a whole-exome sequencing study on 68 primary HSTCL tumors, mutations in chromatin-modifying genes, including *SETD2*, *INO80*, and *ARID1B* (occurring almost exclusively in HSTCL compared with other T-cell lymphoma subtypes), were present in 62% of cases.²³ In addition, *STAT5B*, *STAT3*, and *PIK3CD* mutations have also been identified in 31%, 9%, and 9% of cases, respectively.²³ *STAT3* and *STAT5* mutations, however, are not unique to HSTCL and have also been identified in large granular lymphocytic leukemia (LGLL) and other T-cell lymphoma subtypes.^{24–27}

It is essential to consider other natural killer (NK)/T-cell neoplasms with significant overlapping features with HSTCL in the differential diagnosis (eg, γδ T-cell LGLL, T-lymphoblastic leukemia, primary cutaneous γδ T-cell lymphoma, intestinal monomorphic epitheliotropic intestinal T-cell lymphoma, aggressive NK-cell leukemia, Epstein-Barr virus [EBV]-positive T-cell and NK-cell lymphoproliferative diseases of childhood, and, rarely, other T-cell lymphomas with expression of TCRγδ).^{14,28} Fluorescence in situ hybridization and/or karyotyping to identify isochromosome 7q and trisomy 8 and mutation analysis to identify *SETD2*, *INO80*, and *TET3* mutations would be useful for the differential diagnosis for HSTCL (see HSTCL-1, page 1462).^{19,23} Nonneoplastic, transient

conditions leading to an increase in $\gamma\delta$ T-cells with a similar phenotype, including infections such as ehrlichiosis and other tick-borne diseases, should also be considered in the differential diagnosis of HSTCL.^{29,30}

Workup

Initial workup should include a comprehensive medical history and physical examination, including a full skin evaluation and routine laboratory studies (bone marrow biopsy $\pm$ aspirate, CBC with differential, comprehensive metabolic panel, and assessment of serum uric acid and lactate dehydrogenase levels) (see HSTCL-2, page 1463). FDG-PET/CT and/or CT of chest/abdomen/pelvis with contrast of diagnostic quality are essential for workup. In the absence of lymphadenopathy, normal FDG uptake in lymph nodes is a pertinent negative finding on PET/CT scan that could differentiate HSTCL from other lymphomas.³¹ CT scan of the neck and CT or MRI of the head may be useful in some cases. Multigated acquisition (MUGA) scan or echocardiogram is recommended under certain circumstances. Quantitative PCR for EBV and cytomegalovirus reactivation as well as serology testing for HIV and human T-cell lymphotropic virus (HTLV-1) may be useful in selected cases. Human leukocyte antigen (HLA) typing is recommended for all patients eligible for transplant, because HSTCL is associated with a poor outcome in the absence of a consolidative allogeneic hematopoietic

cell transplant (HCT). Early referral to transplant is advisable for planning purposes.¹³

Hemophagocytic lymphohistiocytosis (HLH) is a rare but potentially life-threatening hyperinflammatory syndrome. HLH in adults is most often associated with an underlying hematologic malignancy, especially T-cell lymphomas.⁷ HSTCL should be considered in the differential diagnosis when evaluating patients presenting with symptoms associated with HLH. Diagnostic workup to confirm the lymphoma subtype and prompt initiation of treatment of underlying T-cell lymphoma is often required (see TCLYM-B, 2 of 3, available with this article at JNCCN.org).

Treatment

HSTCL has been underrepresented in prospective clinical studies, and treatment recommendations are based on the evidence mainly from small case reports or case series and single-center retrospective studies.⁴ Outcomes are poor with CHOP (cyclophosphamide/doxorubicin/vincristine/prednisone)-based chemotherapy regimens. More intensive non-CHOP-based chemotherapy regimens, such as ICE (ifosfamide/carboplatin/etoposide) or IVAC (ifosfamide/etoposide/cytarabine) have been associated with potentially improved outcomes.^{32–35} Purine analogs (pentostatin or cladribine) either as monotherapy or in combination with alemtuzumab have also demonstrated modest activity.^{36–41}

SUGGESTED TREATMENT REGIMENS

FIRST-LINE THERAPY/ADDITIONAL THERAPY ^{a,b}
<ul style="list-style-type: none"> • Clinical trial (preferred) Preferred regimen • ICE
Other recommended regimens (alphabetical order) <ul style="list-style-type: none"> • Alemtuzumab^c + pentostatin • CHOEP • Dose-adjusted EPOCH • HyperCVAD alternating with high-dose methotrexate and cytarabine • IVAC (ifosfamide, etoposide, and cytarabine)
Useful in certain circumstances <ul style="list-style-type: none"> • Brentuximab vedotin + CHP (cyclophosphamide, doxorubicin, and prednisone) for CD30+ cases^d <ul style="list-style-type: none"> ▶ DHAP (dexamethasone, cytarabine, cisplatin) ▶ DHAX (dexamethasone, cytarabine, oxaliplatin)

^a CHOP is not adequate therapy.

^b Voss MH, Lunning MA, Maragulia JC, et al. Intensive induction chemotherapy followed by early high-dose therapy and hematopoietic stem cell transplantation results in improved outcome for patients with hepatosplenic T-cell lymphoma: a single institution experience. Clin Lymphoma Myeloma Leuk 2013;13:8-14; Klebaner D, Koura D, Tzachanis D, et al. Intensive induction therapy compared with CHOP for Hepatosplenic T-cell Lymphoma. Clin Lymphoma Myeloma Leuk 2020;20:431-437 e432.

^c While alemtuzumab is no longer commercially available, it may be obtained for clinical use. Recommend CMV prophylaxis (See TCLYM-B).

^d Patients with HSTCL were eligible for the ECHELON-2 study (Horwitz S, et al. Lancet 2019;393:229-240), but no patients were enrolled. Also see Supportive Care (TCLYM-B).

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

HSTCL-A

Few studies have reported improved survival outcomes with autologous or allogeneic HCT as consolidation therapy for patients with disease in first or second remission.^{34,42,43} Some studies have also reported that graft-versus-lymphoma effect conferred by allogeneic HCT may result in long-term survival in a significant proportion of patients with HSTCL, and that active disease at the time of transplant was not necessarily associated with poor outcomes.^{42,43} The efficacy of allogeneic HCT in relapsed or refractory disease has also been demonstrated in several case reports.⁴⁴⁻⁴⁶

A more recent individual-level meta-analysis (which represents the largest aggregation of all published studies and case reports to date: 166 patients with a diagnosis of HSTCL) compared the response rates and overall survival outcomes of 84 patients with HSTCL treated with CHOP or CHOP-like regimens (n=50) or non-CHOP-based regimens, specifically those containing cytarabine, platinum, and etoposide (n=34).³⁵ Non-CHOP-based regimens were associated with an overall response rate of 82% compared with 52% for CHOP or CHOP-like regimens ($P=.006$), and median survival was 37 and 18 months, respectively ($P=.00014$). Use of non-CHOP-based regimens was a significant predictor of higher response rate ($P=.049$) and improved survival ($P=.026$).

This study also demonstrated a benefit for HCT on survival and the superiority of allogeneic HCT over

autologous HCT. The 2-year survival rate was 12% for patients who did not receive HCT compared with 41% and 56%, respectively, for those who received autologous HCT and allogeneic HCT. In the absence of data from prospective and randomized studies, the results of this largest meta-analysis support the use of induction therapy with non-CHOP-based regimens followed by consolidation with allogeneic HCT as an effective treatment approach (associated with improved survival) for all eligible patients with HSTCL. Autologous HCT has also been shown to provide some benefit for patients when an allogeneic HCT is not feasible.³⁴ In a retrospective series of 14 patients with HSTCL, induction therapy with ICE or IVAC followed by consolidation with autologous HCT was associated with improved outcomes compared with use of a CHOP or CHOP-like regimen.³⁴

NCCN Recommendations

The optimal treatment approach for HSTCL remains undefined given the absence of data from prospective randomized clinical studies. Enrollment in a clinical trial, if an appropriate one is available, is the preferred initial treatment option for all patients with HSTCL. The goal of initial therapy is to induce complete or near-complete response to allow successful bridging to HCT, preferably an allogeneic HCT. Because HSTCL is non-nodal, Lugano response criteria do not apply for response assessment,

PRINCIPLES OF MOLECULAR ANALYSIS IN T-CELL LYMPHOMAS^a

- Genetic testing, including high-throughput sequencing (HTS), array-based comparative genomic hybridization (CGH), NGS, karyotype, or FISH to detect somatic mutations or genetic abnormalities are often informative and in some cases essential for an accurate and precise diagnostic and prognostic assessment of T-cell lymphomas.

T-Cell Antigen Receptor (TCR) Gene Rearrangements

- TCR gene rearrangement testing is recommended to support a diagnosis of T-cell lymphoma.

Diseases:

- PTCLs; mycosis fungoides/Sézary syndrome; primary cutaneous CD30+ T-cell LPD; T-LGL leukemia; T-cell prolymphocytic leukemia (T-PLL); extranodal NK/T-cell lymphoma, nasal type; and hepatosplenic T-cell lymphoma.

Description:

- TCR gene rearrangement is indicative of T-cell clonal expansion. The test targets the gamma and/or beta TCR genes using PCR methods with capillary or gel electrophoresis detection methods. Alternatively, HTS methods are increasingly utilized. HTS methods are more sensitive, precise, and capable of providing a unique sequence of the T-cell clone, which allows for comparison and confirmation of disease evolution and monitoring during remission. Clonal T-cell expansions can also be detected using V beta families in blood or tissue with flow cytometry methods.

Diagnostic value:

- Clonal TCR gene rearrangements without histopathologic and immunophenotypic evidence of abnormal T-cell population does not constitute a diagnosis of T-cell lymphoma since it can be identified in patients with non-malignant conditions. Conversely, a negative result does not exclude the diagnosis of T-cell lymphoma, which occasionally may fail TCR amplification. Nonetheless, it often provides essential information and increased precision for many of these complex diagnoses.

Prognostic value:

- Identification of clonal TCR gene rearrangement has no definitive established prognostic value; however, it could be helpful when used to determine clinical staging or assess relapsed or residual disease.

Continued

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

TCLYM-A
1 OF 4

and PET-negative response should be confirmed by bone marrow biopsy and, in selected cases, liver biopsy.

CHOP is not considered adequate therapy.³⁵ The NCCN Guidelines for T-Cell Lymphomas have included ICE as the preferred regimen for induction therapy because it is used in most NCCN Member Institutions (see HSTCL-A, page 1465). Other intensive induction therapy regimens, such as IVAC and hyper-CVAD (cyclophosphamide/vincristine/doxorubicin/dexamethasone) alternating with high-dose methotrexate and cytarabine, may also be appropriate.^{34,35} Alemtuzumab + pentostatin, CHOEP (cyclophosphamide/doxorubicin/vincristine/etoposide/prednisone), dose-adjusted EPOCH (etoposide/prednisone/vincristine/cyclophosphamide/doxorubicin), and DHAP or DHAX (dexamethasone/cytarabine) + cisplatin or oxaliplatin are included as alternative options.

Consolidation therapy with allogeneic HCT is recommended for eligible patients experiencing complete or partial response after initial induction or second-line

therapy (see HSTCL-3, page 1464).^{35,40,43} Consolidation therapy with autologous HCT can be considered if a suitable donor is not available or in patients who are ineligible for allogeneic HCT.³⁴

Patients with disease not responding to primary treatment or those with progressive disease should be treated with alternate induction therapy regimens before receiving treatment for relapsed/refractory disease.⁴ Purine analogs or regimens recommended for second-line therapy for primary T-cell lymphoma, not otherwise specified may be appropriate for the management of patients with relapsed/refractory HSTCL. Responses have been observed with alemtuzumab, pralatrexate, and ESHAP (etoposide/methylprednisone/cytarabine/cisplatin).⁴

To participate in this journal CE activity, go to
<https://education.nccn.org/node/88590>

References

- Swerdlow SH, Harris NL, Jaffe ES, et al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, Revised 4th ed. Lyon, France: IARC; 2017.
- Foss FM, Horwitz SM, Civallero M, et al. Incidence and outcomes of rare T cell lymphomas from the T Cell Project: hepatosplenic, enteropathy associated and peripheral gamma delta T cell lymphomas. *Am J Hematol* 2020;95:151–155.
- Gratzinger D, de Jong D, Jaffe ES, et al. T- and NK-cell lymphomas and systemic lymphoproliferative disorders and the immunodeficiency setting: 2015 SH/EAHP workshop report—part 4. *Am J Clin Pathol* 2017;147:188–203.
- Pro B, Allen PB, Behdad A. Hepatosplenic T-cell lymphoma: a rare but challenging entity. *Blood* 2020;blood.2019004118.

5. Parakkal D, Sifuentes H, Semer R, et al. Hepatosplenic T-cell lymphoma in patients receiving TNF- α inhibitor therapy: expanding the groups at risk. *Eur J Gastroenterol Hepatol* 2011;23:1150–1156.
6. Yabe M, Medeiros LJ, Daneshbod Y, et al. Hepatosplenic T-cell lymphoma arising in patients with immunodysregulatory disorders: a study of 7 patients who did not receive tumor necrosis factor- α inhibitor therapy and literature review. *Ann Diagn Pathol* 2017;26:16–22.
7. Daver N, McClain K, Allen CE, et al. A consensus review on malignancy-associated hemophagocytic lymphohistiocytosis in adults. *Cancer* 2017;123:3229–3240.
8. Yabe M, Miranda RN, Medeiros LJ. Hepatosplenic T-cell lymphoma: a review of clinicopathologic features, pathogenesis, and prognostic factors. *Hum Pathol* 2018;74:5–16.
9. Farcet JP, Gaulard P, Marolleau JP, et al. Hepatosplenic T-cell lymphoma: sinusoidal/sinusoidal localization of malignant cells expressing the T-cell receptor gamma delta. *Blood* 1990;75:2213–2219.
10. Weidmann E. Hepatosplenic T cell lymphoma. A review on 45 cases since the first report describing the disease as a distinct lymphoma entity in 1990. *Leukemia* 2000;14:991–997.
11. Suarez F, Wlodarska I, Rigal-Huguet F, et al. Hepatosplenic alphabeta T-cell lymphoma: an unusual case with clinical, histologic, and cytogenetic features of gammadelta hepatosplenic T-cell lymphoma. *Am J Surg Pathol* 2000;24:1027–1032.
12. Macon WR, Levy NB, Kurtin PJ, et al. Hepatosplenic alphabeta T-cell lymphomas: a report of 14 cases and comparison with hepatosplenic gammadelta T-cell lymphomas. *Am J Surg Pathol* 2001;25:285–296.
13. Yabe M, Medeiros LJ, Tang G, et al. Prognostic factors of hepatosplenic T-cell lymphoma: clinicopathologic study of 28 cases. *Am J Surg Pathol* 2016;40:676–688.
14. Shi Y, Wang E. Hepatosplenic T-cell lymphoma: a clinicopathologic review with an emphasis on diagnostic differentiation from other T-cell/natural killer-cell neoplasms. *Arch Pathol Lab Med* 2015;139:1173–1180.
15. Cooke CB, Krenacs L, Stetler-Stevenson M, et al. Hepatosplenic T-cell lymphoma: a distinct clinicopathologic entity of cytotoxic gamma delta T-cell origin. *Blood* 1996;88:4265–4274.
16. Wang CC, Tien HF, Lin MT, et al. Consistent presence of isochromosome 7q in hepatosplenic T gamma/delta lymphoma: a new cytogenetic-clinicopathologic entity. *Genes Chromosomes Cancer* 1995;12:161–164.
17. Yao M, Tien HF, Lin MT, et al. Clinical and hematological characteristics of hepatosplenic T gamma/delta lymphoma with isochromosome for long arm of chromosome 7. *Leuk Lymphoma* 1996;22:495–500.
18. Alonzoza EL, Stamberg J, Kumar D, et al. Isochromosome 7q: the primary cytogenetic abnormality in hepatosplenic gammadelta T cell lymphoma. *Leukemia* 1997;11:1367–1372.
19. Wlodarska I, Martin-Garcia N, Achten R, et al. Fluorescence in situ hybridization study of chromosome 7 aberrations in hepatosplenic T-cell lymphoma: isochromosome 7q as a common abnormality accumulating in forms with features of cytologic progression. *Genes Chromosomes Cancer* 2002;33:243–251.
20. Finalet Ferreira J, Rouhigharabaei L, Urbankova H, et al. Integrative genomic and transcriptomic analysis identified candidate genes implicated in the pathogenesis of hepatosplenic T-cell lymphoma. *PLoS One* 2014;9:e102977.
21. Travert M, Huang Y, de Leval L, et al. Molecular features of hepatosplenic T-cell lymphoma unravels potential novel therapeutic targets. *Blood* 2012;119:5795–5806.
22. Nicolae A, Xi L, Pittaluga S, et al. Frequent STAT5B mutations in $\gamma\delta$ hepatosplenic T-cell lymphomas. *Leukemia* 2014;28:2244–2248.
23. McKinney M, Moffitt AB, Gaulard P, et al. The genetic basis of hepatosplenic T-cell lymphoma. *Cancer Discov* 2017;7:369–379.
24. Jerez A, Clemente MJ, Makishima H, et al. STAT3 mutations unify the pathogenesis of chronic lymphoproliferative disorders of NK cells and T-cell large granular lymphocyte leukemia. *Blood* 2012;120:3048–3057.
25. Koskela HL, Eldfors S, Ellonen P, et al. Somatic STAT3 mutations in large granular lymphocytic leukemia. *N Engl J Med* 2012;366:1905–1913.
26. Ohgami RS, Ma L, Merker JD, et al. STAT3 mutations are frequent in CD30+ T-cell lymphomas and T-cell large granular lymphocytic leukemia. *Leukemia* 2013;27:2244–2247.
27. Blombery P, Thompson ER, Jones K, et al. Whole exome sequencing reveals activating JAK1 and STAT3 mutations in breast implant-associated anaplastic large cell lymphoma anaplastic large cell lymphoma. *Haematologica* 2016;101:e387–390.
28. Yabe M, Medeiros LJ, Wang SA, et al. Distinguishing between hepatosplenic T-cell lymphoma and $\gamma\delta$ T-cell large granular lymphocytic leukemia: a clinicopathologic, immunophenotypic, and molecular analysis. *Am J Surg Pathol* 2017;41:82–93.
29. Malani A, Weigand R, Gupta V, et al. Ehrlichiosis mimicking T-cell lymphoma/leukemia [abstract] *Blood* 2005;106:Abstract 4343.
30. Marko D, Perry AM, Ponnampalam A, et al. Cytopenias and clonal expansion of gamma/delta T-cells in a patient with anaplasmosis: a potential diagnostic pitfall. *J Clin Exp Hematop* 2017;56:160–164.
31. Cho MW, Chin BB. 18 F-FDG PET/CT findings in hepatosplenic gamma-delta T-cell lymphoma: case reports and review of the literature. *Am J Nucl Med Mol Imaging* 2018;8:137–142.
32. Belhadj K, Reyes F, Farcet JP, et al. Hepatosplenic gammadelta T-cell lymphoma is a rare clinicopathologic entity with poor outcome: report on a series of 21 patients. *Blood* 2003;102:4261–4269.
33. Falchook GS, Vega F, Dang NH, et al. Hepatosplenic gamma-delta T-cell lymphoma: clinicopathological features and treatment. *Ann Oncol* 2009;20:1080–1085.
34. Voss MH, Lunning MA, Maragulia JC, et al. Intensive induction chemotherapy followed by early high-dose therapy and hematopoietic stem cell transplantation results in improved outcome for patients with hepatosplenic T-cell lymphoma: a single institution experience. *Clin Lymphoma Myeloma Leuk* 2013;13:8–14.
35. Klebaner D, Koura D, Tzachanis D, et al. Intensive induction therapy compared with CHOP for hepatosplenic T-cell lymphoma. *Clin Lymphoma Myeloma Leuk* 2020;20:431–437.e2.
36. Gopcsa L, Bányai A, Tamáska J, et al. Hepatosplenic gamma delta T-cell lymphoma with leukemic phase successfully treated with 2-chlorodeoxyadenosine. *Haematologia (Budap)* 2002;32:519–527.
37. Iannitto E, Barbera V, Quintini G, et al. Hepatosplenic gammadelta T-cell lymphoma: complete response induced by treatment with pentostatin. *Br J Haematol* 2002;117:995–996.
38. Corazzelli G, Capobianco G, Russo F, et al. Pentostatin (2'-deoxycoformycin) for the treatment of hepatosplenic gammadelta T-cell lymphomas. *Haematologica* 2005;90:ECR14.
39. Jaeger G, Bauer F, Brezinschek R, et al. Hepatosplenic gammadelta T-cell lymphoma successfully treated with a combination of alemtuzumab and cladribine. *Ann Oncol* 2008;19:1025–1026.
40. Ravandi F, Aribi A, O'Brien S, et al. Phase II study of alemtuzumab in combination with pentostatin in patients with T-cell neoplasms. *J Clin Oncol* 2009;27:5425–5430.
41. Bennett M, Matutes E, Gaulard P. Hepatosplenic T cell lymphoma responsive to 2'-deoxycoformycin therapy. *Am J Hematol* 2010;85:727–729.
42. Rashidi A, Cashen AF. Outcomes of allogeneic stem cell transplantation in hepatosplenic T-cell lymphoma. *Blood Cancer J* 2015;5:e318.
43. Tanase A, Schmitz N, Stein H, et al. Allogeneic and autologous stem cell transplantation for hepatosplenic T-cell lymphoma: a retrospective study of the EBMT Lymphoma Working Party. *Leukemia* 2015;29:686–688.
44. Domm JA, Thompson M, Kuttesch JF, et al. Allogeneic bone marrow transplantation for chemotherapy-refractory hepatosplenic gammadelta T-cell lymphoma: case report and review of the literature. *J Pediatr Hematol Oncol* 2005;27:607–610.
45. Sumi M, Takeda W, Kaiume H, et al. Successful treatment with reduced-intensity cord blood transplant in a patient with relapsed refractory hepatosplenic T-cell lymphoma. *Leuk Lymphoma* 2015;56:1140–1142.
46. Pan H, Huang J, Li JN, et al. Successful second allogeneic stem-cell transplantation from the same sibling donor for a patient with recurrent hepatosplenic gamma-delta ($\gamma\delta$) T-cell lymphoma: a case report. *Medicine (Baltimore)* 2018;97:e12941.

See [JNCCN.org](https://www.nccn.org) for supplemental online content.

Supplemental online content for:

NCCN Guidelines Insights: T-Cell Lymphomas, Version 1.2021

Steven M. Horwitz, MD; Stephen Ansell, MD, PhD; Weiyun Z. Ai, MD, PhD; Jeffrey Barnes, MD, PhD; Stefan K. Barta, MD, MRCP, MS; Mark W. Clemens, MD; Ahmet Dogan, MD, PhD; Aaron M. Goodman, MD; Gaurav Goyal, MD; Joan Guitart, MD; Ahmad Halwani, MD; Bradley M. Haverkos, MD, MPH, MS; Richard T. Hoppe, MD; Eric Jacobsen, MD; Deepa Jagadeesh, MD, MPH; Allison Jones, MD; Youn H. Kim, MD; Neha Mehta-Shah, MD; Elise A. Olsen, MD; Barbara Pro, MD; Saurabh A. Rajguru, MD; Sima Rozati, MD, PhD; Jonathan Said, MD; Aaron Shaver, MD, PhD; Andrei Shustov, MD; Lubomir Sokol, MD, PhD; Pallawi Torka, MD; Carlos Torres-Cabala, MD; Ryan Wilcox, MD, PhD; Basem M. William, MD; Jasmine Zain, MD; Mary A. Dwyer, MS, CGC; and Hema Sundar, PhD

J Natl Compr Canc Netw 2020;18(11):1460–1467

Page 1: TCLYM-B 1 of 3

Page 2: TCLYM-B 2 of 3

Page 3: TCLYM-B 3 of 3

SUPPORTIVE CARE

Tumor Lysis Syndrome (TLS)

- Laboratory hallmarks of TLS:
 - High potassium
 - High uric acid
 - High phosphorous
 - Low calcium
 - Elevated creatinine
- Symptoms of TLS:
 - Nausea and vomiting, shortness of breath, irregular heartbeat, clouding of urine, lethargy, and/or joint discomfort.
- TLS features:
 - Consider TLS prophylaxis for patients with the following risk factors:
 - ◊ Spontaneous TLS
 - ◊ High tumor burden or bulky disease
 - ◊ Elevated WBC count
 - ◊ Bone marrow involvement
 - ◊ Pre-existing elevated uric acid
 - ◊ Renal disease or renal involvement by tumor
- Treatment of TLS:
 - TLS is best managed if anticipated and treatment is started prior to chemotherapy.
 - Centerpiece of treatment includes:
 - ◊ Rigorous hydration
 - ◊ Management of hyperuricemia
 - ◊ Frequent monitoring of electrolytes and aggressive correction (essential)
 - First-line and at retreatment for hyperuricemia
 - ◊ Allopurinol or febuxostat beginning 2–3 days prior to chemotherapy and continued for 10–14 days or
 - Rasburicase (Doses of 3 to 6 mg are usually effective.^a One dose of rasburicase is frequently adequate. Re-dosing should be individualized.) is indicated for patients with any of the following risk factors:
 - Urgent need to initiate therapy in a high-bulk patient
 - Situations where adequate hydration may be difficult or impossible
 - Acute renal failure
 - If TLS is untreated, its progression may cause acute kidney failure, cardiac arrhythmias, seizures, loss of muscle control, and death.

^a There are data to support that fixed-dose rasburicase is very effective in adult patients.

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

Continued

TCLYM-B
1 OF 3

SUPPORTIVE CARE

Hemophagocytic lymphohistiocytosis (HLH)^b

- Syndrome of extreme immune activation resulting in life-threatening inflammation
- Clinical signs and symptoms may include: (these may overlap with features of underlying lymphoma)
 - Fever
 - Hepatosplenomegaly
 - Cytopenias (affecting 2 of 3 lineages in the peripheral blood)
 - ◊ Hemoglobin <9 g/dL
 - ◊ Platelets <100 k/dL
 - ◊ Neutrophils <1000/dL
 - Hypertriglyceridemia and/or hypofibrinogenemia
 - ◊ Fasting triglycerides >3.0 mmol/L (ie, >265 mg/dL)
 - ◊ Fibrinogen <1.5 g/L
 - Hemophagocytosis in bone marrow or spleen or lymph nodes
 - Ferritin >500 mg/L
 - Soluble interleukin-2 (IL-2) receptor (eg, sCD25) >2400 U/mL
 - Elevated transaminases and bilirubin
 - Elevated LDH
 - Elevated D-dimer
 - Elevated cerebrospinal fluid (CSF) cells and/or protein
- Diagnostic evaluation
 - Labs including CBC with differential, triglycerides, fibrinogen, ferritin, sCD25, liver function tests (LFTs), LDH, and D-dimer
 - Bone marrow biopsy
 - ◊ Consider repeat bone marrow biopsy if strong suspicion of HLH
 - Consider liver biopsy
- Management
 - Recommend expert consultation
 - Treatment of the underlying T-cell lymphoma with preference for etoposide- and steroid-containing regimens
 - Antiviral therapy - See Monoclonal Antibody Therapy and Viral Reactivation ([TCLYM-B](#))

Continued

^b HLH in adults is often associated with an underlying T-cell lymphoma. Diagnostic workup to confirm the lymphoma subtype and prompt initiation of treatment for underlying T-cell lymphoma is often required.

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

TCLYM-B
2 OF 3

SUPPORTIVE CARE

For other immunosuppressive situations, see NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections.

Monoclonal Antibody Therapy and Viral Reactivation

• Brentuximab Vedotin (anti-CD30 antibody-drug conjugate)

▸ Progressive multifocal leukoencephalopathy (PML):

- ◊ Caused by reactivation of the JC virus (JCV) and is usually fatal.
- ◊ Diagnosis made by PCR of CSF and in some cases brain biopsy.
- ◊ No known effective treatment.
- ◊ Clinical indications may include changes in behavior such as confusion, dizziness or loss of balance, difficulty talking or walking, and vision problems.

• Anti-CD52 Antibody Therapy: Alemtuzumab

▸ CMV reactivation:

- ◊ The current appropriate management is controversial; some NCCN Member Institutions use ganciclovir (PO or IV) preemptively if viremia is present, others only if viral load is rising.
- ◊ Herpes simplex virus (HSV) prophylaxis with acyclovir or equivalent.
- ◊ PJP prophylaxis with sulfamethoxazole/trimethoprim or equivalent.
- ◊ Consider antifungal prophylaxis.
- ◊ CMV viremia should be measured by quantitative PCR at least every 2 to 3 weeks.
- ◊ Consultation with an infectious disease expert may be necessary. See NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections.

Management of Tumor Flare Recommended for Patients Receiving Lenalidomide

- Tumor flare reactions: Painful lymph node enlargement or lymph node enlargement with evidence of local inflammation, occurring with treatment initiation; may also be associated with spleen enlargement, low-grade fever, and/or rash.
- Treatment: Steroids (eg, prednisone 25–50 mg PO for 5–10 days); antihistamines for rash and pruritus (eg, cetirizine 10 mg PO once daily or loratadine 10 mg PO daily).
- Prophylaxis: Consider in patients with bulky lymph nodes (>5 cm); administer steroids (eg, prednisone 20 mg PO for 5–7 days followed by rapid taper over 5–7 days).

Prevention of Pralatrexate-Induced Mucositis^{a,b,c}

- Vitamin B12 (cyanocobalamin) at a dose of 1,000 mcg intramuscular to be started no more than 10 weeks prior to starting therapy with pralatrexate and then every 8 to 10 weeks.
- Oral folic acid 1 to 1.25 mg daily to be started within 10 days of starting therapy and continuing for 30 days after the last dose of pralatrexate.
- Consider use of oral leucovorin 25 mg three times daily for two consecutive days (total of 6 doses), starting 24 hours after each dose of pralatrexate.

^a Mould DR, Sweeney K, Duffull SB et al. A population pharmacokinetic and pharmacodynamic evaluation of pralatrexate in patients with relapsed or refractory non-Hodgkin's or Hodgkin's lymphoma. Clin Pharmacol Ther 2009;86:190-196.

^b Shustov AR, Shinohara MM, Dakhil SR, et al. Management of mucositis with the use of leucovorin as adjunct to pralatrexate in treatment of peripheral t-cell lymphomas (PTCL) – Results from a prospective multicenter phase 2 clinical trial. Blood 2018;132:2910.

^c Koch E, Story SK, Geskin L. Preemptive leucovorin administration minimizes pralatrexate toxicity without sacrificing efficacy. Leuk Lymphoma 2013;54:2448-2451.

Version 1.2021 © National Comprehensive Cancer Network, Inc. 2020. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

TCLYM-B
3 OF 3