NCCN: Continuing Education

Target Audience: This activity is designed to meet the educational needs of oncologists, nurses, pharmacists, and other healthcare professionals who manage patients with cancer.

Accreditation Statements

In support of improving patient care, National Comprehensive Cancer Network (NCCN) is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medicine (ACCME): NCCN designates this journal-based CME activity for a maximum of 1.0 AMA PRA Category 1 Credit™. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Nursing (ANCC): NCCN designates this educational activity for a maximum of 1.0 contact hour.

Pharmacy (ACPE): NCCN designates this knowledge-based continuing education activity for 1.0 contact hour (0.1 CEUs) of continuing education credit. UAN: JA4008196-0000-21-006-H01-P

All clinicians completing this activity will be issued a certificate of participation. To participate in this journal CE activity: (1) review the educational content; (2) take the posttest with a 66% minimum passing score and complete the evaluation at <https://education.nccn.org/node/89409>; and (3) view/print certificate.

Pharmacists: You must complete the posttest and evaluation within 30 days of the activity. Continuing pharmacy education credit is reported to the CPE Monitor once you have completed the posttest and evaluation and claimed your credits. Before completing these requirements, be sure your NCCN profile has been updated with your NAPB e-profile ID and date of birth. Your credit cannot be reported without this information. If you have any questions, please e-mail education@nccn.org.

Release date: March 10, 2021; Expiration date: March 10, 2022

Learning Objectives:

Upon completion of this activity, participants will be able to:

- Integrate into professional practice the updates to the NCCN Guidelines for Non–Small Cell Lung Cancer
- Describe the rationale behind the decision-making process for developing the NCCN Guidelines for Non–Small Cell Lung Cancer

Disclosure of Relevant Financial Relationships

The NCCN staff listed below discloses no relevant financial relationships:

Kerrin M. Rosenthal, MA; Kimberly Callan, MS; Genevieve Emberger Hartzman, MA; Erin Hesler; Kristina M. Gregory, RN, MSN, OCN; Rashmi Kumar, PhD; Karen Kanefield; and Kathy Smith.

Individuals Who Provided Content Development and/or Authorship Assistance:

David S. Ettinger, MD, Panel Chair, has disclosed that he is as scientific advisor for BeyondSpring Pharmaceuticals and Takeda Pharmaceuticals North America, Inc.

Miranda Hughes, PhD, Oncology Scientist/Senior Medical Writer, NCCN, has disclosed that she has no relevant financial relationships.

To view all of the conflicts of interest for the NCCN Guidelines panel, go to [NCCN.org/disclosures/guidelinepanellisting.aspx](https://www.nccn.org/disclosures/guidelinepanellisting.aspx).

This activity is supported by educational grants from Agios Pharmaceuticals; AstraZeneca; Clovis Oncology, Inc.; Daiichi Sankyo; Eisai; Epizyme Inc.; Novartis; and Pharmacycics LLC, an AbbVie Company and Janssen Biotech, Inc., administered by Janssen Scientific Affairs, LLC. This activity is supported by an independent medical education grant from Bristol-Myers Squibb, and Regeneron Pharmaceuticals, Inc. and Sanofi Genzyme. This activity is supported by an independent medical educational grant from Mylan Inc. This activity is supported by a medical education grant from Karyopharm Therapeutics. This activity is supported by an independent educational grant from AbbVie.

Non–Small Cell Lung Cancer, Version 2.2021

Featured Updates to the NCCN Guidelines

David S. Ettinger, MD^{1,*}; Douglas E. Wood, MD²; Dara L. Aisner, MD, PhD³; Wallace Akerley, MD⁴; Jessica R. Bauman, MD⁵; Ankit Bharat, MD⁶; Debora S. Bruno, MD, MS⁷; Joe Y. Chang, MD, PhD⁸; Lucian R. Chirieac, MD⁹; Thomas A. D'Amico, MD¹⁰; Thomas J. Dilling, MD, MS¹¹; Jonathan Dowell, MD¹²; Scott Gettinger, MD¹³; Matthew A. Gubens, MD, MS¹⁴; Aparna Hegde, MD¹⁵; Mark Hennon, MD¹⁶; Rudy P. Lackner, MD¹⁷; Michael Lanuti, MD¹⁸; Ticiana A. Leal, MD¹⁹; Jules Lin, MD²⁰; Billy W. Loo Jr, MD, PhD²¹; Christine M. Lovly, MD, PhD²²; Renato G. Martins, MD, MPH²³; Erminia Massarelli, MD²⁴; Daniel Morgensztern, MD²⁴; Thomas Ng, MD²⁵; Gregory A. Otterson, MD²⁶; Sandip P. Patel, MD²⁷; Gregory J. Riely, MD, PhD²⁸; Steven E. Schild, MD²⁹; Theresa A. Shapiro, MD, PhD¹; Aditi P. Singh, MD³⁰; James Stevenson, MD⁷; Alda Tam, MD⁸; Jane Yanagawa, MD³¹; Stephen C. Yang, MD¹; Kristina M. Gregory, RN, MSN, OCN^{32,*}; and Miranda Hughes, PhD^{32,*}

ABSTRACT

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for Non–Small Cell Lung Cancer (NSCLC) address all aspects of management for NSCLC. These NCCN Guidelines Insights focus on recent updates to the NCCN Guidelines regarding targeted therapies, immunotherapies, and their respective biomarkers.

J Natl Compr Canc Netw 2021;19(3):254–266
doi: 10.6004/jnccn.2021.0013

NCCN CATEGORIES OF EVIDENCE AND CONSENSUS

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Clinical trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

PLEASE NOTE

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines[®]) are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. **The NCCN Guidelines Insights highlight important changes in the NCCN Guidelines recommendations from previous versions. Colored markings in the algorithm show changes and the discussion aims to further the understanding of these changes by summarizing salient portions of the panel's discussion, including the literature reviewed.**

The NCCN Guidelines Insights do not represent the full NCCN Guidelines; further, the National Comprehensive Cancer Network[®] (NCCN[®]) makes no representations or warranties of any kind regarding their content, use, or application of the NCCN Guidelines and NCCN Guidelines Insights and disclaims any responsibility for their application or use in any way.

The complete and most recent version of these NCCN Guidelines is available free of charge at NCCN.org.

© National Comprehensive Cancer Network, Inc. 2021. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN.

¹The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins; ²Fred Hutchinson Cancer Research Center/Seattle Cancer Care Alliance; ³University of Colorado Cancer Center; ⁴Huntsman Cancer Institute at the University of Utah; ⁵Fox Chase Cancer Center; ⁶Robert H. Lurie Comprehensive Cancer Center of Northwestern University; ⁷Case Comprehensive Cancer Center/University Hospitals Seidman Cancer Center and Cleveland Clinic Taussig Cancer Institute; ⁸The University of Texas MD Anderson Cancer Center; ⁹Dana-Farber/Brigham and Women's Cancer Center; ¹⁰Duke Cancer Institute; ¹¹Moffitt Cancer Center; ¹²UT Southwestern Simmons Comprehensive Cancer Center; ¹³Yale Cancer Center/Smilow Cancer Hospital; ¹⁴UCSF Helen Diller Family Comprehensive Cancer Center; ¹⁵O'Neal Comprehensive Cancer Center at UAB; ¹⁶Roswell Park Comprehensive Cancer Center; ¹⁷Fred & Pamela Buffett Cancer Center; ¹⁸Massachusetts General Hospital Cancer Center; ¹⁹University of Wisconsin Carbone Cancer Center; ²⁰University of Michigan Rogel Cancer Center; ²¹Stanford Cancer Institute; ²²Vanderbilt-Ingram Cancer Center; ²³City of Hope National Medical Center; ²⁴Siteman Cancer Center at Barnes-Jewish Hospital and Washington University School of Medicine; ²⁵The University of Tennessee Health Science Center; ²⁶The Ohio State University Comprehensive Cancer Center - James Cancer Hospital and Solove Research Institute; ²⁷UC San Diego Moores Cancer Center; ²⁸Memorial Sloan Kettering Cancer Center; ²⁹Mayo Clinic Cancer Center; ³⁰Abramson Cancer Center at the University of Pennsylvania; ³¹UCLA Jonsson Comprehensive Cancer Center; and ³²National Comprehensive Cancer Network.

*Provided content development and/or authorship assistance.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
 The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-4

Overview

Lung cancer is the leading cause of cancer death in the United States.¹ In 2021, an estimated 235,760 new cases (119,100 in men and 116,660 in women) of lung and bronchial cancer will be diagnosed, and 131,880 deaths (69,410 in men and 62,470 in women) are estimated to occur.¹ Only 26% of all patients with non–small cell lung cancer (NSCLC) are alive ≥5 years after diagnosis.² The 5-year relative survival rate for metastatic disease is approximately 6% when patients receive historic cytotoxic chemotherapy regimens.² However, certain patients with metastatic NSCLC who are eligible for newer targeted therapies or immunotherapies are now surviving longer, with 5-year survival rates ranging from 15% to 50%, depending on the biomarker.^{3–13}

These NCCN Guidelines Insights focus on recent updates in targeted therapies, immunotherapies, and their respective biomarkers for eligible patients with NSCLC. The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for NSCLC address all aspects of management for NSCLC. For the 2020 and 2021 updates, several new targeted therapies (or new indications for therapies), including capmatinib, lorlatinib, pralsetinib, seliperatinib, and fam-trastuzumab

deruxtecan, are now recommended in the NCCN Guidelines for eligible patients with metastatic NSCLC who have certain actionable biomarkers.^{14–20} These NCCN Guidelines Insights detail the reasons behind the recent revisions and provide a valuable resource for busy healthcare providers who need to quickly learn about the recent recommendations to improve outcomes for their patients with metastatic NSCLC. Unless otherwise indicated, all NCCN recommendations are category 2A (the complete version of these guidelines is available at NCCN.org).

Biomarkers

A *predictive* (also known as *actionable*) biomarker has a corresponding specific targeted therapy(ies) that has been shown to improve outcomes in patients with the predictive biomarker (eg, *ALK* rearrangements are targeted by alectinib, brigatinib, lorlatinib, and other *ALK* inhibitors). A *prognostic* biomarker is indicative of patient survival independent of the treatment received, because the biomarker is an indicator of the innate tumor aggressiveness (eg, *KRAS* mutations). In the NCCN Guidelines, key established predictive molecular biomarkers include *ALK* rearrangements, *BRAF*V600E point

^a See Principles of Pathologic Review (NSCL-A).

^c Temel JS, et al. *N Engl J Med* 2010;363:733-742.

^{kk} If there is insufficient tissue to allow testing for all of *EGFR*, *ALK*, *ROS1*, *BRAF*, *NTRK1/2/3*, *MET*, and *RET*, repeat biopsy and/or plasma testing should be done. If these are not feasible, treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.

^{ll} See Principles of Molecular and Biomarker Analysis (NSCL-H).

^{mmm} The NCCN NSCLC Guidelines Panel strongly advises broader molecular profiling with the goal of identifying rare driver mutations for which effective drugs may already be available, or to appropriately counsel patients regarding the availability of clinical trials. Broad molecular profiling is a key component of the improvement of care of patients with NSCLC. See Emerging Biomarkers to Identify Patients for Therapies (NSCL-I).

ⁿⁿ Lam VK, et al. *Clin Lung Cancer* 2019;20:30-36.e3; Sands JM, et al. *Lung Cancer* 2020;140:35-41.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-18

mutations, sensitizing *EGFR* mutations, *MET* exon 14 skipping mutations, *NTRK1/2/3* gene fusions, *RET* rearrangements, and *ROS1* rearrangements; PD-L1 expression is a key established immune biomarker (see NSCL-18, page 257). The NCCN NSCLC Panel recommends testing for these key established predictive biomarkers after patients have been diagnosed with metastatic NSCLC and ideally before initial treatment, because effective targeted therapy or immunotherapy is available depending on biomarker test results.¹⁴

Emerging predictive biomarkers also have corresponding specific targeted therapies, but fewer data are available to support use of these targeted therapies compared with the targeted therapies for the key established predictive biomarkers (see NSCL-I, page 264). Several emerging biomarkers have become established biomarkers in the NCCN Guidelines after more clinical trial data were published for their corresponding targeted therapies. For example, *MET* exon 14 skipping mutations and *RET* rearrangements were moved from the emerging biomarkers section to the established biomarker section of the algorithm for the 2020 update (see NSCL-18, page 257).^{14,16–18} For the version 1.2021 (v1.2021) update, the panel clarified that biomarker testing is recommended in certain patients with metastatic (stage IV) disease, including M1a, M1b, and

M1c. The panel also recommends considering biomarker testing for *EGFR* mutations in surgical tissue or biopsies from patients with completely resected stage IB–IIIA NSCLC to determine whether adjuvant osimertinib can be considered for these patients (see NSCL-4, page 256, and next section).^{14,21}

Molecular Biomarkers

The panel recommends molecular testing using a validated test(s) that assesses a minimum of the following potential genetic variants: *ALK* rearrangements (category 1), *BRAF* mutations, *EGFR* mutations (category 1), *MET* exon 14 skipping mutations, *NTRK1/2/3* gene fusions, *RET* rearrangements, and *ROS1* rearrangements.¹⁴ Both FDA-approved companion diagnostics and laboratory-developed test platforms are available to evaluate for these and other analytes. The NCCN Guidelines for NSCLC provide recommendations for specific biomarkers that should be assessed in patients who have been diagnosed with NSCLC, and recommend testing techniques for the key established biomarkers, but do not endorse any specific commercially available biomarker assays. For the 2020 update, the panel recommends that molecular testing be performed via a broad, panel-based

ALK REARRANGEMENT POSITIVE^{II}

^{II} See Principles of Molecular and Biomarker Analysis (NSCL-H).

^{OO} See Targeted Therapy or Immunotherapy for Advanced or Metastatic Disease (NSCL-J).

^{PP} For performance status 0–4.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-23

approach, most typically performed by next-generation sequencing (NGS), so that testing is done for all of the actionable biomarkers at the same time, including the established and emerging biomarkers. This testing technique helps ensure that there is sufficient tissue to test for all of the actionable biomarkers. However, some gene fusions are difficult to detect using DNA-based NGS. For patients who, in broad panel testing, do not have identifiable driver oncogenes (especially never smokers), RNA-based NGS should be considered, if not already performed, to maximize detection of fusion events.²²

ALK, *EGFR*, *BRAF*, *METex14*, *NTRK1/2/3*, *RET*, and *ROS1* status should be known before deciding whether to use either targeted therapy or immunotherapy with or without chemotherapy regimens. If it is not feasible to perform molecular testing, then patients are treated as though they do not have driver oncogenes.^{23–27} Note that the panel recommends that testing for PD-L1 expression levels, which is an immune biomarker, be performed using an immunohistochemistry test that is an FDA companion diagnostic or has been shown to have equivalent performance to an approved companion diagnostic.^{28,29} The immunohistochemical evaluation of PD-L1 that guides use of pembrolizumab is based on tumor proportion score

(TPS), which is the percentage of viable tumor cells showing partial or complete membrane staining at any intensity.

For the 2021 and 2020 updates, new content was added for *EGFR* mutations and *NTRK1/2/3* gene fusions; recommended testing techniques were added for *METex14* skipping mutations, *NTRK1/2/3* gene fusions, and *RET* fusions (see NSCL-H 2 and 4, pages 263 and 264)¹⁴; and content was revised for the key established biomarkers, such as *EGFR* mutations, and for the emerging biomarkers, such as tumor mutational burden (TMB) (see NSCL-18 and NSCL-I, pages 257 and 264). For the v1.2021 update, the panel decided that routine molecular testing should be considered in all patients with metastatic NSCLC squamous cell carcinoma (see NSCL-18, page 257).¹⁴ Therefore, characteristics—such as smoking status, small biopsy specimens, and mixed histology—should no longer be used when considering whether to perform biomarker testing on patients with metastatic NSCLC squamous cell carcinoma. This decision is based on recent data showing that patients with metastatic squamous cell carcinoma also have actionable biomarkers, such as *EGFR* mutations, although at a lower incidence than those with metastatic NSCLC adenocarcinoma.^{23,30–32} The cumulative incidence

ROS1 REARRANGEMENT POSITIVE^{II}

^{II} See Principles of Molecular and Biomarker Analysis (NSCL-H).

^{OO} See Targeted Therapy or Immunotherapy for Advanced or Metastatic Disease (NSCL-J).

^{PP} For performance status 0–4.

^{TT} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{EEE} Entrectinib may be better for patients with brain metastases.

^{FFF} Entrectinib is primarily for patients with CNS progression after crizotinib.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-26

of actionable alterations in tumors carrying a diagnosis of metastatic squamous cell carcinoma is sufficient to justify consideration of molecular testing. The panel now feels that molecular testing should be considered in patients with metastatic squamous cell carcinoma based on the effectiveness of targeted therapies.^{23,32} The panel also clarified that *NTRK1/2/3* gene fusions are established predictive molecular biomarkers for the 2021 update. Typically, the key established molecular biomarkers do not overlap; most patients with actionable mutations only have one actionable mutation.^{23,32} Although *KRAS* mutations are not actionable at this time, inclusion on panel-based testing is informative, because it generally excludes the presence of an actionable alteration.

Immunotherapy with or without chemotherapy is recommended for patients who do not have actionable molecular biomarkers. If patients have both a molecular biomarker and high PD-L1 expression levels, targeted therapy is usually recommended over immunotherapy with or without chemotherapy based on data showing that targeted therapy yields higher response rates compared with immunotherapy in the first-line setting, targeted therapy is better tolerated, and most patients with an actionable molecular biomarker will only have a modest or

slight response to immunotherapy.^{23–27} Response rates for immunotherapy are lower in patients with *EGFR* and *ALK* variants; however, pembrolizumab with or without chemotherapy may be considered for a heavy smoker with PD-L1 levels of 100% and a *BRAF* V600E mutation.

Recent data show that certain patients with completely resected early-stage NSCLC who have sensitizing *EGFR* mutations have longer duration of disease-free survival if they receive adjuvant osimertinib versus placebo.²¹ ADAURA, a phase III randomized trial in 682 patients, showed that 90% (95% CI, 84%–93%) of patients with stage II–IIIA NSCLC receiving osimertinib were alive and disease-free at 24 months compared with 44% (95% CI, 37%–51%) receiving placebo (HR, 0.17; 99.06% CI, 0.11–0.26; *P* < .001).²¹ Disease-free survival was also improved in patients who received adjuvant chemotherapy before osimertinib compared with placebo. Nine patients in the osimertinib group and 20 in the placebo group died. It will be interesting to see whether patients with other actionable molecular biomarkers, such as *ALK* rearrangements, also have improved survival with targeted agents in the same setting. For the v1.2021 update, the panel recommends considering adjuvant osimertinib for patients with completely resected *EGFR* mutation–positive stage

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-31

IIB–IIIA and high-risk stage IB–IIA NSCLC who received previous adjuvant chemotherapy or are ineligible to receive platinum chemotherapy based on FDA approval and clinical trial results (see NSCL-E, page 261).²¹ High-risk features are described in the algorithm (see NSCL-4, page 256).

For the version 2.2021 (v2.2021) update, the panel now recommends lorlatinib (category 1) as another preferred first-line therapy option for patients with *ALK* rearrangement–positive metastatic NSCLC (see NSCL-23, page 258).¹⁵ The panel revised the preference stratification for brigatinib (category 1) to a preferred first-line therapy option for patients with *ALK* rearrangement–positive metastatic NSCLC in the v1.2021 update (see NSCL-23, page 258).^{33,34} The panel clarified that entrectinib may be better for patients with *ROS1* rearrangement–positive metastatic NSCLC who have brain metastases, and entrectinib is now recommended as a subsequent therapy option for patients with *ROS1*-positive disease who have CNS progression after crizotinib (see NSCL-26, page 259).^{35,36} Entrectinib was designed to cross the blood–brain barrier.³⁵ However, entrectinib is less effective for certain resistant *ROS1* variants.³⁵ For the v1.2021 update, the panel added capmatinib as a treatment option for patients with high-level *MET* amplification, which is an emerging

biomarker (see NSCL-I, page 264).¹⁶ The panel also added fam-trastuzumab deruxtecan as a treatment option for patients with *ERBB2* (*HER2*) mutation–positive metastatic NSCLC, which is another emerging biomarker (see NSCL-I, page 264).¹⁹ The panel deleted single-agent dabrafenib as a treatment option for patients with *BRAF* V600E mutation–positive metastatic NSCLC who cannot tolerate combination therapy with dabrafenib + trametinib; single-agent vemurafenib remains an option in this setting.^{37,38} Although less toxic, dabrafenib monotherapy is less effective than combination therapy with dabrafenib + trametinib.³⁹

For the 2020 updates, the panel added selpercatinib and pralsetinib as preferred first-line therapy options for patients with *RET* rearrangement–positive metastatic NSCLC.^{17,18} Selpercatinib and pralsetinib are also recommended as preferred subsequent therapy options in this setting if *RET* inhibitors have not been previously given as first-line therapy. The panel also added capmatinib as a preferred first-line therapy option for patients with *METex14* skipping mutations.¹⁶ Likewise, capmatinib is also recommended as a preferred subsequent therapy option in this setting if *MET* inhibitors have not been previously given as first-line therapy.

SYSTEMIC THERAPY REGIMENS FOR NEOADJUVANT AND ADJUVANT THERAPY

Preferred (nonsquamous)

- Cisplatin 75 mg/m² day 1, pemetrexed 500 mg/m² day 1 every 21 days for 4 cycles¹

Preferred (squamous)

- Cisplatin 75 mg/m² day 1; gemcitabine 1250 mg/m² days 1 and 8, every 21 days for 4 cycles²
- Cisplatin 75 mg/m² day 1; docetaxel 75 mg/m² day 1 every 21 days for 4 cycles³

Other Recommended

- Cisplatin 50 mg/m² days 1 and 8; vinorelbine 25 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles⁴
- Cisplatin 100 mg/m² day 1; vinorelbine 30 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles^{5,6}
- Cisplatin 75–80 mg/m² day 1; vinorelbine 25–30 mg/m² days 1 and 8, every 21 days for 4 cycles
- Cisplatin 100 mg/m² day 1; etoposide 100 mg/m² days 1–3, every 28 days for 4 cycles⁵

Useful in Certain Circumstances**Chemotherapy Regimens for Patients with Comorbidities or Patients Not Able to Tolerate Cisplatin**

- Carboplatin AUC 6 day 1, paclitaxel 200 mg/m² day 1, every 21 days for 4 cycles⁷
- Carboplatin AUC 5 day 1, gemcitabine 1000 mg/m² days 1 and 8, every 21 days for 4 cycles⁸
- Carboplatin AUC 5 day 1, pemetrexed 500 mg/m² day 1 for nonsquamous every 21 days for 4 cycles⁹

All regimens can be used for sequential chemotherapy/RT.**Previous Adjuvant Chemotherapy or Ineligible for Platinum-Based Chemotherapy**

- Osimertinib 80 mg daily¹⁰

► Consider osimertinib for patients with completely resected Stage IB–IIIA *EGFR* mutation-positive NSCLC who received previous adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy.

¹ Kreuter M, Vansteenkiste J, Fishcer JR, et al. Randomized phase 2 trial on refinement of early-stage NSCLC adjuvant chemotherapy with cisplatin and pemetrexed versus cisplatin and vinorelbine: the TREAT study. *Ann Oncol* 2013;24:986-992.

² Pérol M, Chouaid C, Pérol D, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin-gemcitabine induction chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 2012;30:3516-3524.

³ Fossella F, Pereira JR, von Pawel J, et al. Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. *J Clin Oncol* 2003;21:3016-3024.

⁴ Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-lung cancer. *N Engl J Med* 2005;352:2589-2597.

⁵ Ariagada R, Bergman B, Dunant A, et al. The International Adjuvant Lung Cancer Trial Collaborative Group. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small cell lung cancer. *N Engl J Med* 2004;350:351-360.

⁶ Douillard JY, Rosell R, De Lena M, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB–IIIA non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. *Lancet Oncol* 2006;7:719-727.

⁷ Strauss GM, Herndon III JE, Maddaus MA, et al. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small cell lung cancer: CALGB 9633 with the Cancer and Leukemia Group B, Radiation Therapy Oncology Group, and North Central Cancer Treatment Group Study Groups. *J Clin Oncol* 2008;26:5043-5051.

⁸ Usami N, Yokoi K, Hasegawa Y, et al. Phase II study of carboplatin and gemcitabine as adjuvant chemotherapy in patients with completely resected non-small cell lung cancer: a report from the Central Japan Lung Study Group, CJLSG 0503 trial. *Int J Clin Oncol* 2010;15:583-587.

⁹ Zhang L, Ou W, Liu Q, et al. Pemetrexed plus carboplatin as adjuvant chemotherapy in patients with curative resected non-squamous non-small cell lung cancer. *Thorac Cancer* 2014;5:50-56.

¹⁰ Wu Y-L, Tsuboi M, He J, et al. Osimertinib in resected *EGFR*-mutated non-small-cell lung cancer. *N Engl J Med* 2020;383:1711-1723.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved. The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-E

Immune Biomarkers

Several immune biomarkers, including PD-L1 expression levels and TMB, have been assessed in clinical trials to see if they are useful for predicting whether patients with metastatic NSCLC will respond to various immunotherapy regimens, such as single-agent pembrolizumab or nivolumab + ipilimumab.⁴⁰⁻⁴³ The response rate to immune checkpoint inhibitors varies depending on the regimen and setting. The highest response rates occur in patients with PD-L1 levels $\geq 50\%$ in the first-line setting and no actionable molecular biomarkers (approximately 40% response rate to single-agent immunotherapy; approximately 60% to chemoimmunotherapy).^{40,41,44-46} However, only approximately 30% of patients with metastatic NSCLC have PD-L1 levels $\geq 50\%$.^{44,47} Although PD-L1 expression level is widely used as an immune biomarker, it is not an ideal biomarker because some patients with low PD-L1 expression levels respond to immunotherapy and others with high levels do not respond to immunotherapy.^{48,49} PD-L1 expression levels are useful for deciding whether to use single-agent immunotherapy or combination immunotherapy.

TMB is an approximate measure of the total number of somatic mutations.⁵⁰ Theoretically, high TMB levels

will correlate with high neoantigen levels that will activate an antitumor immune response.⁴⁸ TMB levels are typically high in patients with NSCLC who are smokers or former smokers. Low TMB is more commonly detected in never-smokers.^{22,51} Preliminary data for progression-free survival from CheckMate 227, a phase III randomized trial with a complex design, suggested that TMB might be a useful immune biomarker for deciding whether to use immunotherapy in patients with metastatic NSCLC.⁴² However, updated data from this trial showed that overall survival was improved with nivolumab + ipilimumab regardless of TMB or PD-L1 expression levels.⁴³ In addition, combining TMB with PD-L1 expression level also did not correlate with overall survival.

Several trials have shown that high TMB levels do not correlate with PD-L1 expression levels in patients with NSCLC.^{42,43,52,53} KEYNOTE-158, a phase II trial, assessed TMB levels in patients with solid tumors who received pembrolizumab as second-line therapy; however, none of the patients had NSCLC.⁵⁴ TMB does not identify patients who will respond to chemotherapy; therefore, it has limited value for assessing combination immunotherapy + chemotherapy regimens.⁴⁸ TMB is also

CONCURRENT CHEMORADIATION REGIMENS

Concurrent Chemoradiation Regimens[¶]**Preferred (nonsquamous)**

- Carboplatin AUC 5 on day 1, pemetrexed 500 mg/m² on day 1 every 21 days for 4 cycles; concurrent thoracic RT^{1,*,†,‡}
- Cisplatin 75 mg/m² on day 1, pemetrexed 500 mg/m² on day 1 every 21 days for 3 cycles; concurrent thoracic RT^{2,3,*,†,‡} ± additional 4 cycles of pemetrexed 500 mg/m²^{†,§}
- Paclitaxel 45–50 mg/m² weekly; carboplatin AUC 2, concurrent thoracic RT^{4,*,†,‡} ± additional 2 cycles every 21 days of paclitaxel 200 mg/m² and carboplatin AUC 6^{†,§}
- Cisplatin 50 mg/m² on days 1, 8, 29, and 36; etoposide 50 mg/m² days 1–5 and 29–33; concurrent thoracic RT^{5,6,*,†,‡}

Preferred (squamous)

- Paclitaxel 45–50 mg/m² weekly; carboplatin AUC 2, concurrent thoracic RT^{6,*,†,‡} ± additional 2 cycles every 21 days of paclitaxel 200 mg/m² and carboplatin AUC 6^{†,§}
- Cisplatin 50 mg/m² on days 1, 8, 29, and 36; etoposide 50 mg/m² days 1–5 and 29–33; concurrent thoracic RT^{5,6,*,†,‡}

Consolidation Immunotherapy for Patients with Unresectable Stage II/III NSCLC, PS 0–1, and No Disease Progression After 2 or More Cycles of**Definitive Concurrent Chemoradiation**

Durvalumab 10 mg/kg IV every 2 weeks or 1500 mg every 4 weeks for up to 12 months (patients with a body weight of ≥30 kg)^{7,8} (category 1 for stage III; category 2A for stage II)

[¶] For patients with superior sulcus tumors, the recommendation is for 2 cycles concurrent with radiation therapy and 2 more cycles after surgery. Rusch VW, Giroux DJ, Kraut MJ, et al. Induction chemoradiation and surgical resection for superior sulcus non-small-cell lung carcinomas: long-term results of Southwest Oncology Group Trial 9416 (Intergroup Trial 0160). *J Clin Oncol* 2007;25:313-318.

* Regimens can be used as preoperative/adjuvant chemotherapy/RT.

† Regimens can be used as definitive concurrent chemotherapy/RT.

‡ For eligible patients, durvalumab may be used after noted concurrent chemo/RT regimens.

§ If using durvalumab, an additional 2 cycles of chemotherapy is not recommended, if patients have not received full-dose chemotherapy concurrently with RT.

¹ Govindan R, Bogart J, Stinchcombe T, et al. Randomized phase II study of pemetrexed, carboplatin, and thoracic radiation with or without cetuximab in patients with locally advanced unresectable non-small-cell lung cancer: Cancer and Leukemia Group B trial 30407. *J Clin Oncol* 2011;29:3120-3125.

² Choy H, Gerber DE, Bradley JD, et al. Concurrent pemetrexed and radiation therapy in the treatment of patients with inoperable stage III non-small cell lung cancer: a systematic review of completed and ongoing studies. *Lung Cancer* 2015;87:232-240.

³ Senan S, Brade A, Wang LH, et al. PROCLAIM: randomized phase III trial of pemetrexed-cisplatin or etoposide-cisplatin plus thoracic radiation therapy followed by consolidation chemotherapy in locally advanced nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2016;34:953-962.

⁴ Bradley JD, Paulus R, Komaki R, et al. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (RTOG 0617): a randomised, two-by-two factorial phase 3 study. *Lancet Oncol* 2015;16:187-199.

⁵ Albain KS, Crowley JJ, Turrisi AT III, et al. Concurrent cisplatin, etoposide, and chest radiotherapy in pathologic stage IIIB non-small-cell lung cancer: A Southwest Oncology Group Phase II Study. *SWOG 9019. J Clin Oncol* 2002;20:3454-3460.

⁶ Curran WJ Jr, Paulus R, Langer CJ, et al. Sequential vs. concurrent chemoradiation for stage III non-small cell lung cancer: randomized phase III trial RTOG 9410. *J Natl Cancer Inst* 2011;103:1452-1460.

⁷ Antonia SJ, Villegas A, Daniel D, et al. Overall survival with durvalumab after chemoradiotherapy in stage III NSCLC. *N Engl J Med* 2018;379:2342-2550.

⁸ Baverel PG, Dubois VFS, Jin CY, et al. Population pharmacokinetics of durvalumab in cancer patients and association with longitudinal biomarkers of disease status. *Clin Pharmacol Ther* 2018;103:631-642.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-F

not an ideal immune biomarker, because some patients with low levels experience a response to immunotherapy and others with high levels do not.⁴⁸

In addition to the lack of clinical data to support use of TMB as an immune biomarker, there are technical problems with measuring TMB,⁵⁰ including lack of agreement on the definition of a cutoff for designating high TMB levels and lack of standardization of TMB measurements across laboratories.⁵⁰ PD-L1 expression level is a more useful immune biomarker than TMB for deciding how to use immunotherapy, because test results are obtained more quickly, less tissue is needed for testing, and data demonstrate relative reproducibility across platforms and individuals. For the v1.2021 update, the panel decided to remove TMB as an emerging immune biomarker for patients with metastatic NSCLC based on clinical trial data, concerns about variable TMB measurements, and other issues as previously described (see NSCL-I, page 264).^{14,43,50} Currently, the NCCN Guidelines do not recommend measurement of TMB levels before deciding whether to use nivolumab + ipilimumab combined with or without chemotherapy or to use other immune checkpoint inhibitors, such as pembrolizumab.¹⁴

For the v1.2021 update, the panel added a recommendation (category 2A) for consolidation immunotherapy with single-agent durvalumab for patients with unresectable stage II NSCLC who have not experienced disease progression after definitive concurrent chemoradiation. Previously, the durvalumab recommendation had been restricted to patients with unresectable stage III NSCLC (category 1) (see NSCL-F, page 262).⁵⁵ The panel also revised the recommendation for atezolizumab monotherapy to category 1 (from category 2A) as a preferred treatment option for patients with PD-L1 expression levels of ≥50% and negative for actionable biomarkers (see NSCL-31, page 260).⁴⁵ The recommendation for nivolumab + ipilimumab was also revised to category 1 (from category 2A) for patients with PD-L1 expression levels of ≥1% and negative for actionable biomarkers (see NSCL-31, page 260).

Summary

These NCCN Guidelines Insights focus on recent updates in targeted therapies, immunotherapies, and their respective biomarkers for eligible patients with NSCLC. For a list of the recent updates, see the complete version of these guidelines at NCCN.org. The NCCN Guidelines for

PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS

• Molecular Targets for Analysis

- ▶ In general, the mutations/alterations described below are seen in a non-overlapping fashion, although between 1%–3% of NSCLC may harbor concurrent alterations.
- ▶ **EGFR** (Epidermal Growth Factor Receptor) Gene Mutations: EGFR is a receptor tyrosine kinase normally found on the surface of epithelial cells and is often overexpressed in a variety of human malignancies.
 - ◊ The most commonly described mutations in *EGFR* (exon 19 deletions, *p.L858R* point mutation in exon 21) are associated with responsiveness to oral EGFR tyrosine kinase inhibitor (TKI) therapy; most recent data indicate that tumors that do not harbor a sensitizing *EGFR* mutation should not be treated with EGFR TKI in any line of therapy.
 - ◊ Consider adding molecular testing for *EGFR* mutation to be performed on diagnostic biopsy or post-surgical resection sample to ensure the *EGFR* mutation results are available for adjuvant treatment decisions for patients with stage IB to IIIA NSCLC.
 - ◊ Many of the less commonly observed alterations in EGFR, which cumulatively account for ~10% of *EGFR*-mutation positive NSCLC (ie, exon 19 insertions, *p.L861Q*, *p.G719X*, *p.S768I*) are also associated with responsiveness to EGFR TKI therapy, although the number of studied patients is lower.
 - ◊ *EGFR* exon 20 (*EGFR*ex20) mutations are a heterogeneous group, some of which are responsive to targeted therapy and that require detailed knowledge of the specific alteration.
 - *EGFR* *p.T790M* is most commonly observed as a mutation that arises in response to and as a mechanism of resistance to first- and second-generation EGFR TKI. In patients with progression on first- or second-generation TKI with *p.T790M* as the primary mechanism of resistance, third-generation TKIs are typically efficacious. If *p.T790M* is observed in the absence of prior EGFR TKI therapy, genetic counseling and possible germline genetic testing is warranted.
 - Most other *EGFR*ex20 alterations are a diverse group of in-frame duplication or insertion mutations.
 - These are generally associated with lack of response to EGFR TKI therapy, with select exceptions:
 - p.A763_Y764insFQEA* is associated with sensitivity to TKI therapy
 - p.A763_Y764insLQEA* may be associated with sensitivity to TKI therapy
 - For this reason, the specific sequence of *EGFR*ex20 insertion mutations is important, and some assays will identify the presence of an *EGFR*ex20 insertion without specifying the sequence. In this scenario, additional testing to further clarify the *EGFR*ex20 insertion is indicated.
 - ◊ As use of NGS testing increases, additional *EGFR* variants are increasingly identified; however, the clinical implications of individual alterations are unlikely to be well established.
 - ◊ Some clinicopathologic features—such as smoking status, ethnicity, and histology—are associated with the presence of an *EGFR* mutation; however, these features should not be utilized in selecting patients for testing.
 - ◊ Testing Methodologies: Real-time PCR, Sanger sequencing (ideally paired with tumor enrichment), and NGS are the most commonly deployed methodologies for examining *EGFR* mutation status.

Continued

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-H
2 OF 5

NSCLC address all aspects of management for NSCLC. For the 2020 and 2021 updates, several new targeted therapies or new indications for therapies, including capmatinib, lorlatinib, pralsetinib, selipercatinib, and fam-trastuzumab deruxtecan, are now recommended in the NCCN Guidelines for eligible patients with metastatic NSCLC who have certain actionable biomarkers.^{14,16–20} For the v2.2021 update, the panel now recommends lorlatinib (category 1) as another preferred first-line therapy option for patients with *ALK* rearrangement–positive metastatic NSCLC.¹⁵ For the v1.2021 update, the panel recommends considering adjuvant osimertinib for patients with completely resected *EGFR* mutation–positive stage IIB–IIIA and high-risk stage IB–IIA NSCLC who received previous adjuvant chemotherapy or are ineligible to receive platinum chemotherapy.^{14,21}

The panel revised the preference stratification for brigatinib (category 1) to a preferred first-line therapy option for patients with *ALK* rearrangement–positive metastatic NSCLC.^{33,34} The panel clarified that entrectinib may be better for patients with *ROS1* rearrangement–positive metastatic NSCLC who have brain metastases; entrectinib is recommended as a

subsequent therapy option for patients with *ROS1*-positive disease who have central nervous system progression after crizotinib.^{35,36} For the v1.2021 update, the panel decided that routine molecular biomarker testing should be considered in all patients with metastatic NSCLC squamous cell carcinoma; *EGFR* mutation biomarker testing can also be considered for patients with completely resected stage IB–IIIA NSCLC.^{14,21,30,31} Content was revised for the key established biomarkers, such as *EGFR* mutations. TMB, which is an immune biomarker, was deleted from the NCCN Guidelines based on clinical trial data, concerns about variable TMB measurements, and other issues.^{14,43,50} TMB was previously listed as an emerging biomarker. Currently, the NCCN Guidelines do not recommend measurement of TMB levels before deciding on the use of nivolumab + ipilimumab combined with or without chemotherapy or the use of other immune checkpoint inhibitors, such as pembrolizumab.¹⁴

To participate in this journal CE activity, go to
<https://education.nccn.org/node/89409>

PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS

- **Molecular Targets for Analysis (continued)**
 - ▶ **MET** (mesenchymal-epithelial transition) *exon 14 (METex14)* skipping variants: MET is a receptor tyrosine kinase. A mutation that results in loss of *exon 14* can occur in NSCLC. Loss of *METex14* leads to dysregulation and inappropriate signaling.
 - ◊ The presence of *METex14* skipping mutation is associated with responsiveness to oral MET TKIs.
 - ◊ A broad range of molecular alterations lead to *METex14* skipping.
 - ◊ Testing Methodologies: NGS-based testing is the primary method for detection of *METex14* skipping events, with RNA-based NGS demonstrating improvement in detection. IHC is not a method for detection of *METex14* skipping.
 - ▶ **RET** (rearranged during transfection) Gene Rearrangements: RET is a receptor tyrosine kinase that can be rearranged in NSCLC, resulting in dysregulation and inappropriate signaling through the RET kinase domain.
 - ◊ Common fusion partners are *KIF5B*, *NCOA4*, and *CCDC6*; however, numerous other fusion partners have been identified.
 - ◊ The presence of a *RET* rearrangement is associated with responsiveness to oral RET TKIs regardless of fusion partner.
 - ◊ Testing Methodologies: FISH break-apart probe methodology can be deployed; however, it may under-detect some fusions. Targeted real-time reverse-transcriptase PCR assays are utilized in some settings, although they are unlikely to detect fusions with novel partners. NGS-based methodology has a high specificity, and RNA-based NGS is preferable to DNA-based NGS for fusion detection.
 - ▶ **NTRK1/2/3** (neurotrophic tyrosine receptor kinase) gene fusions
 - ◊ *NTRK1/2/3* are tyrosine receptor kinases that are rarely rearranged in NSCLC as well as in other tumor types, resulting in dysregulation and inappropriate signaling.
 - ◊ Numerous fusion partners have been identified.
 - ◊ To date, no specific clinicopathologic features, other than absence of other driver alterations, have been identified in association with these fusions.
 - ◊ Point mutations in *NTRK1/2/3* are generally non-activating and have not been studied in association with targeted therapy.
 - ◊ Testing Methodologies: Various methodologies can be used to detect *NTRK1/2/3* gene fusions, including: FISH, IHC, PCR, and NGS; false negatives may occur. IHC methods are complicated by baseline expression in some tissues. FISH testing may require at least 3 probe sets for full analysis. NGS testing can detect a broad range of alterations. DNA-based NGS may under-detect *NTRK1* and *NTRK3* fusions.
- In the event that a complete assessment for all biomarkers cannot be reasonably accomplished prior to initiation of therapy, consider repeat panel testing or selected biomarker testing at progression on first-line therapy if a lesion can be accessed for sampling and testing.
- Testing in the Setting of Progression on Targeted Therapy:
 - ▶ For many of the above listed analytes, there is growing recognition of the molecular mechanisms of resistance to therapy. Re-testing of a sample from a tumor that is actively progressing while exposed to targeted therapy can shed light on appropriate next therapeutic steps:
 - ◊ For patients with an underlying *EGFR* sensitizing mutation who have been treated with *EGFR* TKI, minimum appropriate testing includes high-sensitivity evaluation for *p.T790M*; when there is no evidence of *p.T790M*, testing for alternate mechanisms of resistance (*MET* amplification, *ERBB2* amplification) may be used to direct patients for additional therapies. The presence of *p.T790M* can direct patients to third-generation *EGFR* TKI therapy.
 - Assays for the detection of *EGFR p.T790M* should be designed to have an analytic sensitivity of a minimum of 5% allelic fraction. The original sensitizing mutation can be utilized as an internal control in many assays to determine whether a *p.T790M* is within the range of detection if present as a sub-clonal event.
 - ◊ For patients with underlying *ALK* rearrangement who have been treated with *ALK* TKI, it is unclear whether identification of specific tyrosine kinase domain mutation can identify appropriate next steps in therapy, although some preliminary data suggest that specific kinase domain mutations can impact next line of therapy.

Continued

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-H
4 OF 5

EMERGING BIOMARKERS TO IDENTIFY NOVEL THERAPIES FOR PATIENTS WITH METASTATIC NSCLC

Genetic Alteration (ie, Driver event)	Available Targeted Agents with Activity Against Driver Event in Lung Cancer
High-level <i>MET</i> amplification	Crizotinib ¹⁻² Capmatinib ³
<i>ERBB2</i> (<i>HER2</i>) mutations	Ado-trastuzumab emtansine ⁴ Fam-trastuzumab deruxtecan-nxki ⁵

¹ Ou SH, Kwak EL, Siwak-Tapp C, et al. Activity of crizotinib (PF02341066), a dual mesenchymal-epithelial transition (MET) and anaplastic lymphoma kinase (ALK) inhibitor, in a non-small cell lung cancer patient with de novo MET amplification. *J Thorac Oncol* 2011;6:942-946.

² Camidge RD, Ou S-H, Shapiro G, et al. Efficacy and safety of crizotinib in patients with advanced c-MET-amplified non-small cell lung cancer. *J Clin Oncol* 2014;32(Suppl 5):Abstract 8001.

³ Wolf J, Seto T, Han JY, et al; GEOMETRY mono-1 Investigators. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med* 2020;383:944-957.

⁴ Li BT, Shen R, Buonocore D, et al. Ado-trastuzumab emtansine in patients with HER2 mutant lung cancers: Results from a phase II basket trial. *J Clin Oncol* 2018;36:2532-2537.

⁵ Smit EF, Nakagawa K, Nagasaka M, et al. Trastuzumab deruxtecan (T-DXd; DS-8201) in patients with HER2-mutated metastatic non-small cell lung cancer: interim results of DESTINY-Lung01[abstract]. *J Clin Oncol* 2020;38:Abstract 9504.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
The NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN®.

NSCL-I

References

1. Siegel RL, Miller KD, Fuchs HE, et al. Cancer statistics, 2021. *CA Cancer J Clin* 2021;71:7–33.
2. Howlader N, Noone AM, Krapcho M, et al, eds. SEER Cancer Statistics Review, 1975–2017. Accessed January 10, 2021. Available at: https://seer.cancer.gov/csr/1975_2017/
3. Nadal E, Massutí B, Dómine M, et al. Immunotherapy with checkpoint inhibitors in non-small cell lung cancer: insights from long-term survivors. *Cancer Immunol Immunother* 2019;68:341–352.
4. Garon EB, Hellmann MD, Rizvi NA, et al. Five-year overall survival for patients with advanced non-small-cell lung cancer treated with pembrolizumab: results from the phase I KEYNOTE-001 study. *J Clin Oncol* 2019;37:2518–2527.
5. Leigh NB, Hellmann MD, Hui R, et al. Pembrolizumab in patients with advanced non-small-cell lung cancer (KEYNOTE-001): 3-year results from an open-label, phase 1 study. *Lancet Respir Med* 2019;7:347–357.
6. Lin JJ, Cardarella S, Lydon CA, et al. Five-year survival in EGFR-mutant metastatic lung adenocarcinoma treated with EGFR-TKIs. *J Thorac Oncol* 2016;11:556–565.
7. Pacheco JM, Gao D, Smith D, et al. Natural history and factors associated with overall survival in stage IV ALK-rearranged non-small cell lung cancer. *J Thorac Oncol* 2019;14:691–700.
8. Shaw AT, Riely GJ, Bang YJ, et al. Crizotinib in ROS1-rearranged advanced non-small-cell lung cancer (NSCLC): updated results, including overall survival, from PROFILE 1001. *Ann Oncol* 2019;30:1121–1126.
9. Singhi EK, Horn L, Sequist LV, et al. Advanced non-small cell lung cancer: sequencing agents in the EGFR-mutated/ALK-rearranged populations. *Am Soc Clin Oncol Educ Book* 2019;39:e187–e197.
10. Zhao D, Chen X, Qin N, et al. The prognostic role of EGFR-TKIs for patients with advanced non-small cell lung cancer. *Sci Rep* 2017;7:40374.
11. Johung KL, Yeh N, Desai NB, et al. Extended survival and prognostic factors for patients with ALK-rearranged non-small-cell lung cancer and brain metastasis. *J Clin Oncol* 2016;34:123–129.
12. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Updated analysis of KEYNOTE-024: pembrolizumab versus platinum-based chemotherapy for advanced non-small-cell lung cancer with PD-L1 tumor proportion score of 50% or greater. *J Clin Oncol* 2019;37:537–546.
13. Antonia SJ, Borghaei H, Ramalingam SS, et al. Four-year survival with nivolumab in patients with previously treated advanced non-small-cell lung cancer: a pooled analysis. *Lancet Oncol* 2019;20:1395–1408.
14. Ettinger DS, Wood DE, Aisner DL, et al. NCCN Clinical Practice Guidelines in Oncology: Non-Small Cell Lung Cancer. Version 2.2021. Accessed January 10, 2021. To view the most recent version, visit [NCCN.org](https://www.nccn.org)
15. Shaw AT, Bauer TM, de Marinis F, et al. First-line lorlatinib or crizotinib in advanced ALK-positive lung cancer. *N Engl J Med* 2020;383:2018–2029.
16. Wolf J, Seto T, Han JY, et al. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med* 2020;383:944–957.
17. Gainor JF, Curigliano G, Kim DW, et al. Registrational dataset from the phase I/II ARROW trial of pralsetinib (BLU-667) in patients (pts) with advanced RET fusion+ non-small cell lung cancer (NSCLC) [abstract]. *J Clin Oncol* 2020;38(Suppl):Abstract 9515.
18. Drilon A, Oxnard GR, Tan DSW, et al. Efficacy of selpercatinib in RET fusion-positive non-small-cell lung cancer. *N Engl J Med* 2020;383:813–824.
19. Smit EF, Nakagawa K, Nagasaka M, et al. Trastuzumab deruxtecan (T-DXd; DS-8201) in patients with HER2-mutated metastatic non-small cell lung cancer (NSCLC): interim results of DESTINY-Lung01 [abstract]. *J Clin Oncol* 2020;38(Suppl):Abstract 9504.
20. Tsurutani J, Iwata H, Krop I, et al. Targeting HER2 with trastuzumab deruxtecan: a dose-expansion, phase I study in multiple advanced solid tumors. *Cancer Discov* 2020;10:688–701.
21. Wu YL, Tsuboi M, He J, et al. Osimertinib in resected EGFR-mutated non-small-cell lung cancer. *N Engl J Med* 2020;383:1711–1723.
22. Benayed R, Offin M, Mullaney K, et al. High yield of RNA sequencing for targetable kinase fusions in lung adenocarcinomas with no mitogenic driver alteration detected by DNA sequencing and low tumor mutation burden. *Clin Cancer Res* 2019;25:4712–4722.
23. Mazieres J, Drilon A, Lusque A, et al. Immune checkpoint inhibitors for patients with advanced lung cancer and oncogenic driver alterations: results from the IMMUNOTARGET registry. *Ann Oncol* 2019;30:1321–1328.
24. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med* 2018;378:113–125.
25. Lisberg A, Cummings A, Goldman JW, et al. A phase II study of pembrolizumab in EGFR-mutant, PD-L1+, tyrosine kinase inhibitor naive patients with advanced NSCLC. *J Thorac Oncol* 2018;13:1138–1145.
26. Camidge DR, Dziadziuszko R, Peters S, et al. Updated efficacy and safety data and impact of the EML4-ALK fusion variant on the efficacy of alectinib in untreated ALK-positive advanced non-small cell lung cancer in the global phase III ALEX study. *J Thorac Oncol* 2019;14:1233–1243.
27. Gainor JF, Shaw AT, Sequist LV, et al. EGFR mutations and ALK rearrangements are associated with low response rates to PD-1 pathway blockade in non-small cell lung cancer: a retrospective analysis. *Clin Cancer Res* 2016;22:4585–4593.
28. Tsao MS, Kerr KM, Kockx M, et al. PD-L1 immunohistochemistry comparability study in real-life clinical samples: results of Blueprint Phase 2 Project. *J Thorac Oncol* 2018;13:1302–1311.
29. Hirsch FR, McElhinny A, Stanforth D, et al. PD-L1 immunohistochemistry assays for lung cancer: results from phase 1 of the Blueprint PD-L1 IHC Assay Comparison Project. *J Thorac Oncol* 2017;12:208–222.
30. Lam VK, Tran HT, Banks KC, et al. Targeted tissue and cell-free tumor DNA sequencing of advanced lung squamous-cell carcinoma reveals clinically significant prevalence of actionable alterations. *Clin Lung Cancer* 2019;20:30–36.e3.
31. Sands JM, Nguyen T, Shivdasani P, et al. Next-generation sequencing informs diagnosis and identifies unexpected therapeutic targets in lung squamous cell carcinomas. *Lung Cancer* 2020;140:35–41.
32. Sholl LM, Aisner DL, Varella-Garcia M, et al. Multi-institutional oncogenic driver mutation analysis in lung adenocarcinoma: the Lung Cancer Mutation Consortium experience. *J Thorac Oncol* 2015;10:768–777.
33. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in advanced ALK inhibitor-naïve ALK-positive non-small cell lung cancer: second interim analysis of the phase III ALTA-1L trial. *J Clin Oncol* 2020;38:3592–3603.
34. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in ALK-positive non-small-cell lung cancer. *N Engl J Med* 2018;379:2027–2039.
35. D'Angelo A, Sobhani N, Chapman R, et al. Focus on ROS1-positive non-small cell lung cancer (NSCLC): crizotinib, resistance mechanisms and the newer generation of targeted therapies. *Cancers (Basel)* 2020;12:12.
36. Drilon A, Siena S, Dziadziuszko R, et al. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:261–270.
37. Mu Y, Yang K, Hao X, et al. Clinical characteristics and treatment outcomes of 65 patients with BRAF-mutated non-small cell lung cancer. *Front Oncol* 2020;10:603.
38. Auliac JB, Bayle S, Do P, et al. Efficacy of dabrafenib plus trametinib combination in patients with BRAF V600E-mutant NSCLC in real-world setting: GFPC 01-2019. *Cancers (Basel)* 2020;12:12.
39. Khunger A, Khunger M, Velcheti V. Dabrafenib in combination with trametinib in the treatment of patients with BRAF V600-positive advanced or metastatic non-small cell lung cancer: clinical evidence and experience. *Ther Adv Respir Dis* 2018;12:1753466618767611.
40. Reck M, Rodríguez-Abreu D, Robinson AG, et al. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* 2016;375:1823–1833.
41. Mok TSK, Wu YL, Kudaba I, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819–1830.
42. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden. *N Engl J Med* 2018;378:2093–2104.
43. Hellmann MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. *N Engl J Med* 2019;381:2020–2031.
44. Gandhi L, Rodríguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378:2078–2092.
45. Herbst RS, Giaccone G, de Marinis F, et al. Atezolizumab for first-line treatment of PD-L1-selected patients with NSCLC. *N Engl J Med* 2020;383:1328–1339.
46. Socinski MA, Jotte RM, Cappuzzo F, et al. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N Engl J Med* 2018;378:2288–2301.
47. Paz-Ares L, Luft A, Vicente D, et al. Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. *N Engl J Med* 2018;379:2040–2051.

48. Sesma A, Pardo J, Cruellas M, et al. From tumor mutational burden to blood T cell receptor: looking for the best predictive biomarker in lung cancer treated with immunotherapy. *Cancers (Basel)* 2020;12:12.
49. Borghaei H, Langer CJ, Paz-Ares L, et al. Pembrolizumab plus chemotherapy versus chemotherapy alone in patients with advanced non-small cell lung cancer without tumor PD-L1 expression: a pooled analysis of 3 randomized controlled trials. *Cancer* 2020;126:4867–4877.
50. Sholl LM, Hirsch FR, Hwang D, et al. The promises and challenges of tumor mutation burden as an immunotherapy biomarker: a perspective from the International Association for the Study of Lung Cancer Pathology Committee. *J Thorac Oncol* 2020;15:1409–1424.
51. Offin M, Rizvi H, Tenet M, et al. Tumor mutation burden and efficacy of EGFR-tyrosine kinase inhibitors in patients with EGFR-mutant lung cancers. *Clin Cancer Res* 2019;25:1063–1069.
52. Rizvi H, Sanchez-Vega F, La K, et al. Molecular determinants of response to anti-programmed cell death (PD)-1 and anti-programmed death-ligand 1 (PD-L1) blockade in patients with non-small-cell lung cancer profiled with targeted next-generation sequencing. *J Clin Oncol* 2018;36:633–641.
53. Carbone DP, Reck M, Paz-Ares L, et al. First-line nivolumab in stage IV or recurrent non-small-cell lung cancer. *N Engl J Med* 2017;376:2415–2426.
54. Marabelle A, Fakih M, Lopez J, et al. Association of tumour mutational burden with outcomes in patients with advanced solid tumours treated with pembrolizumab: prospective biomarker analysis of the multicohort, open-label, phase 2 KEYNOTE-158 study. *Lancet Oncol* 2020;21:1353–1365.
55. Antonia SJ, Villegas A, Daniel D, et al. Overall survival with durvalumab after chemoradiotherapy in stage III NSCLC. *N Engl J Med* 2018;379:2342–2350.