NCCN: Continuing Education

Target Audience: This activity is designed to meet the educational needs of oncologists, nurses, pharmacists, and other healthcare professionals who manage patients with cancer.

Accreditation Statements

In support of improving patient care, National Comprehensive Cancer Network (NCCN) is jointly accredited by the Accreditation Council for Continuing Medical Education (ACCME), the Accreditation Council for Pharmacy Education (ACPE), and the American Nurses Credentialing Center (ANCC), to provide continuing education for the healthcare team.

Medicine (ACCME): NCCN designates this journal-based CME activity for a maximum of 1.0 AMA PRA Category 1 Credit™. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Nursing (ANCC): NCCN designates this educational activity for a maximum of 1.0 contact hour.

Pharmacy (ACPE): NCCN designates this knowledge-based continuing education activity for 1.0 contact hour (0.1 CEUs) of continuing education credit. UAN: JA4008196-0000-21-010-H01-P

All clinicians completing this activity will be issued a certificate of participation. To participate in this journal CE activity: (1) review the educational content; (2) take the posttest with a 66% minimum passing score and complete the evaluation at <https://education.nccn.org/node/90502>; and (3) view/print certificate.

Pharmacists: You must complete the posttest and evaluation within 30 days of the activity. Continuing pharmacy education credit is reported to the CPE Monitor once you have completed the posttest and evaluation and claimed your credits. Before completing these requirements, be sure your NCCN profile has been updated with your NAPB e-profile ID and date of birth. Your credit cannot be reported without this information. If you have any questions, please e-mail education@nccn.org.

Release date: July 10, 2021; Expiration date: July 10, 2022

Learning Objectives:

Upon completion of this activity, participants will be able to:

- Integrate into professional practice the updates to the NCCN Guidelines for Palliative Care
- Describe the rationale behind the decision-making process for developing the NCCN Guidelines for Palliative Care

Disclosure of Relevant Financial Relationships

The NCCN staff listed below discloses no relevant financial relationships:

Kerrin M. Rosenthal, MA; Kimberly Callan, MS; Genevieve Emberger Hartzman, MA; Erin Hesler; Kristina M. Gregory, RN, MSN, OCN; Rashmi Kumar, PhD; Karen Kanefield; and Kathy Smith.

Individuals Who Provided Content Development and/or Authorship Assistance:

Maria Dans, MD, Panel Chair, has disclosed no relevant financial relationships.

Jean S. Kutner, MD, MSPH, Panel Vice Chair, has disclosed no relevant financial relationships.

Mary Anne Bergman, Guidelines Coordinator, NCCN, has disclosed no relevant financial relationships.

Ndiya Ogbu, PhD, Oncology Scientist/Medical Writer, NCCN, has disclosed no relevant financial relationships.

Mallory Campbell, PhD, Oncology Scientist/Medical Writer, NCCN, has disclosed no relevant financial relationships.

To view all of the conflicts of interest for the NCCN Guidelines Panel, go to [NCCN.org/guidelines/guidelines-panels-and-disclosure/disclosure-panels](https://www.nccn.org/guidelines/guidelines-panels-and-disclosure/disclosure-panels).

This activity is supported by educational grants from Agios Pharmaceuticals; AstraZeneca; Clovis Oncology, Inc.; Daiichi Sankyo; Eisai; Epizyme Inc.; Novartis; and Pharmacyclis LLC, an AbbVie Company and Janssen Biotech, Inc., administered by Janssen Scientific Affairs, LLC. This activity is supported by an independent medical education grant from Bristol-Myers Squibb, and Regeneron Pharmaceuticals, Inc. and Sanofi Genzyme. This activity is supported by an independent medical educational grant from Mylan Inc. This activity is supported by a medical education grant from Karyopharm Therapeutics. This activity is supported by an independent educational grant from AbbVie.

Palliative Care, Version 2.2021

Featured Updates to the NCCN Guidelines

Maria Dans, MD^{1,*}; Jean S. Kutner, MD, MSPH^{2,*}; Rajiv Agarwal, MD³; Justin N. Baker, MD⁴; Jessica R. Bauman, MD⁵; Anna C. Beck, MD⁶; Toby C. Campbell, MD, MSCI⁷; Elise C. Carey, MD⁸; Amy A. Case, MD⁹; Shalini Dalal, MD¹⁰; Danielle J. Doberman, MD, MPH¹¹; Andrew S. Epstein, MD¹²; Leslie Fecher, MD¹³; Joshua Jones, MD, MA¹⁴; Jennifer Kapo, MD¹⁵; Richard T. Lee, MD¹⁶; Elizabeth T. Loggers, MD, PhD¹⁷; Susan McCammon, MD¹⁸; William Mitchell, MD¹⁹; Adeboye B. Ogunseitan, MD²⁰; Diane G. Portman, MD²¹; Kavitha Ramchandran, MD²²; Linda Sutton, MD²³; Jennifer Temel, MD²⁴; Melissa L. Teply, MD²⁵; Stephanie Y. Terauchi, MD²⁶; Jane Thomas, MD²⁷; Anne M. Walling, MD, PhD²⁸; Finly Zachariah, MD²⁹; Mary Anne Bergman^{30,*}; Ndiya Ogbia, PhD^{30,*}; and Mallory Campbell, PhD^{30,*}

ABSTRACT

Palliative care has evolved to be an integral part of comprehensive cancer care with the goal of early intervention to improve quality of life and patient outcomes. The NCCN Guidelines for Palliative Care provide recommendations to help the primary oncology team promote the best quality of life possible throughout the illness trajectory for each patient with cancer. The NCCN Palliative Care Panel meets annually to evaluate and update recommendations based on panel members' clinical expertise and emerging scientific data. These NCCN Guidelines Insights summarize the panel's recent discussions and highlights updates on the importance of fostering adaptive coping strategies for patients and families, and on the role of pharmacologic and non-pharmacologic interventions to optimize symptom management.

J Natl Compr Canc Netw 2021;19(7):780–788
doi: 10.6004/jnccn.2021.0033

NCCN CATEGORIES OF EVIDENCE AND CONSENSUS

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Clinical trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

PLEASE NOTE

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. **The NCCN Guidelines Insights highlight important changes in the NCCN Guidelines recommendations from previous versions. Colored markings in the algorithm show changes and the discussion aims to further the understanding of these changes by summarizing salient portions of the panel's discussion, including the literature reviewed.**

The NCCN Guidelines Insights do not represent the full NCCN Guidelines; further, the National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use, or application of the NCCN Guidelines and NCCN Guidelines Insights and disclaims any responsibility for their application or use in any way.

The complete and most recent version of these NCCN Guidelines is available free of charge at [NCCN.org](https://www.nccn.org).

© National Comprehensive Cancer Network, Inc. 2021. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN.

¹Siteman Cancer Center at Barnes-Jewish Hospital and Washington University School of Medicine; ²University of Colorado Cancer Center; ³Vanderbilt-Ingram Cancer Center; ⁴St. Jude Children's Research Hospital/The University of Tennessee Health Science Center; ⁵Fox Chase Cancer Center; ⁶Huntsman Cancer Institute at the University of Utah; ⁷University of Wisconsin Carbone Cancer Center; ⁸Mayo Clinic Cancer Center; ⁹Roswell Park Comprehensive Cancer Center; ¹⁰The University of Texas MD Anderson Cancer Center; ¹¹The Sidney Kimmel Comprehensive Cancer Center at Johns Hopkins; ¹²Memorial Sloan Kettering Cancer Center; ¹³University of Michigan Rogel Cancer Center; ¹⁴Abramson Cancer Center at the University of Pennsylvania; ¹⁵Yale Cancer Center/Smilow Cancer Hospital; ¹⁶Case Comprehensive Cancer Center/University Hospitals Seidman Cancer Center and Cleveland Clinic Taussig Cancer Institute; ¹⁷Fred Hutchinson Cancer Research Center/Seattle Cancer Care Alliance; ¹⁸O'Neal Comprehensive Cancer Center at UAB; ¹⁹UC San Diego Moores Cancer Center; ²⁰Robert H. Lurie Comprehensive Cancer Center of Northwestern University; ²¹Moffitt Cancer Center; ²²Stanford Cancer Institute; ²³Duke Cancer Institute; ²⁴Massachusetts General Hospital Cancer Center; ²⁵Fred & Pamela Buffett Cancer Center; ²⁶UT Southwestern Simmons Comprehensive Cancer Center; ²⁷Dana-Farber Cancer Institute; ²⁸UCLA Jonsson Comprehensive Cancer Center; ²⁹City of Hope National Medical Center; and ³⁰National Comprehensive Cancer Network.

* Provided content development and/or authorship assistance.

Definition of Palliative Care^{a,b}

Palliative care is an approach to patient/family/caregiver-centered health care that focuses on optimal management of distressing symptoms, while incorporating psychosocial and spiritual care according to patient/family/caregiver needs, values, beliefs, and cultures. The goal of palliative care is to anticipate, prevent, and reduce suffering; **promote adaptive coping**; and support the best possible quality of life for patients/families/caregivers, regardless of the stage of the disease or the need for other therapies. Palliative care can begin at diagnosis; be delivered concurrently with disease-directed, life-prolonging therapies; and facilitate patient autonomy, access to information, and choice. Palliative care becomes the main focus of care when disease-directed, life-prolonging therapies are no longer effective, appropriate, or desired. Palliative care should be provided by the primary oncology team and augmented as needed by collaboration with an interprofessional team of palliative care experts.

Standards of Palliative Care^{c,d}

- Institutions should develop processes for integrating palliative care into cancer care, both as part of usual oncology care and for patients with specialty palliative care needs.
- All cancer patients should be screened for palliative care needs at their initial visit, at appropriate intervals, and as clinically indicated.
- Patients/families/caregivers should be informed that palliative care is an integral part of their comprehensive cancer care.
- Educational programs should be provided to all health care professionals and trainees so that they can develop effective palliative care knowledge, skills, and attitudes.
- Palliative care specialists and **interprofessional palliative care teams**, including but not limited to board-certified palliative care physicians, advanced practice nurses, physician assistants, nurses, dietitians, social workers, chaplains, and pharmacists, should be readily available to provide consultative or direct care to patients/families/caregivers and/or health care professionals who request or require their expertise.
- Quality of palliative care should be monitored by institutional quality improvement programs.

^aHui D, et al. *J Pain Symptom Manage* 2012;43:582-592.

^bSeaman JB, et al. *J Palliat Med* 2020;23:1157-1158.

^cFerris FD, et al. *J Clin Oncol* 2009;27:3052-3058.

^dIOM (Institute of Medicine). 2014 *Dying in America: Improving quality and honoring individual preferences near the end of life*. Washington, DC: The National Academies Press. Available at: www.nap.edu/read/18748/chapter/1.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.

NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

Overview

More than 1.8 million people are expected to be diagnosed with cancer in the United States in 2021, and 608,570 people are expected to die of the disease.¹ Global cancer rates are increasing, with an associated rise in the number of cancer survivors living with symptoms and disabilities as a result of their disease and/or its treatment.¹⁻³ More than one-third of patients with cancer in a large observational cohort study reported moderate to severe symptoms in most categories (pain, nausea, anxiety, depression, shortness of breath, drowsiness, well-being, loss of appetite, and tiredness) in the last weeks of life.⁴ Several groups, including the WHO, ASCO, and American Society of Hematology, recommend integrating palliative care with disease-directed care beginning at the time of diagnosis.⁵⁻⁷ This allows for more holistic and timely identification of challenges facing patients and their families, and more robust and restorative problem-solving.

Palliative Care in Oncology

The NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) define palliative care as an approach to patient/family/caregiver-centered healthcare that

focuses on optimal management of distressing symptoms, while incorporating psychosocial and spiritual care according to patient/family/caregiver needs, values, beliefs, and cultures. The goal of palliative care is to anticipate, prevent, and reduce suffering; **promote adaptive coping**; and support the best possible quality of life (QoL) for patients/families/caregivers, regardless of the stage of the disease or the need for other therapies. Palliative care should begin at diagnosis; be delivered concurrently with disease-directed, life-prolonging therapies; and facilitate patient autonomy, access to information, and choice. The value of early integration of palliative care into the continuum of cancer care has been well demonstrated.⁸⁻¹² During the recent meeting to discuss updates to the NCCN Guidelines for Palliative Care, the panel highlighted the importance of applying palliative care practices in a way that includes not only the patient but also families and caregivers. In particular, the panel stressed that the promotion of adaptive coping strategies (defined as problem-solving strategies as opposed to avoidant coping strategies) for patients, families, and caregivers should be a component of all stages of the care plan. A recent study examining patients' perception of treatment goals showed that patients who acknowledged the incurable nature of their

⁹See FEV-D in the NCCN Guidelines for Prevention and Treatment of Cancer-Related Infections for an approach to assess for the risk of serious complications in patients with neutropenic fever.

¹Look for opportunities to use single agents to treat multiple symptoms.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

PAL-5

disease reported higher psychologic distress compared with those who perceived their treatment goals as curative, regardless of accuracy of the perception.¹³ Additionally, patients whose treatment goals were discordant with their oncologist's also reported increased distress symptoms.¹³ In contrast, another study showed that patients with an accurate understanding that their disease was incurable who engaged in specific active coping strategies had fewer depressive symptoms than those not engaged in coping strategies.¹⁴ The panel recommends encouraging adaptive coping methods for patients, family, and caregivers, particularly when cultivating prognostic awareness to help improve both goal-concordant care and QoL.

Palliative care should be initiated by the primary oncology team and should be augmented through collaboration with palliative care specialists in an interprofessional team (including but not limited to board-certified palliative care physicians, advanced practice providers, nurses, dietitians, social workers, psychologists, psychiatrists, chaplains, and pharmacists) to address intractable symptoms and/or complex psychosocial issues.¹⁵ Additionally, palliative care should reach beyond the patient

to include family and caregivers. During the recent meeting to discuss updates to the NCCN Guidelines, the panel acknowledged that palliative care is a collaborative practice, and therefore recommends recognition of all expertise with the term "interprofessional." Using language that allows for a better understanding of expectations can improve teamwork during interactions among multiple areas of practice.¹⁵

Comprehensive Assessment

Patients should undergo a comprehensive palliative care assessment by their primary oncology team, including an evaluation of the benefits and burdens of anticancer therapy; the patient's and family's decision-making style and capacity; their coping strategies; any physical symptoms; psychosocial or spiritual distress; personal goals, values, and expectations; educational and informational needs; potential financial toxicity; and cultural factors affecting care.^{16–18}

Assessment of Psychosocial Distress

Assessment of psychosocial distress should include both the patient and the caregivers, and should focus on

PAL-9

illness-related distress and psychosocial, spiritual, or existential needs according to the NCCN Guidelines for Distress Management (available at NCCN.org). Concerns regarding social support and resources (ie, home, family, community, or financial) must also be assessed. While discussing recent updates to the Guidelines, the panel emphasized the effect that employment or the loss of employment may have on psychologic distress. Loss of employment, in particular, may lead to decreased feelings of personal purpose, increased social isolation (or loneliness) due to loss of one's workplace community, as well as financial toxicity.¹⁹ Interventions that may help to reverse some of this distress include but are not limited to promotion of adaptive coping methods for both patients and caregivers, and provision of additional psychologic and financial support, when possible (see PAL-5, page 783).

Palliative Care Interventions

The oncology team should initiate treatments following the specific recommendations described in these guidelines for common symptoms. Comorbid physical and

psychosocial conditions should be treated by appropriate clinicians. Consultation or collaboration with palliative care specialists or teams is recommended for patients with more complex concerns to improve their QoL.^{20–23} Referrals should be made as needed to mental health, social and psychosocial support services, spiritual care, healthcare interpreters, hospice services, financial counselors, or other specialists. Finally, the oncology team may find collaboration with a palliative care team helpful in identifying additional support through religious organizations, schools, or other agencies in the community.

Anticancer Therapy

Primary palliative care, including both anticancer treatments and disease-related symptom management, should be provided for all patients. Anticancer therapy should be in line with standard practice as well as stated patient goals and priorities and be accompanied by appropriate prevention/management of adverse effects as well as palliation of physical and psychosocial symptoms. A longitudinal cohort study evaluated the impact of palliative chemotherapy on QoL near the end of life (median 3.8 months before

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

PAL-10

death) as a function of patient performance status. For patients with progressive metastatic cancer in this study, palliative chemotherapy failed to improve QoL near death for those with moderate or poor performance status and worsened QoL near death for those with initially good performance status.²⁴ Additional studies have reported that anticancer therapy may continue beyond what is evidence-based for patients with advanced or metastatic disease.^{25–27} This suggests a need for better prognostic awareness and iterative discussions of patient/family/caregiver values and goals of treatment.

Symptom Management

Symptoms should be comprehensively assessed in all patients. In noncommunicative patients, symptom intensity may be assessed using physical signs. Education should be provided to the patient/family/caregiver on the patient's condition and the risk/benefit of treatment options. At the most recent meeting to update the NCCN Guidelines, the panel added a recommendation that providing resources for the psychosocial and emotional support of not only the patient but also family/caregiver(s) be considered an integral part of symptom management. As a general

principle, when appropriate, providers should try to use palliative interventions that may address multiple symptoms.

Pain

The patient and family/caregiver should be educated on the role of pain medications, customized to the patient's particular situation. In addition, it is important to note that dying patients in their last weeks of life may have different needs than patients at earlier disease stages. For instance, if feasible or desired, pain management may need to be balanced against reduced level of consciousness. In advanced cancer, opioids may need to be titrated aggressively for moderate/severe acute/chronic pain.²⁸ Single-fraction palliative radiotherapy may also be used to address pain associated with bone metastases.^{29–32} A recent systematic review also found that terminally ill patients receiving nonpharmacologic therapy in addition to medication experienced greater physical symptom relief and overall mood improvement.³³ The panel recommends consideration of nonpharmacologic and/or integrative interventions as an additional source of pain management. Examples include cognitive behavioral therapy and acceptance and commitment therapy; massage; art or music

^kSee Drug Appendix (PAL-A) for specific recommendations for medical management of symptoms.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

PAL-13

therapy; application of hot or cold packs; repositioning; and use of supportive braces.^{31–39} For pain refractory to all of these interventions, palliative sedation may be considered following consultation with pain management or palliative care specialists, consistent with local institutional policies.

Anorexia/Cachexia

Cachexia is physical wasting with loss of skeletal and visceral muscle mass and is very common among patients with advanced cancer.^{40,41} Many patients with cancer lose the desire to eat (anorexia), which contributes to cachexia. Cachexia can also occur independently from anorexia, as proinflammatory cytokines and tumor-derived factors directly lead to muscle proteolysis.^{40,41} Cachexia leads to asthenia (weakness), hypoalbuminemia, emaciation, immune system impairment, metabolic dysfunction, and autonomic failure. Cancer-related cachexia has also been associated with failure of anticancer treatment, increased treatment toxicity, delayed treatment initiation, early treatment termination, shorter survival, and psychosocial distress.^{40–42} A study that examined cancer cachexia in a cohort of 1,473 patients across all weight ranges showed that muscle depletion conveys a similarly

poor prognosis as involuntary weight loss, regardless of body mass index.⁴³

Initial steps include evaluating the rate and severity of weight loss and associated symptoms, educating the patient/family/caregiver on the patient's condition, and discussing the risks/benefits of available treatment options. During assessment, the panel strongly recommends considering a formal swallowing evaluation for patients who report dysphagia or exhibit symptoms that suggest possible swallowing dysfunction. The panel also suggests that clinicians assess for conditions or symptoms that interfere with food intake, such as dysgeusia, xerostomia, dental/jaw issues, intractable hiccups, mucositis, oropharyngeal candidiasis, depression, early satiety, nausea and vomiting, and pain. A psychiatric evaluation may be considered if there is suspicion or a history of an eating disorder. Treatment may include management of symptoms or discontinuation of medications that interfere with food intake, as well as improving a bowel regimen to optimize gut motility.^{44–46} Strengthening social supports, dietary counseling, and feeding assistance may also be offered as nonpharmacologic management of cachexia.⁴⁷

If increased appetite is an important aspect of the patient's QoL, appetite stimulants may be helpful (eg,

PALLIATIVE CARE DRUG APPENDIX

Condition	Recommended Agents and Dosage by Estimated Life Expectancy and Symptom Etiology
Dyspnea (PAL-11)	Life Expectancy: Years; Year to Months; and Months to Weeks • General: Morphine, 2.5–10 mg PO q2h PRN or 1–3 mg IV q2h PRN for opioid naïve, increase dose by 25% for opioid tolerant ▶ For acute progressive dyspnea, or for patients who are not opioid naïve, more aggressive titration may be required • Anxiety: Lorazepam, 0.25–1 mg PO q4h PRN for benzodiazepine naïve
Dyspnea (PAL-12)	Life Expectancy: Weeks to Days (dying patient) • General: Morphine, 2.5–10 mg PO q2h PRN or 1–3 mg IV q2h PRN if opioid naïve, increase dose by 25% for opioid tolerant ▶ For acute progressive dyspnea, or for patients who are not opioid naïve, more aggressive titration may be required • Anxiety: Lorazepam, 0.25–1 mg PO q4h PRN if benzodiazepine naïve • Fluid overload: Furosemide
Secretions (PAL-12)	• Excessive secretions: Scopolamine, 0.4 mg SC q4h PRN/1.5 mg patches, 1–3 patches q72h OR atropine, 1% ophthalmic solution 1–2 drops SL q4h PRN OR glycopyrrolate, 0.2–0.4 mg IV or SC q4h PRN
Anorexia/Cachexia ¹ (PAL-13)	Life Expectancy: Years; Year to Months • Depression/anorexia: Mirtazapine, 7.5–30 mg PO QHS • Gastroparesis (early satiety): Metoclopramide 5–10 mg PO QID 30 min before meals and at bedtime • Low/no appetite: Megestrol acetate, 200–600 mg/d PO OR olanzapine, 5 mg/d PO
Anorexia/Cachexia ¹ (PAL-14)	Life Expectancy: Months to Weeks; Weeks to Days (dying patient) • Low/no appetite: Megestrol acetate, 400–800 mg/d PO OR olanzapine, 5 mg/d PO OR dexamethasone, 3–4 mg/d PO OR consider cannabinoid • Depression: Mirtazapine, 7.5–30 mg PO QHS

¹Roeland EJ, Bohlke K, Baracos VE, et al. Management of Cancer Cachexia; ASCO Guideline. J Clin Oncol 2020;38(21):2438-2453.

Version 2.2021 © National Comprehensive Cancer Network, Inc. 2021. All rights reserved.
NCCN Guidelines® and this illustration may not be reproduced in any form without the express written permission of NCCN.

PAL-A
1 OF 5

megestrol acetate, dexamethasone, olanzapine).^{48–52} Systematic reviews and meta-analyses of megestrol acetate demonstrate improved appetite and slight improvements in weight gain.^{47,49} However, benefits of using megestrol acetate should be balanced with the potential increased risk for venous thromboembolic events, fluid retention, and death, particularly in older patients or those who are nonambulatory.^{53–55} Combining megestrol acetate therapy with dietary counseling and nutritional supplements may yield improved outcomes for patients with preserved functional status and cancer cachexia.⁴⁷

Summary

This article highlights recent updates to the 2021 version of the NCCN Guidelines for Palliative Care. Several

recommendations were made by the panel to enhance the QoL of patient/family/caregivers throughout the disease trajectory. These include effectively assessing and addressing psychosocial distress, promoting adaptive coping strategies for patients/family/caregivers, and emphasizing the need for holistic cancer care plans that include patient/family/caregiver education and support. The panel acknowledged and recommended describing the palliative care team as “interprofessional” to emphasize its collaborative nature. Updates also included suggestions for a multimodal approach to the management of anorexia/cachexia.

To participate in this journal CE activity, go to <https://education.nccn.org/node/90502>

References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. CA Cancer J Clin 2020;70:7–30.
- Centers for Disease Control and Prevention. Cancer survivors—United States, 2007. MMWR Morb Mortal Wkly Rep 2011;60:269–272.
- Ferris FD, Bruera E, Cherny N, et al. Palliative cancer care a decade later: accomplishments, the need, next steps – from the American Society of Clinical Oncology. J Clin Oncol 2009;27:3052–3058.
- Seow H, Barbera L, Sutradhar R, et al. Trajectory of performance status and symptom scores for patients with cancer during the last six months of life. J Clin Oncol 2011;29:1151–1158.
- Murray SA, Kendall M, Mitchell G, et al. Palliative care from diagnosis to death. BMJ 2017;356:j878.
- Ferrell BR, Temel JS, Temin S, et al. Integration of palliative care into standard oncology care: American Society of Clinical Oncology clinical practice guideline update. J Clin Oncol 2017;35:96–112.

7. LeBlanc TW, El-Jawahri A. When and why should patients with hematologic malignancies see a palliative care specialist? *Hematology (Am Soc Hematol Educ Program)* 2015;2015:471–478.
8. The SUPPORT Principal Investigators. A controlled trial to improve care for seriously ill hospitalized patients. The study to understand prognoses and preferences for outcomes and risks of treatments (SUPPORT). *JAMA* 1995; 274:1591–1598.
9. Malin JL. Bridging the divide: integrating cancer-directed therapy and palliative care. *J Clin Oncol* 2004;22:3438–3440.
10. Meyers FJ, Linder J. Simultaneous care: disease treatment and palliative care throughout illness. *J Clin Oncol* 2003;21:1412–1415.
11. Greer JA, Jackson VA, Meier DE, et al. Early integration of palliative care services with standard oncology care for patients with advanced cancer. *CA Cancer J Clin* 2013;63:349–363.
12. Davis MP, Bruera E, Morganstern D. Early integration of palliative and supportive care in the cancer continuum: challenges and opportunities. *Am Soc Clin Oncol Educ Book* 2013;33:144–150.
13. El-Jawahri A, Forst D, Fenech A, et al. Relationship between perceptions of treatment goals and psychological distress in patients with advanced cancer. *J Natl Compr Canc Netw* 2020;18:849–855.
14. Nipp RD, Greer JA, El-Jawahri A, et al. Coping and prognostic awareness in patients with advanced cancer. *J Clin Oncol* 2017;35:2551–2557.
15. Seaman JB, Lakin JR, Anderson E, et al. Interdisciplinary or interprofessional: why terminology in teamwork matters to hospice and palliative care. *J Palliat Med* 2020;23:1157–1158.
16. Berger A, Portenoy R, Weissman D, eds. *Practice and Principles of Supportive Oncology*. Philadelphia, PA: Lippincott-Raven; 1998.
17. Doyle D, Hanks GWC, MacDonald N, eds. *Oxford Textbook of Palliative Medicine*, 2nd ed. Oxford, England: Oxford University; 1998.
18. Emanuel LL, von Gunten CF, Ferris FD, eds. *The Education for Physicians on End-of-Life Care (EPEC) Curriculum*. Princeton, NJ: The Robert Wood Johnson Foundation; 1999.
19. Carrera PM, Kantarjian HM, Blinder VS. The financial burden and distress of patients with cancer: Understanding and stepping-up action on the financial toxicity of cancer treatment. *CA Cancer J Clin* 2018;68:153–165.
20. Smith TJ, Temin S, Alesi ER, et al. American Society of Clinical Oncology provisional clinical opinion: the integration of palliative care into standard oncology care. *J Clin Oncol* 2012;30:880–887.
21. Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010;363:733–742.
22. Bakitas MA, Tosteson TD, Li Z, et al. Early versus delayed initiation of concurrent palliative oncology care: patient outcomes in the ENABLE III randomized controlled trial. *J Clin Oncol* 2015;33:1438–1445.
23. Hamano J, Yamaguchi T, Maeda I, et al. Multicenter cohort study on the survival time of cancer patients dying at home or in a hospital: does place matter? *Cancer* 2016;122:1453–1460.
24. Prigerson HG, Bao Y, Shah MA, et al. Chemotherapy use, performance status, and quality of life at the end of life. *JAMA Oncol* 2015;1:778–784.
25. Chen AB, Cronin A, Weeks JC, et al. Palliative radiation therapy practice in patients with metastatic non-small-cell lung cancer: a Cancer Care Outcomes Research and Surveillance Consortium (CanCORS) study. *J Clin Oncol* 2013;31:558–564.
26. Zafar SY, Malin JL, Grambow SC, et al. Chemotherapy use and patient treatment preferences in advanced colorectal cancer: a prospective cohort study. *Cancer* 2013;119:854–862.
27. Sano M, Fushimi K. Association of palliative care consultation with reducing inpatient chemotherapy use in elderly patients with cancer in Japan: analysis using a nationwide administrative database. *Am J Hosp Palliat Care* 2017;34:685–691.
28. Ferrell B, Levy MH, Paice J. Managing pain from advanced cancer in the palliative care setting. *Clin J Oncol Nurs* 2008;12:575–581.
29. Lutz S, Balboni T, Jones J, et al. Palliative radiation therapy for bone metastases: update of an ASTRO evidence-based guideline. *Pract Radiat Oncol* 2017;7:4–12.
30. Westhoff PG, de Graeff A, Monnikhof EM, et al. Quality of life in relation to pain response to radiation therapy for painful bone metastases. *Int J Radiat Oncol Biol Phys* 2015;93:694–701.
31. McDonald R, Ding K, Brundage M, et al. Effect of radiotherapy on painful bone metastases: a secondary analysis of the NCIC Clinical Trials Group Symptom Control Trial SC.23. *JAMA Oncol* 2017;3:953–959.
32. Park KR, Lee CG, Tseng YD, et al. Palliative radiation therapy in the last 30 days of life: a systematic review. *Radiother Oncol* 2017;125:193–199.
33. Qi He Mabel L, Drury VB, Hong PW. The experience and expectations of terminally ill patients receiving music therapy in the palliative setting: a systematic review. *JBI Library Syst Rev* 2010;8:1088–1111.
34. Gutsell KJ, Schluchter M, Margevicius S, et al. Music therapy reduces pain in palliative care patients: a randomized controlled trial. *J Pain Symptom Manage* 2013;45:822–831.
35. Kutner JS, Smith MC, Corbin L, et al. Massage therapy versus simple touch to improve pain and mood in patients with advanced cancer: a randomized trial. *Ann Intern Med* 2008;149:369–379.
36. Lim JA, Choi SH, Lee WJ, et al. Cognitive-behavioral therapy for patients with chronic pain: Implications of gender differences in empathy. *Medicine (Baltimore)* 2018;97:e10867.
37. Feliu-Soler A, Montesinos F, Gutiérrez-Martínez O, et al. Current status of acceptance and commitment therapy for chronic pain: a narrative review. *J Pain Res* 2018;11:2145–2159.
38. Syrjala KL, Jensen MP, Mendoza ME, et al. Psychological and behavioral approaches to cancer pain management. *J Clin Oncol* 2014;32:1703–1711.
39. Coelho A, Parola V, Cardoso D, et al. Use of non-pharmacological interventions for comforting patients in palliative care: a scoping review. *JBI Database Syst Rev Implement Reports* 2017;15:1867–1904.
40. Kumar NB, Kazi A, Smith T, et al. Cancer cachexia: traditional therapies and novel molecular mechanism-based approaches to treatment. *Curr Treat Options Oncol* 2010;11:107–117.
41. Laviano A, Meguid MM, Inui A, et al. Therapy insight: cancer anorexia-cachexia syndrome—when all you can eat is yourself. *Nat Clin Pract Oncol* 2005;2:158–165.
42. Oberholzer R, Hopkinson JB, Baumann K, et al. Psychosocial effects of cancer cachexia: a systematic literature search and qualitative analysis. *J Pain Symptom Manage* 2013;46:77–95.
43. Martin L, Birdsall L, MacDonald N, et al. Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index. *J Clin Oncol* 2013;31:1539–1547.
44. Dy SM, Lorenz KA, Naeim A, et al. Evidence-based recommendations for cancer fatigue, anorexia, depression, and dyspnea. *J Clin Oncol* 2008;26: 3886–3895.
45. Dy SM, Apostol CC. Evidence-based approaches to other symptoms in advanced cancer. *Cancer J* 2010;16:507–513.
46. Mercadante S, Aielli F, Adile C, et al. Prevalence of oral mucositis, dry mouth, and dysphagia in advanced cancer patients. *Support Care Cancer* 2015;23:3249–3255.
47. Roeland EJ, Bohlke K, Baracos VE, et al. Management of cancer cachexia: ASCO guideline. *J Clin Oncol* 2020;38:2438–2453.
48. Pascual López A, Roqué i Figuls M, Urrutia Cuchi G, et al. Systematic review of megestrol acetate in the treatment of anorexia-cachexia syndrome. *J Pain Symptom Manage* 2004;27:360–369.
49. Ruiz Garcia V, López-Briz E, Carbonell Sanchis R, et al. Megestrol acetate for treatment of anorexia-cachexia syndrome. *Cochrane Database Syst Rev* 2013;3:CD004310.
50. Yavuzsen T, Davis MP, Walsh D, et al. Systematic review of the treatment of cancer-associated anorexia and weight loss. *J Clin Oncol* 2005;23:8500–8511.
51. Miller S, McNutt L, McCann MA, et al. Use of corticosteroids for anorexia in palliative medicine: a systematic review. *J Palliat Med* 2014;17:482–485.
52. Navari RM, Brenner MC. Treatment of cancer-related anorexia with olanzapine and megestrol acetate: a randomized trial. *Support Care Cancer* 2010;18:951–956.
53. By the 2019 American Geriatrics Society Beers Criteria Update Expert Panel. American Geriatrics Society 2019 Updated AGS Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc* 2019;67:674–694.
54. Bolen JC, Andersen RE, Bennett RG. Deep vein thrombosis as a complication of megestrol acetate therapy among nursing home residents. *J Am Med Dir Assoc* 2000;1:248–252.
55. Ruiz Garcia V, López-Briz E, Carbonell Sanchis R, et al. Megestrol acetate for treatment of anorexia-cachexia syndrome. *Cochrane Database Syst Rev* 2013;2013:CD004310.