

Supplementary Table S1. Overview of Geriatric Assessment Measures by Domain

GA Domain	Patient-Reported Measures
FUNCTION	OARS Instrumental Activities of Daily Living (IADL) ¹ OARS Activities of Daily Living (ADL) ¹ Patient-reported ECOG Performance Status ² No. of falls in last 6 months ³
NUTRITION	Patient-Generated Subjective Global Assessment ^{2,4}
COGNITION	PROMIS Cognitive Function ⁵
PSYCHOLOGICAL	PROMIS Anxiety ^{6,7} PROMIS Depression ^{6,7}
SOCIAL SUPPORT	MOS Social Support Survey ⁸ (Emotional/Informational Support subscales)
COMORBIDITY	No. of medications ⁹ OARS comorbidity assessment ^{1,10} Vision and Hearing Assessment
HRQOL	PROMIS 10-item Global Health ¹¹

References

1. Fillenbaum GG, Smyer MA. The development, validity, and reliability of the OARS multidimensional functional assessment questionnaire. *J Gerontol* 1981;36:428–434.
2. Bauer J, Capra S, Ferguson M. Use of the scored Patient-Generated Subjective Global Assessment (PG-SGA) as a nutrition assessment tool in patients with cancer. *Eur J Clin Nutr* 2002;56:779–785.
3. Teno J, Kiel D, Mor V. Multiple stumbles: a risk factor for falls in community-dwelling elderly. A prospective study. *J Am Geriatr Soc* 1990;38:1321–1325.

4. Viganò AL, di Tomasso J, Kilgour RD, et al. The abridged patient-generated subjective global assessment is a useful tool for early detection and characterization of cancer cachexia. *J Acad Nutr Diet* 2014;114:1088–1098.
5. Saffer BY, Lanting SC, Koehle MS, et al. Assessing cognitive impairment using PROMIS® applied cognition-abilities scales in a medical outpatient sample. *Psychiatry Res* 2015;226:169–172.
6. Pilkonis PA, Choi SW, Reise SP, et al. Item banks for measuring emotional distress from the Patient-Reported Outcomes Measurement Information System (PROMIS(R)): depression, anxiety, and anger. *Assessment* 2011;18:263–283.
7. Riley WT, Pilkonis P, Cella D. Application of the National Institutes of Health Patient-reported Outcome Measurement Information System (PROMIS) to mental health research. *J Ment Health Policy Econ* 2011;14:201–208.
8. Moser A, Stuck AE, Silliman RA, et al. The eight-item modified Medical Outcomes Study Social Support Survey: psychometric evaluation showed excellent performance. *J Clin Epidemiol* 2012;65:1107–1116.
9. Lees J, Chan A. Polypharmacy in elderly patients with cancer: clinical implications and management. *Lancet Oncol* 2011;12:1249–1257.
10. Klepin HD, Pitcher BN, Ballman KV, et al. Comorbidity, chemotherapy toxicity, and outcomes among older women receiving adjuvant chemotherapy for breast cancer on a clinical trial: CALGB 49907 and CALGB 361004 (Alliance). *J Oncol Pract* 2014;10:e285–292.
11. Hays RD, Bjorner JB, Revicki DA, et al. Development of physical and mental health summary scores from the patient-reported outcomes measurement information system (PROMIS) global items. *Qual Life Res* 2009;18:873–880.